

Eludril

Solution

Formula :

Chlorhexidine (DCI) digluconate.....	0,10 g
Chlorobutanol (DCI).....	0,10 g
Sodium Docusate (DCI).....	0,10 g
Chloroform.....	0,5 ml
Excipient q.s.....	100 ml
(alcohol at 95° , glycerin, menthol, mint spirit, methyl parahydroxybenzoate, propyl parahydroxybenzoate, coccine, purified water).	

Therapeutic Class : Local Antiseptic, Local Anesthetic (A : Digestive tract and metabolism).

Indications : • Tonsillitis • Pre and post-operative care in ENT and stomatology
• Gingivitis, stomatitis, mouth ulcer, parodontitis.

Contraindication : Do not use in infants.

SHOULD THERE BE ANY DOUBT, IT IS ABSOLUTELY NECESSARY TO ASK YOUR PHYSICIAN'S ADVICE.

Precautions : KEEP OUT OF REACH OF CHILDREN.

Eludril contains alcohol. Take care medication is not swallowed when administered to children.

Medical interactions : To avoid medical interactions, other treatments have to be indicated to the physician or pharmacist.

Dosage : Gargle or use as mouthwash 2 to 3 times daily or more depending on your physician's advice. Mean dosage = 15 ml of Eludril to dilute with water.

Administration: Dilute 2-4 coffee spoonfuls of Eludril solution in half a glass of water.

Side effects : Embarrassing effects have to be indicated to the physician.

Duration of stability : Do not exceed the use by date stated on the packaging.

* Store 15° - 25° C away from light & humidity .

Reg. N° Lebanon : N° 38908/96

Presentation : Bottle of 90 ml.

Pharmaline

905700-A

Produced by Pharmaline - Lebanon
Licensed by Pierre Fabre - France



Pierre Fabre