

Febradol

Brand of Paracetamol, BP

Analgesic / Antipyretic

DESCRIPTION Febradol, brand of paracetamol, is an analgesic and antipyretic agent.

ACTIONS Febradol produces analgesia through its selective depressant action on the pain-perception apparatus in the central nervous system. Reduction of fever results by its direct effect on the hypothalamic heat-regulating center to produce peripheral vasodilation resulting in increased blood flow through the skin, sweating and heat loss. It is metabolized in the liver and excreted in the urine. Protein binding is 25%.

INDICATIONS Febradol is indicated for the relief of painful disorders such as simple headaches, minor muscular aches, tooth extractions, dental work, teething and menstrual cramps. It is also indicated in conditions accompanied by discomfort and fever such as common colds and viral infections.

CONTRAINDICATIONS Febradol is contraindicated in patients with known hypersensitivity to the drug.

PRECAUTIONS If pain persists for more than 10 days or fever remains for more than 3 days, consult a physician immediately.

ADVERSEREACTIONS Skin reactions such as urticaria are rarely reported.

DOSAGE & ADMINISTRATION

ADULTS

500-1000 mg, 3-4 times daily as needed.

CHILDREN

AGE

Below 1 Yr. :	60 mg	3-4 times daily as needed
1-3 Yrs:	120 mg	3-4 times daily as needed
4-6 Yrs:	240 mg	3-4 times daily as needed
7- 12 Yrs:	250-500 mg	3-4 times daily as needed

AVAILABILITY

* Easy to Swallow Tablets: Packs of 20 tablets, each containing Paracetamol 500 mg , Excipient q.s.1 tablet. Reg N° Lebanon: 208322

* Syrup Alcohol & Sugar free : Bottles of 60 ml containing in each 5 ml (teaspoonful) Paracetamol 120 mg, Excipient q.s. 5 ml. Reg N° Lebanon: 28555

* Elixir: Bottles of 100 ml containing in each 5 ml (teaspoonful) Paracetamol 120 mg, Excipient q.s. 5 ml. Reg N° Lebanon: 26596

This is a medicament

-A medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.

-Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.

-The doctor and the pharmacist are experts in medicine, its benefits and risks.

-Do not by yourself interrupt the period of treatment prescribed for you.

-Do not repeat the same prescription without consulting your doctor.

Keep medicaments out of reach of children

Pharmaline -Lebanon

Council of Arab Health Ministers Union of Arab Pharmacists

Febradol is a Registered Trademark.

REV. 11/09 905261-B