

HÉLICIDINE[®]

SYRUP 10 per cent

QUALITATIVE AND QUANTITATIVE COMPOSITION

HELICIDINE 10 ml.

per 100 ml of syrup.

Excipients: saccharose, glycerol, raspberry flavoured, parahydroxy methyl benzoate, sodic parahydroxypropyl benzoate, purified water.

1 teaspoon contains 3.4 grams of saccharose.

1 tablespoon contains 10.3 grams of saccharose.

PHARMACEUTICAL FORM

Syrup, bottle of 125 ml or 250 ml.

PHARMACO-THERAPEUTIC CLASS

Peripheral acting anti-tussive

(R: respiratory system)

CASES IN WHICH THIS MEDICINE CAN BE USED

This medicine is recommended for the relief of dry and irritating coughs.

CASES IN WHICH THIS MEDICINE SHOULD NOT BE USED

Allergy to one of these constituents.

IF IN DOUBT, SEEK ADVICE FROM YOUR DOCTOR OR PHARMACIST.

SPECIAL WARNING

- Do not use this medicine for the relief of loose coughs. In this case, coughing is a natural means of defence necessary for the evacuation of bronchial secretions.
- If the cough becomes loose causing blockage, coughing up of phlegm or fever, consult your doctor.
- In the event of chronic disorders (after a period of time) of the bronchial tract or the lungs with coughing up of phlegm, you must seek medical advice.

PRECAUTIONS WHEN USING

In the event of diabetes or low sugar diet, bear in mind the contribution of 3.4 grams of saccharose per teaspoon and of 10.3 grams per tablespoon.

IF IN DOUBT, DO NOT HESITATE TO SEEK THE ADVICE OF YOUR DOCTOR OR PHARMACIST.

MEDICINAL AND OTHER INTERACTIONS

IN ORDER TO AVOID ANY POSSIBLE INTERACTIONS BETWEEN SEVERAL MEDICINES, YOU MUST SYSTEMATICALLY INFORM YOUR DOCTOR OR PHARMACIST OF ANY OTHER TREATMENT YOU MAY BE FOLLOWING.

LIST OF EXCIPIENTS THE KNOWLEDGE OF WHICH IS NECESSARY FOR A RISK-FREE USE BY CERTAIN PATIENTS

- Saccharose
- Parahydroxy methyl benzoate
- Sodic parahydroxy propyl benzoate
- Glycerol

HOW TO USE THIS MEDICINE

Dosage

If no improvement is observed, do not increase the dose beyond that recommended and do not take any other cough mixture. Consult your doctor.

Usual dose is:

Adults: 2 tablespoons 3 times a day.

Children from 25 to 50 Kg (about 8 to 15 years of age): 3 to 5 tablespoons per day.

Children from 15 to 25 Kg (about 4 to 8 years of age): 1 tablespoon 3 times a day.

Children under 4 years of age: Follow your doctor's advice.

THIS MEDICINE IS PROVIDED FOR YOUR PERSONAL USE FOR A SPECIFIC SITUATION:

- IT MAY BE NOT SUITABLE FOR ANOTHER CASE.

- DO NOT RECOMMEND IT TO ANOTHER PERSON.

METHOD AND ROUTE OF ADMINISTRATION

Orally

DURATION OF TREATMENT

Do not exceed 5 to 6 days treatment without medical advice.

UNPLEASANT AND UNDESIRABLE EFFECTS

LIKE ALL ACTIVE PRODUCTS, THIS MEDICINE CAN CAUSE UNWANTED EFFECTS IN SOME PERSONS.

Possibility of allergic reactions.

DO NOT HESITATE TO SEEK THE ADVICE OF YOUR DOCTOR OR PHARMACIST AND MENTION ANY UNWANTED OR SIDE EFFECTS NOT INDICATED IN THIS PACKAGE INSERT.

STORAGE: DO NOT USE AFTER THE EXPIRY DATE INDICATED ON THE OUTER PACKAGING.

INFORM YOUR PHARMACIST OF ANY VISIBLE SIGN OF DETERIORATION.

KEEP OUT OF REACH OF CHILDREN

Name and address of producer

THERABEL LUCIEN PHARMA

123, rue Jules Guesde

92309 LEVALLOIS-PERRET cedex

France

DATE OF REVISION OF INSERT

11/97

125 ml Authorized medicine n° 304 765-7

250 ml Authorized medicine n° 340 316-4

Manufacturer :

Laboratoires THISSEN

B-1420 Braine-l'Alleud

