

Use in the paediatric population

For paediatric patients Bridion can be diluted using sodium chloride 9 mg/mL (0.9%) to a concentration of 10 mg/mL (see section 6.3).

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. MANUFACTURER

Patheon Manufacturing Services LLC,
5900 Martin Luther King Jr. Highway, Greenville, North
Carolina 27834, USA

8. MARKETING AUTHORISATION HOLDER

Merck Sharp & Dohme Limited Hertford Road, Hoddesdon Hertfordshire EN11 9BU
United Kingdom

9. DATE OF REVISION OF THE TEXT

This leaflet was last approved in April 2020

Detailed information on this medicinal product is available on the website of the European Medicines Agency
<http://www.ema.europa.eu>.

Prescription Only Medicine

THIS IS A MEDICAMENT

- Medicament is a product which affects your health and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament. The doctor and the pharmacist are the experts in medicines, their benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed.
- Do not repeat the same prescription without consulting your doctor.
- Keep all medicaments out of reach of children.

Council of Arab Health Ministers & Union of Arab Pharmacists