

Fortum™

Ceftazidime

QUALITATIVE AND QUANTITATIVE COMPOSITION

FORTUM injection contains 250 mg, 500 mg, 1 g, 2 g or 3 g of ceftazidime (as pentahydrate).
FORTUM MONOVIAL™ contains 1 g or 2 g of ceftazidime (as pentahydrate).

PHARMACEUTICAL FORM

Powder for injection/infusion

CLINICAL PARTICULARS

Indications

Treatment of single or multiple infections caused by susceptible organisms.
May be used alone as first choice drug before the results of sensitivity tests are available.
May be used in combination with an aminoglycoside or most other beta-lactam antibiotics.
May be used with an antibiotic against anaerobes when the presence of *Bacteroides fragilis* is suspected.
Indications include:

- severe infections e.g.
 - septicaemia, bacteraemia, peritonitis, meningitis.
 - infections in immunosuppressed patients
 - infections in patients in intensive care, e.g. infected burns

- respiratory tract infections including lung infections in cystic fibrosis
- ear, nose and throat infections
- urinary tract infections
- skin and soft tissue infections
- gastrointestinal, biliary and abdominal infections
- bone and joint infections
- infections associated with haemo- and peritoneal dialysis and with continuous ambulatory peritoneal dialysis (CAPD)
- prophylaxis: prostatic surgery (transurethral resection).

Dosage and Administration

Dosage depends upon the severity, sensitivity, site and type of infection and upon the age and renal function of the patient. Use **FORTUM** injection i.v. or by deep i.m. injection. Recommended i.m. injection sites are the upper outer quadrant of the gluteus maximus or lateral part of the thigh.

FORTUM solutions may be given directly into the vein or introduced into the tubing of a giving set if the patient is receiving parenteral fluids.

FORTUM MONOVIAL is for i.v. infusion only.

• Adults

1 to 6 g/day in two or three divided doses by i.v. or i.m. injection.

Urinary tract and less severe infections:

- 500 mg or 1 g every 12 h.

Most infections:

- 1 g every 8 h or 2 g every 12 h.

Very severe infections particularly in immunocompromised patients including those with neutropenia:

- 2 g every eight or 12 h, or 3 g every 12 h.

Fibrocystic adults with pseudomonal lung infections:

- 100 to 150 mg/kg/day in three divided doses.

In adults with normal renal function 9 g/day has been used without ill effect.

When used as a prophylactic agent in prostatic surgery, 1 g should be given at the induction of anaesthesia. A second dose should be considered at the time of catheter removal.

• Infants and children (greater than 2 months)

30 to 100 mg/kg/day in two or three divided doses.

Doses up to 150 mg/kg/day (maximum 6 g/day) in three divided doses may be given to infected immunocompromised or fibrocystic children or children with meningitis.

• Neonates (0 to 2 months)

25 to 60 mg/kg/day in two divided doses.

In neonates, the serum half life of ceftazidime can be three to four times that in adults.

• Elderly

In view of the reduced clearance of ceftazidime in acutely ill elderly patients, the daily dosage should not normally exceed 3 g, especially in those over 80 years of age.

• Renal Impairment

Ceftazidime is excreted unchanged by the kidneys. Therefore, in patients with impaired renal function, the dosage should be reduced. An initial loading dose of 1 g should be given. Maintenance doses should be based on creatinine clearance:

Recommended maintenance doses of **FORTUM** in renal insufficiency:

Creatinine Clearance (ml / min)	Approx. Serum creatinine (micromoles / l) (mg / dl)	Recommended unit dose of FORTUM (g)	Frequency of dosing (hourly)
> 50	< 150 (<1.7)	Normal dosage	
50 – 31	150 – 200 (1.7 – 2.3)	1.0	12
30 – 16	200 – 350 (2.3 – 4.0)	1.0	24
15 – 6	350 – 500 (4.0 – 5.6)	0.5	24
< 5	> 500 (> 5.6)	0.5	48

In patients with severe infections the unit dose should be increased by 50% or the dosing frequency increased. In such patients the ceftazidime serum levels should be monitored and trough levels should not exceed 40 mg/l.

In children the creatinine clearance should be adjusted for body surface area or lean body mass.

Haemodialysis

The serum half-life during haemodialysis ranges from 3 to 5 h.

Following each haemodialysis period, the maintenance dose of **FORTUM** recommended in the above table should be repeated.

Peritoneal dialysis

FORTUM may be used in peritoneal dialysis and continuous ambulatory peritoneal dialysis (CAPD).

In addition to i.v. use, **FORTUM** can be incorporated into the dialysis fluid (usually 125 to 250 mg for 2 litres of dialysis solution).

For patients in renal failure on continuous arteriovenous haemodialysis or high-flux haemofiltration in intensive therapy units; 1 g daily either as a single dose or in divided doses. For low-flux haemofiltration, follow the dosage recommended under impaired renal function.

For patients on venovenous haemofiltration and venovenous haemodialysis, follow the dosage recommendations in the tables below.

Continuous venovenous haemofiltration dosage guidelines for **FORTUM**

Residual renal function (creatinine clearance in ml/min)	Maintenance dose (mg) for a ultrafiltration rate (ml/min) of \geq :				
	5	16.7	33.3	50	
0	250	250	500	500	
5	250	250	500	500	500
10	250	500	500	500	750
15	250	500	500	500	750
20	500	500	500	500	750

^a Maintenance dose to be administered every 12 h.

FORTUM dosage guidelines during continuous venovenous haemodialysis

Residual renal function (creatinine clearance in ml/min)	Maintenance dose (mg) for a dialysate in flow rate of \geq :					
	1.0 litre/h			2.0 litres/h		
	Ultrafiltration rate (litre/h)			Ultrafiltration rate (litres/h)		
	0.5	1.0	2.0	0.5	1.0	2.0
0	500	500	500	500	500	750
5	500	500	750	500	500	750
10	500	500	750	500	750	1000
15	500	750	750	750	750	1000
20	750	750	1000	750	750	1000

^a Maintenance dose to be administered every 12 h.

Contraindications

- Patients with known hypersensitivity to cephalosporin antibiotics.
- Hypersensitivity to ceftazidime pentahydrate or to any of the excipients of the injection.

Warnings and Precautions

Before beginning treatment establish whether the patient has a history of hypersensitivity reactions to ceftazidime, cephalosporins, penicillins or other drugs.

Special care is indicated in patients who have experienced an allergic reaction to penicillins or other beta-lactams.

If an allergic reaction to **FORTUM** occurs discontinue the drug. Serious hypersensitivity reactions may require epinephrine (adrenaline), hydrocortisone, antihistamine or other emergency measures.

Concurrent treatment with high doses of cephalosporins and nephrotoxic drugs such as aminoglycosides or potent diuretics (e.g. frusemide) may adversely affect renal function. Clinical experience has shown that this is not likely to be a problem with **FORTUM** at the recommended dose levels. There is no evidence that **FORTUM** adversely affects renal function at normal therapeutic doses.

Ceftazidime is eliminated via the kidneys, therefore the dosage should be reduced according to the degree of renal impairment. Neurological sequelae have occasionally been reported when the dose has not been reduced in patients with renal impairment (see *Dosage and Administration – Renal Impairment and Adverse Reactions*).

As with other broad spectrum antibiotics, prolonged use may result in the overgrowth of non-susceptible organisms (e.g. *Candida*, enterococci) which may require interruption of treatment or appropriate measures. Repeated evaluation of the patient's condition is essential.

As with other extended-spectrum cephalosporins and penicillins, some initially susceptible strains of *Enterobacter* spp. and *Serratia* spp. may develop resistance during **FORTUM** therapy. When clinically appropriate during therapy of such infections, periodic susceptibility testing should be considered.

Interactions

Concurrent use of high doses with nephrotoxic drugs may adversely affect renal function (see *Warnings and Precautions*).

Chloramphenicol is antagonistic in vitro with ceftazidime and other cephalosporins. The clinical relevance of this finding is unknown, but if concurrent administration of **FORTUM** with chloramphenicol is proposed, the possibility of antagonism should be considered.

In common with other antibiotics, ceftazidime may affect the gut flora, leading to lower oestrogen reabsorption and reduced efficacy of combined oral contraceptives.

Ceftazidime does not interfere with enzyme-based tests for glycosuria but slight interference may occur with copper reduction methods (Benedict's, Fehling's, Clinitest).

Ceftazidime does not interfere in the alkaline picrate assay for creatinine.

Pregnancy and Lactation

There is no experimental evidence of embryopathic or teratogenic effects, but as with all drugs, **FORTUM** should be administered with caution during the early months of pregnancy and early infancy.

Ceftazidime is excreted in human milk in small quantities and should be used with caution in breast feeding.

Effects on Ability to Drive and Use Machines

None reported.

Adverse Reactions

Data from large clinical trials (internal and published) were used to determine the frequency of very common to uncommon undesirable effects. The frequencies assigned to all other undesirable effects were mainly determined using post-marketing data and refer to a reporting rate rather than a true frequency.

The following convention has been used for the classification of frequency:

very common \geq 1/10,

common \geq 1/100 and <1/10,

uncommon \geq 1/1,000 and <1/100,

rare \geq 1/10,000 and <1/1,000,

very rare <1/10,000.

Infections and infestations

Uncommon: Candidiasis (including vaginitis and oral thrush).

Blood and lymphatic system disorders

Common: Eosinophilia and thrombocytosis.

Uncommon: Leucopenia, neutropenia, and thrombocytopenia.

Very rare: Lymphocytosis, haemolytic anaemia, and agranulocytosis.

Immune system disorders

Very rare: Anaphylaxis (including bronchospasm and/or hypotension).

Nervous system disorders

Uncommon: Headache and dizziness.

Very rare: Paraesthesia.

There have been reports of neurological sequelae including tremor, myoclonia, convulsions, encephalopathy, and coma in patients with renal impairment in whom the dose of **FORTUM** has not been appropriately reduced.

Vascular disorders

Common: Phlebitis or thrombophlebitis with i.v. administration.

Gastrointestinal disorders

Common: Diarrhoea.

Uncommon: Nausea, vomiting, abdominal pain, and colitis.

Very rare: Bad taste.

As with other cephalosporins, colitis may be associated with *Clostridium difficile* and may present as pseudomembranous colitis.

Hepatobiliary disorders

Common: Transient elevations in one or more of the hepatic enzymes, ALT (SGPT), AST (SOGT), LDH, GGT and alkaline phosphatase.

Very rare: Jaundice.

Skin and subcutaneous tissue disorders

Common: Maculopapular or urticarial rash.

Uncommon: Pruritus.

Very rare: Angioedema, erythema multiforme, Stevens-Johnson syndrome, and toxic epidermal necrolysis.

General disorders and administration site conditions

Common: Pain and/or inflammation after i.m. injection.

Uncommon: Fever.

Investigations

Common: Positive Coombs test.

Uncommon: As with some other cephalosporins, transient elevations of blood urea, blood urea nitrogen and/or serum creatinine have been observed.

A positive Coombs test develops in about 5% of patients and may interfere with blood cross-matching.

Overdose

Overdosage can lead to neurological sequelae including encephalopathy, convulsions and coma.

Serum levels of ceftazidime can be reduced by haemodialysis or peritoneal dialysis

PHARMACOLOGICAL PROPERTIES

Pharmacodynamics

Mechanism of Action

Ceftazidime is bactericidal in action. It acts by inhibiting bacterial cell wall synthesis.

Pharmacodynamic Effects

Bacteriology

A wide range of pathogenic strains and isolates are susceptible *in vitro* including strains resistant to gentamicin and other aminoglycosides. Ceftazidime is highly stable to most clinically important beta-lactamases produced by both Gram-positive and Gram-negative organisms, therefore it is active against many ampicillin- and cephalothin-resistant strains. Ceftazidime has high intrinsic activity *in vitro* and acts within a narrow MIC range for most genera with minimal changes in MIC at varied inoculum levels. *In vitro* the activities of ceftazidime and aminoglycosides in combination are additive. There is evidence of synergy in some strains.

Ceftazidime is active *in vitro* against the following organisms:

Gram-negative:

Pseudomonas aeruginosa

Pseudomonas spp (including *Ps. pseudomallei*)

Escherichia coli

Klebsiella spp. (including *Klebsiella pneumoniae*)

Proteus mirabilis

Proteus vulgaris

Morganela morgani (formerly *Proteus morgani*)

Proteus rettgeri

Providencia spp.

Enterobacter spp.

Citrobacter spp.

Serratia spp.

Salmonella spp.

Shigella spp.

Yersinia enterocolitica

Pasteurella multocida

Acinetobacter spp.

Neisseria gonorrhoeae

Neisseria meningitidis

Haemophilus influenzae (including ampicillin resistant strains)

Haemophilus parainfluenzae (including ampicillin resistant strains).

Gram-positive:

Staphylococcus aureus (methicillin-sensitive strains)

Staphylococcus epidermidis (methicillin-sensitive strains)

Micrococcus spp.

Streptococcus pyogenes (Group A beta-haemolytic streptococci)

Streptococcus Group B (*S. agalactiae*)

Streptococcus pneumoniae

Streptococcus mitis

Streptococcus spp (excluding *Enterococcus (Streptococcus faecalis)*)

Anaerobic strains:

Peptococcus spp.

Peptostreptococcus spp.

Streptococcus spp.

Propionibacterium spp.

Clostridium perfringens

Fusobacterium spp.

Bacteroides spp (many strains of *Bacteroides fragilis* resistant).

Ceftazidime is not active *in vitro* against the following organisms:

Methicillin-resistant staphylococci.

Enterococcus (Streptococcus) faecalis and many other enterococci.

Clostridium difficile

Listeria monocytogenes

Campylobacter spp.

Pharmacokinetics

Absorption

After i.m. administration of 500 mg and 1 g, peak levels of 18 and 37 mg/l, respectively, are achieved rapidly. Five minutes after i.v. bolus injection of 500 mg, 1 g or 2 g, serum levels are, respectively, 46, 87 and 170 mg/l.

Distribution

Therapeutically effective concentrations are still present in the serum 8 to 12 h after either i.v. or i.m. administration. Serum protein binding is about 10%. Concentrations in excess of the MIC for common pathogens can be achieved in tissues such as bone, heart, bile, sputum, aqueous humour, synovial, pleural and peritoneal fluids. Ceftazidime crosses the placenta readily, and is excreted in the breast milk. Penetration of the intact blood-brain barrier is poor resulting in low levels of ceftazidime in the CSF in the absence of inflammation. However, therapeutic levels of 4 to 20 mg/l or more are achieved in the CSF when the meninges are inflamed.

Metabolism

Ceftazidime is not metabolised in the body.

Elimination

Parenteral administration produces high and prolonged serum levels, which decrease with a half-life of about 2 h.

Ceftazidime is excreted unchanged, in active form into the urine by glomerular filtration; approximately 80 to 90% of the dose is recovered in the urine within 24 h. Less than 1% is excreted via the bile, which limits the amount entering the bowel.

Special Patient Populations

Elimination of ceftazidime is decreased in patients with impaired renal function and the dose should be reduced. (See *Dosage and Administration – Renal Impairment, Warnings and Precautions*).

Pre-clinical Safety Data

No additional data of relevance.

PHARMACEUTICAL PARTICULARS

List of Excipients

Sodium carbonate (anhydrous).

Incompatibilities

FORTUM is less stable in Sodium Bicarbonate Injection than in other i.v. fluids. It is not recommended as a diluent. **FORTUM** and aminoglycosides should not be mixed in the same giving set or syringe. Precipitation has been reported with vancomycin added to **FORTUM** in solution. Therefore, it would be prudent to flush giving sets and i.v. lines between administration of these two agents.

Shelf Life

The expiry date is indicated on the packaging.

Special Precautions for Storage

Ceftazidime vials for injection : 3 years below 25°C. Ceftazidime Monovial: 2 years below 30°C.

Occasional storage at temperatures not higher than 30°C for up to two months is not detrimental to the product.

Protect unconstituted vials from light.

Nature and Contents of Container

As registered locally.

Instructions for Use/Handling

FORTUM for injection/infusion is compatible with most commonly used i.v. fluids. However, Sodium Bicarbonate Injection is not recommended as a diluent (see *Incompatibilities*).

All sizes of vials of Fortum Injection and Monovial are supplied under reduced pressure. As the product dissolves, carbon dioxide is released and a positive pressure develops. Small bubbles of carbon dioxide in the constituted solution may be ignored.

Vial Size	Amount of Diluent to be added (ml)	Approximate Concentration (mg/ml)	
250 mg	Intramuscular	1.0 ml	210
	Intravenous	2.5 ml	90
500 mg	Intramuscular	1.5 ml	260
	Intravenous	5 ml	90
1 g	Intramuscular	3 ml	260
	Intravenous bolus	10 ml	90
	Intravenous infusion	50 ml #	20