

Warnings:

Using more than one enema in 24 hours can be harmful.

Ask a doctor before using this product if the child:

- Is on a sodium restricted diet
- Has a kidney disease

Ask a doctor before

using any laxative if the child:

- Has nausea, vomiting, or abdominal pain
- Has a sudden change in bowel habits lasting more than 2 weeks
- Has already used a laxative for more than 1 week

Stop using this product

and consult a doctor if the child:

- Has rectal bleeding
- Has no bowel movement after the enema is given

These symptoms may indicate a serious condition.

Keep out of reach of children.

In case of overdose or accidental ingestion, get medical help right away.

Professional Use Warnings:

Do not use in patients with congenital megacolon, bowel obstruction, imperforate anus, or congestive heart failure. Use with caution in patients with impaired renal function, pre-existing electrolyte disturbances or a colostomy, or in patients on diuretics or other medications that may affect electrolyte levels. Hypocalcemia, hyperphosphatemia, hypernatremia, or acidosis may occur.

Consult **PHYSICIANS' DESK REFERENCE®** for complete professional labeling.

CARTON SEALED FOR SAFETY. IF SEAL WITH FLEET EMBLEM ON TOP OR BOTTOM FLAP IS BROKEN OR MISSING, DO NOT USE.

© C.B. Fleet Company, Inc. 1999
The package design is a trademark of C.B. Fleet Company, Inc.



C.B. FLEET CO., INC.
Lynchburg, Virginia
24502 U.S.A.

BRAND

pharmaceuticals

Fleet® enema for children

ready-to-use
saline laxative



Soft, Flexible
Comfortip®

Sample

LATEX FREE

2.25 Fl Oz (66 mL)

Uses:

For relief of occasional constipation or bowel cleansing before rectal examinations.

Active Ingredients:

(Each 59 mL delivered dose)

Active Ingredient	Purpose
Monobasic Sodium Phosphate 9.5 g	laxative/bowel cleanser
Dibasic Sodium Phosphate 3.5 g	laxative/bowel cleanser

Sodium Content 2.2 g


Single Daily Dosage:

Do not use more unless directed by a doctor.
See Warnings.

Children 5 to 11 years	One Bottle or as directed by a doctor.
Children 2 to under 5 years	One-Half Bottle (See Below)
Children under 2 years	DO NOT USE

One-half bottle preparation: Unscrew cap and remove 2 tablespoons of liquid with a measuring spoon. Replace cap and follow DIRECTIONS on back of carton.

Complete enema in a disposable squeeze bottle with soft, pre-lubricated Comfortip®



Fleet® Enemas are LATEX FREE.
Allergic reaction to materials containing Latex is a growing and serious medical problem.

DIRECTIONS

Positions for using this enema:



- **Left-side position:** Place child on left side with knees bent, and arms resting comfortably.



- **Knee-chest position:** Have child kneel, then lower head and chest forward until left side of face is resting on surface with left arm folded comfortably.

How to use this enema:

- **REMOVE ORANGE PROTECTIVE SHIELD FROM ENEMA COMFORTIP® BEFORE INSERTING.**



- With steady pressure, gently insert enema tip into rectum with a slight side-to-side movement, with tip pointing toward navel. Insertion may be easier if child receiving enema bears down, as if having a bowel movement. This helps relax the muscles around the anus.
- **Do not force the enema tip into rectum as this can cause injury.**
- Squeeze bottle until nearly all liquid is gone. It is not necessary to empty the bottle completely, as it contains more liquid than needed.
- Remove Comfortip® from rectum and keep child in position until urge to evacuate is strong (usually 2 to 5 minutes).

