

WARNINGS (cont.):

DO NOT USE:

- if you have congestive heart failure
- if you have serious kidney problems
- in children under 5 years of age

Ask a doctor before use if you are:

- under a doctor's care for any medical condition
- pregnant or breast-feeding
- on a low salt diet

Ask a doctor or pharmacist before use if you are taking any other prescription or non-prescription drugs.

Ask a doctor before using any laxative if you have:

- abdominal (belly) pain, nausea or vomiting
- a change in your daily bowel movements that lasts more than 2 weeks
- already used another laxative daily for constipation for more than 1 week

Stop use and ask a doctor if you:

- have any rectal bleeding
- do not have a bowel movement within 6 hours of taking this product
- have any symptoms that your body is losing more fluids than you are drinking. This is called dehydration.

Early symptoms of dehydration include:

- feeling thirsty
- dizziness
- urinating less often than normal
- vomiting

These symptoms may be signs of serious problems.


Fleet®

PHOSPHO-SODA®

**oral saline
laxative**

**Ginger-lemon
flavor**

**For relief of
occasional
constipation
or bowel
cleansing
before
medical
procedures**



45 mL

Use as a bowel cleanser before Medical Procedures

Use this product for bowel cleansing only as directed by your doctor. Before using this product for bowel cleansing, make sure you have talked to your doctor about your health and warnings about use of this product as a bowel cleanser.

Follow the special directions from your doctor exactly. Take the dose your doctor has recommended. Do NOT take the laxative dose in the chart. Drink as much extra clear liquids as you can to replace fluids you are losing during bowel movements: minimum 9 (240 ml) glasses.

ACTIVE INGREDIENTS: (in each 15 ml)

Active Ingredient	Purpose
Monobasic Sodium Phosphate 7.2g	saline laxative/ bowel cleanser
Dibasic Sodium Phosphate 2.7g	saline laxative/ bowel cleanser

INACTIVE INGREDIENTS:
flavor, glycerin, purified water, saccharin sodium, sodium benzoate

WARNINGS:

Taking more than the recommended dose in 24 hours can be harmful.
(cont. on side panel)

DIRECTIONS:

Use as a laxative for relief of occasional constipation. Follow the dose and directions below.

Do not take more unless directed by a doctor. See Warnings. Drink as much extra liquids as you can to help replace the fluids you are losing during bowel movements.

LAXATIVE DOSE AND DIRECTIONS

Ages (years)	Step 1	Step 2	24 Hour Maximum Dose*
12 Years & Older	Mix 15 ml (1 tablespoon) in a full glass of cold liquid (240 ml). Drink	Drink at least 1 extra full glass of liquid (240 ml).	45 ml (3 tablespoons)
10 to 11 years			15 ml (1 tablespoon)
5 to 9 years	Mix 7.5 ml (½ tablespoon) in a full glass of cold liquid (240 ml). Drink	Drink at least 1 extra full glass of liquid (240 ml).	7.5 ml (½ tablespoon)
Under 5 years	DO NOT USE	DO NOT USE	DO NOT USE

*Take no more than this amount in a 24 hour period.

USES:

Relief of occasional constipation. To help clean out the bowel before medical procedures.

OTHER INFORMATION:

1 tablespoon contains: sodium 1668 mg. This product is sugar-free. This product usually causes a person to have a bowel movement in 30 minutes to 6 hours. Chill this product in refrigerator to improve the taste. Do not freeze.

The top of the bottle is sealed with foil for your safety. Do not use if foil imprinted "SEALED for YOUR PROTECTON" is broken or missing.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Made in the U.S.A. by Fleet Laboratories™

Fleet
Laboratories

© C.B. Fleet Company, Inc. 2006.

This package design is a trademark of C.B. FLEET CO., INC., Lynchburg, VA 24502 U.S.A.