Clinimycin® T Clindamycin phosphate

Clindamycin phosphate

Properties:

Clinimycin® T is a topical antibacterial, antiacne agent, characterized by minimal percutaneous absorption.

Indications:

Clinimycin® T is indicated in the treatment of acne vulgaris.

It is effective against inflammatory acne lesions such as papules and pustules.

Clinimycin®T is also indicated in the treatment of minor bacterial skin infections as well as dermal ulcers.

Dosage and administration:

The skin should be cleansed prior to application. A thin film of **Clinimycin®T** should be applied to the affected area twice daily. Avoiding contact with the eyes and mouth.

Precautions:

- Medication should be stopped if antibiotic associated pseudomembranous colitis develops during treatment.
- Risk benefit ratio of the medication should be considered in patients with history of antibiotic associated p. colitis, ulcerative colitis or regional enteritis
- Avoid contact with eyes, nose, mouth or other mucous membranes. In case of accidental contact with these areas, wash thoroughly with water.
- Avoid using near heat, open flame or while smoking.
- Pediatric use: Safety and effectiveness in children under the age of 12 years have not been established.
- Adequate and well-controlled studies in pregnant women have not been done.
- It is not known whether topical clindamycin is distributed into breast milk. Since only a small amount is absorbed following topical application of clindamycin phosphate, it is unlikely to be distributed into breast milk in significant amounts.

Contraindications:

Patients with known sensitivity to clindamycin or lincomycin.

Side effects:

It is well tolerated. It causes skin irritation and dryness or hypersensitivity. It may also cause diarrhea, nausea or abdominal pain and in rare cases pseudomembranous colitis

Storage:

Keep container tightly closed.

Presentations:

Clinimycin® T Topical Solution: Each 1 ·ml contains Clindamycin phosphate B.P equivalent to Clindamycin 10 mg in bottles of 50 ml.

Clinimycin® T Topical Gel: Each gram contains Clindamycin phosphate B.P equivalent to Clindamycin 10 mg in tubes of 45 g.

(This is a medicament - Keep medicaments out of the reach of children)

- Medicament is a product that affects your health, and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who dispensed the medicament.
- The doctor and the pharmacist are experts in medicine, its benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed
- Do not repeat the same prescription without consulting your doctor.



Hayat Pharmaceutical Industries Co. PLC P.O. Box 1564 Amman 11118 Jordan