

For the use of a Registered Medical Practitioner or a Hospital or a Laboratory only

# Uri Soda

*Systemic & urinary alkalinizer*



**Contains:**

Sodium Bicarbonate BP 44.03%,  
Sodium Citrate Anhydrous USP 15.75%,  
Tartaric Acid BP 22.25%,  
Citric Acid Anhydrous BP 17.88%,  
Flavoured Excipients Q.S.

**Direction for use:**

Pour entire contents of a sachet in about 3/4th of a glass of water. Drink after effervescence.

**Dosage :**

**Adults :** 1-2 sachets in 3/4th of a glass of water three times daily.

**Children :** 6-12 years of age :  
One sachet in half a glass  
of water two or three times daily.

**Indications:**

1) Urinary alkalinisation where indicated in the treatment of urinary tract infections like cystitis, pyelonephritis, urethritis.

- 2) Symptomatic relief of dysuria.
- 3) To enhance the action of certain antibiotics especially gentamycin, sulfonamides, kanamycin, streptomycin.
- 4) In gout especially when treated with uricosurics and possibly allopurinol.
- 5) Symptomatic treatment of gastric hyperacidity.
- 6) In higher doses exerts mild laxative action.

Store in a dry place at room temperature.

**Presentation**

**Uri Soda**                      Sachets of 4 gm.