

PERFU-MOL® 1 g/100 ml Infusion

Paracetamol

ACTION

This medicine is an analgesic and an antipyretic.

It is restricted to adults and children weighing more than 33 kg (approximately 11 years old).

INDICATIONS

It is indicated for the short-term treatment of moderate pain, especially following surgery and for the short-term treatment of fever.

DOSAGE AND ADMINISTRATION

Dolorgan 1g/100 ml, for Intravenous infusion administration.

Restricted to adults and children weighing more than 33 kg (approximately 11 years old).

Before administration, the product should be visually inspected for any particulate matter and yellowing.

For single use only, any unused solution should be discarded.

Dosage:

Adolescents and adults weighing more than 50 kg:

1g of Paracetamol per administration, i.e. a 100 ml vial, up to 4 times a day.

Leave an interval of at least 4 hours between administrations.

The maximum dose must not exceed 4 g of paracetamol per day, taking into account all the medicines containing paracetamol.

Children weighing more than 33 kg (approximately 11 years old), adolescents and adults weighing less than 50 kg:

15 mg/kg paracetamol per administration, i.e. 1.5 ml solution per kg.

Leave an interval of at least 4 hours between administrations.

The maximum dose must not exceed 60 mg paracetamol per kilo and per day (without exceeding 4 g), taking into account all the medicines containing paracetamol.

Method of administration

The Dolorgan solution is administered in intravenous infusion over 15 minutes.

In children the volume of solution to be infused is 1.5 ml per kg per administration.

If you have the impression that the effect of Dolorgan 1 g/100 ml, solution for infusion is too strong or too weak, talk to your doctor or pharmacist.

If you may have used more Dolorgan 1 g/100 ml, solution for infusion than you should, talk to your doctor or pharmacist immediately.

CONTRAINDICATIONS

DO NOT USE Paracetamol 1 g/100 ml, solution for infusion in the following cases:

- Known allergy to paracetamol, to propacetamol (another analgesic for infusion and a precursor of paracetamol) or to one of the constituents.
- Severe liver disease.

PRECAUTIONS

- Inform your doctor in cases of liver or kidney disease, or alcohol abuse.
- Inform your doctor if you are taking other medicines containing paracetamol.
- In cases of nutrition problems (malnutrition) or dehydration, paracetamol must be used with caution.

Pregnancy

Inform your doctor if you are pregnant. Under normal conditions of use, paracetamol may be used during pregnancy. However, in this case the doctor must evaluate if the treatment is advisable.

Ask your doctor or pharmacist for advice before taking any medicine.

Breast-feeding

Under normal conditions of use, paracetamol may be used during breast-feeding.

Ask your doctor or pharmacist for advice before taking any medicine.

Taking or using other medicines:

This medicine contains paracetamol and this must be taken into account if other medicines containing paracetamol are taken, in order not to exceed the recommended daily dose. Please inform your doctor or pharmacist if you are taking, or have recently taken, any other medicine even one obtained without a prescription.

SIDE EFFECTS

- In rare cases, the following may occur: a malaise, a drop in blood pressure or changes in laboratory test results: abnormally high levels of hepatic enzymes found during blood checks. Should this occur, inform your doctor as regular blood checks may be required later.
- In certain rare cases, a skin rash or allergic reaction may occur. Stop the treatment immediately and inform your doctor.
- In exceptionally rare cases, changes in laboratory test results have been observed which have necessitated regular blood checks: abnormally low levels of some types of blood cells (platelets), possibly leading to bleeding from the nose or gums. Should this occur, inform your doctor.

If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

STORAGE

Store below 30°C.

Do not refrigerate or freeze.

PRESENTATION

Vials

DOLORGAN 1 g/100 ml: Paracetamol 1 g in 100 ml vial

Excipients: mannitol, cysteine hydrochloride monohydrate, disodium phosphate dihydrate, sodium hydroxide, hydrochloric acid, water for injection

THIS IS A MEDICAMENT

- A medicament is a product which affects your health, and its consumption contrary to instructions is dangerous.
- Follow the doctor's prescription strictly, the method of use and the instructions of the pharmacist who sold the medicament.
- The doctor and the pharmacist are experts in medicine, its benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.

