

Paraxone®-JPI
(Chlorzoxazone and Paracetamol)

PROPERTIES

Chlorzoxazone is a long acting effective skeletal muscle relaxant. It is well tolerated as has been proved by long term clinical experience. Paracetamol is an analgesic which is generally non-irritant to the gastro-intestinal tract.

INDICATIONS

Paraxone-JPI is particularly effective in treating pain and spasm of the skeletal muscles arising in:

- Sprains and strains.
- Myalgia
- Torticollis
- Tension headache
- Traumatic muscle damage
- Lumbago
- Spondylarthrosis
- Cervical root syndrome

DOSAGE AND ADMINISTRATION

The recommended dose of Paraxone-JPI is two capsules three times daily after meals (or as prescribed by the physician).

CONTRAINDICATIONS

Known hypersensitivity to one or more of the active ingredients.

PRECAUTIONS

- Use in pregnancy:
Since the safety of Paraxone-JPI in pregnancy is not well established, the treating physician should outweigh the benefit to the mother against the possible risks of the fetus when treating with Paraxone-JPI.
- Change of the urine colour may occur in treatment with Paraxone-JPI, however this change does not implicate any clinical significance.
- In case of hepatic or renal diseases, a routine check up on kidney and liver functions should be performed.
- Treatment with Paraxone-JPI should be discontinued if the patient condition worsened.
- Even in the prescribed dose Paraxone-JPI may delay reaction time, so drivers and people who operate machineries should take this into consideration.
- Concomitant use of Paraxone-JPI with alcohol increases the delay in reaction time.

SIDE EFFECTS

- Less common:
Gastrointestinal irritation, Somnolence, Vertigo, headaches, nausea, over-excitability and allergic skin reactions.
- Rare : Angio-neurotic oedemas, anaphylactic reactions.

PRESENTATION

Capsules

Paraxone-JPI : Chlorzoxazone 250 mg and Paracetamol 300 mg/Capsule.

THIS IS A MEDICAMENT

Medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.
The doctor and the pharmacist are experts in medicine, its benefits and risks.
Do not by yourself interrupt period of treatment prescribed for you.
Do not repeat the same prescription without consulting your doctor.

Keep medicament out of the reach of children