

TERFINIL[®] Cream

Terbinafine Hydrochloride 1%

ACTION

Antifungal agent.

INDICATIONS

Terfinil cream is a medicine (antifungal agent) intended for treatment of skin infections caused by fungi (mycoses) such as athlete's foot or other superficial mycoses of the skin and pityriasis versicolor.

Terfinil cream is also effective against infections caused by yeasts (candidosis of the skin)

Terfinil cream rapidly suppresses symptoms such as itching, desquamation (scaling) and red blotches.

DOSAGE AND ADMINISTRATION

For adults and children above 12 years: before applying Terfinil, carefully wash the affected parts with a cotton wool pad and warm water and then dry carefully.

Apply Terfinil cream to the affected parts and surrounding areas and rub in gently to ensure penetration.

Duration of normal treatment

For adults and children above 12 years:

Athlete's foot, superficial skin mycoses: apply once a day for 1 week.

Skin candidiasis: apply 1-2 times a day for 1 week.

Pityriasis versicolor: apply 1-2 times a day for 2 weeks.

In order to avoid any relapse, follow the treatment as described above.

If after one week there is no improvement, consult your doctor.

Please observe the dosage indicated in this information leaflet or prescribed by your doctor.

If you feel the efficacy of the medicine is too weak or on the contrary too strong, please consult your doctor or pharmacist.

CONTRAINDICATIONS

Do not use this product if you are allergic to one of its components. As the experience in children under 12 years of age is limited, the use of Terfinil cream is not recommended in children under 12 years of age.

PRECAUTIONS

Terfinil cream should only be used externally. Avoid all contact with your eyes.

If by accident Terfinil cream comes into contact with the eyes, it should be carefully flushed with cold water. Consult your doctor, if there is still discomfort.

Please inform your doctor or pharmacist if you suffer from other illnesses, allergies or use other medicines for internal or external use (even under self-medication!).

PREGNANCY AND LACTATION

Terfinil should only be used during pregnancy or lactation under medical authorization.

SIDE EFFECTS

Sometimes, irritations, redness or mild sensation of burning on the skin occur but it is rare that these complaints cause the treatment to be stopped.

Allergic reactions such as itching, rashes and outbreaks of spots are rare. If you notice the appearance of any of these symptoms, stop the treatment and consult your doctor.

STORAGE

Store below 30°C.

PRESENTATIONS

Tube:

TERFINIL Cream 15 g :

Terbinafine Hydrochloride 1%

Excipients: Benzyl alcohol, cetyl alcohol, stearyl alcohol, cetyl palmitate, isopropyl myristate, polysorbate 60, sorbitan monostearate, sodium hydroxide & purified water.

THIS IS A MEDICAMENT

- A medicament is a product which affects your health, and its consumption contrary to instructions is dangerous.
- Follow the doctor's prescription strictly, the method of use and the instructions of the pharmacist who sold the medicament.
- The doctor and the pharmacist are experts in medicine, its benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.

Manufactured by Jazeera Pharmaceutical Industries, Riyadh - Saudi Arabia, for HIKMA Pharmaceuticals, Amman - Jordan



Keep medicament out of the reach of children
2INTFN-AE-04/2006