

CYTAROSE

Antiseptic foaming solution
Antibacterial & Antifungal

Composition:

Each 100 ml contains:

Chlorhexidine di-gluconate 0.10 gm

Hexamidine di-isethionate 0.10 gm

Chlorocresol 0.30 gm

Indications:

Treatment of vaginal infections (that are caused by bacteria and fungi).

Disinfection of birth canal pre and / or post labor.

Other indications as primary skin diseases (e.g. acne, intertrigos, and cutaneous mycosis), disinfections of wounds and hands.

Contraindications:

CYTAROSE is contraindicated in patients with known hypersensitivity for any ingredients of the product.

Precautions:

CYTAROSE is designed for external use only.

There is no evidence of any adverse effects on the fetus arising from the use of **CYTAROSE** during pregnancy or lactation; therefore no special precautions are recommended during pregnancy or lactation.

Adverse effects:

CYTAROSE is well pH balanced tolerated formula but in rare cases sensitivity may develop in certain individuals. If irritation, discomfort, redness, swelling or other signs of sensitivity occur the patient should discontinue treatment.

Dosage & Administration:

CYTAROSE could be used directly or diluted with water to 1:10.

In gynecological diseases :

Use without dilution then wash or use diluted to 1:10 as vaginal douche 1-2 times daily for 7-14 days.

Primary skin diseases:

Use without dilution for cleaning then wash the skin with water or use diluted to 1:10 in dermatological inflammations.

In surgery :

Use as diluted to 1:10 for cleaning and disinfection of wounds .

Hands disinfections:

Use without dilution and rinse thoroughly with water .

Package:

Bottles of 250 ml

Storage:

Store below 25 °C.

Keep away from direct light.
Keep out of reach of children.

For external use only

This is a medicament

Medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.

Follow strictly the doctor's prescription, the method of use and the instructions of the Pharmacist who sold the medicament.

The doctor and the pharmacist are experts in medicine, its benefits and risks.

Do not be hurried: insert the period of treatment prescribed for you.

Do not repeat the same prescription without consulting your doctor.

Keep the medicament out of reach of children.

Council of Arab Health Administrators & Union of Arab Pharmacists



Manufactured by: DEEF Pharmaceutical Industries Co. Saudi Arabia Al Qassim Al Badaye