

suppositories Meclozine Pyridoxine hydrochloride Pyridoxine hydrochloride

PRESENTATION AND FORMULA

- 10 tablets : each tablet contains meclozine hydrochloride 25 mg,
- vitamin B₆ 50 mg. **6 suppositories :** each suppository contains meclozine 30 mg, vitamin B₆ 62,5 mg.

INDICATIONS

Pregnancy nausea and vomiting.

DOSAGE

One tablet or suppository at night, when retiring.

DIRECTIONS FOR USE

The tablets should be swallowed with a little water and, if need be, some food (dry biscuit).

Remove the protective wrapping from the suppositories and administer them after rectal evacuation or, if required, previous enema.

- How to restore accidentally molten suppositories to their original shape 1° Put the plaquettes containing the molten suppositories into cold
- water or into a refrigerator, their tips pointing down.
- 2° If the suppositories have got out of shape because of accidental melting followed by solidification in the wrongly oriented plaquettes, they must be remelted by dipping the plaquettes into water at 40-50 °C and then be treated as specified under 1°.

STORAGE

- Tablets : store below 25 °C in a dry place.
- Suppositories : store in a cool place (8-15 °C).

Should any complaint arise, please mention the batch control number indicated on the package.

UCB S.A. PHARMA SECTOR BRAINE-L'ALLEUD · BELGIUM

© UCB (1984)

R1094422-G

	2-G
 (ان مستلا السيون) (الدين من منك وا تنهذی للدين المسيون عليه وتعليمات بورفك للدين . د العرب محلي المسيد ولي منه المسيب وطريمة الاستعمال المسعوى عليه وتعليمات . د المسيد لاي مالي الحالي . د المسيد المسيد المالي . د المعاد الحالي . د المعاد المالي . د المالي	