movicillin

usilitative and Quantitative Composition (active ingredient or MOXIL Capsules 250 mg contain 250 mg amoxicillin per caps MOXIL Capsules 500 mg contain 500 mg amoxicillin per caps harmaceutical Form MOXIL Capsules: marcon and onld carendae mamrietad **Q**t Al

harmaceutical Form MOVIC Capaules: marcon and gold capsules overprinted GS.JUL: { for 500 mg capsule}. 'GS LEX" { for 250 mg capsule}. UNICAL PARTICULARS dirations

Next of presentations are satisfied in proceynamic. Technology and the second second

Transferret of Intention: Add charge (Including definity plenetable): Add charge (Including definits): Add c

or Omeprazole 40 mg daily, Amoxicillin 750 mg-1 Gm BID, Metronidazole 400 mg TID x 7 days Renal impairment:

Omerand 24 Om gdisky, Amoulialin 720 mpC (Gm BE), Methomataski sau mg reak / kary Real Impairment General Inflation and Schellink Tag Anglutentialities mes Stoping and General Inflation and Collimics. Amountainin, mass. Stoping and General Inflation and Collimics. Amountainin, mass. Stoping and General Inflation and Collimics. Takonologi and anglutes filmes daily for more severe infections. Real Impairment Inflation and et al. (Second Schelling and Sche

riopinyiaus or endocardin	15:			
CONDITION		ADULTS' DOSAGE (INCLUDING ELDERLY)	CHILDREN'S DOSAGE	NOTES
Dental procedures: prophytics' for patients undergoing extraction, scaling or surgery involving ginglival issues and who have not reasived a pericialin in the previous month. (N.B. Patients with proathetic heart valves should be referred to hospital - see below).	Pulsent not having general anaesthetic. Pastent having general ambiotics: considered to be appropriate. Pastent having general anaesthetic: if oral antibiotics not appropriate.	3.9 (*ARCN2: only, 1 hour before products. A second does may be given 8 hours iter, if considered necessary. Taking 3.9 (*ARCN2 coulty + hours prior to ansesthesis, followed by 3.9 con3/pr 1.9 for VII if locates ent belansted) as soon as possible after the speersion. 1.9 (*ARCN2: Yor MI memodatal) before induction, with 500 mg only, 6 hours later.	Under 10: half adult diose. Under 5: quarter adult dose.	Note 1. If prophyticis with "AMCXIC" is given hoice within one morth, mergance of resistant streptococci is unlikely to be a problem. Alternative antibiotics are recommended if more hequest purphylicits is nequined, or if the patient has received a occurse of bushment with a puncifish adampt by previous month. Note 2 To minimize pain on injection, "AMCXII! may be given as two injections of 500 mg dissolved in nuclei 1% Signocal most solution
Dental procedures patients for whom referral to hospital is recommended. 4 Patients to be given a general assachteric who have a Patients to be given a general assachteric who have a possible hart value. 4 Patients who have had one or more attacks of endocardifis.		Initially: 1 g*AMCXXL*N or IM with 120 mg pertamicin N or IM immediately prior to anexthesia (6) given Jon 15 minutes prior to dental procedure. Followed by (8 hours later): 500 mg *AMCXXL* orally.	Under 12: the doses of "AMDXE" should be half the adult dose; the dose of gentamicin should be 2 mg/s. Under 5: the doses of "AMDXE" should be	See Note 2: See Note 2: "AMCXIL" and gentamicin should not be mixed in the same syrings. Note 4: Please consult the appropriate data sheat for full preacribing information on gentamicin.
Gentoxvinary Surgery or Instrumentations prophysias for patients who have no urinary sauri infection and who are to have gento-urinary surgery or instrumentation under general anaethesia. In the case of Obstathic and Gynaecological Procedures and Gastroinsteinal Procedures – routine prophyliais is recommended only for patients with prosthick beaut values.		britistly: 1 g*AMCXXC*/V or MI with 120 mg guntamicin IV or IM, immediately before induction. Followed by (8 hours later): 500 mg "AMCXXC* only or IV or IM according to clinical condition.	quarter the adult door; the doos of gentamicin should be 2 mg/kg.	See Notes 2, 3 and 4 above.
Surpery or Instrumentation of the Upper Respiratory Tract	Patients other than those with prosthetic heart valves.	1 g 'AMCXE' IV or IM immediately before induction; 500 mg 'AMCXE' IV or IM 6 hours later.	Under 10: half adult dose. Under 5: quarter adult drese	See Note 2 above. Note 5. The second dose of 'AMOXE' may be administered onally as 'AMOXE' Syrup GENE
	Patients with prosthetic heart valves.	Initially: 1 gr:AMXD0C VV or M with 120 mg gentamics VV or M, immediately before induction; followed by B hours later 500 mg"AMXDNE" N or R4.	Under 10: the dose of "AMDXIX" should be half the adult dose; the gentamicin dose should be 2 mg kg. Under 5: the dose of "AMDXIX" should be quanter the adult dose; the dose of gentamicin	See Notes 2, 3, 4 and 5 above.
Administration: Out: Treatment should be continued for 2 to 3 days following the disupserance of synctoms. It is economended that at least 10 days treatment by byten for any infection caused by beta harmonity dis streptococci in other to achieve exclusion of the organicm. Anoncellinit is explored in advanced to given to galantees with a history of hypersentitivity to beta-least multikotic leg pencilline, cephalogonitis. Biologoni disped Presentions for the cephalogonitis. Biologoni and Speed Presentions in the nate concenting period hypersentitivity to beta least multikotics and concentinal the presentiality (association and the cephalogonitis). Cephalogonitis. Biologonia are one likely to concer in individual and a history of hypersentitivity to beta least				
Contranacations; A moxicillin should be avoided if infectious mononucleosis is suspected since the occurrence of a morbilliform rash has been associated with this condition following the use of amoxicillin.				
Promogeu use may also occasionairy result in overgrowth or non-subcoptible organisms. Abnormal profongation of prothombin time (increased INP) has been reported arrely in patients receiving amoxicilin and oral anticoogulants. Appropriate monitoring should be undertaken when anticoogulants are prescribed concurrently. Adjustments in the dose				

.ins. These : .ndication: .xicillin shoul this conditio. trigged use ms yrmal protor bagulants il antico 'ge sh' jer' '

In the second second

A second of a second second

(10) Use that 110, excention in one that 1500, test than 1500, test into 1500, test than 1500, ever rare (see 110,000, test has 1500, test than 1500, test ction or in those receiving high

ls. Vers Gastr. # Comm # Uncor Very rai 'ack h

- un en montoria associated calling finalularg paradomenhianeous collis and Sapahala tomo discussion has been reported in children. Good said hypein. Neuro bill report of the second in the second second second second heart of the second second second second second second second second 2 common Silvaria and childrate (second second second second second 2 common Silvaria). A second second second second second second second 2 common Silvaria and second second second second second second second 2 common Silvaria and second second

Nensa and Unitary tract disorders Very rare: Interstitial nephritis, crystalluria (see Overdose) # The incidence of these AEs was derived from clinical studies involving a total of approxit (6,000 adult and paediatric patients tai

ne, tax ic epidern

Overdose Gastrointestinal effects such as nausea, vomiting and diarrhoea may the waterrelectrolyte balance. Amoxicillin crystalluria, in some cases leading to renal failure, has be Amoxicillin crystalluria, in some cases leading to renal failure, has be ManuAccEURCLA PARTICULARS atically with attention to en oi nd p

PHARMACEUTICAL PARTICULARS List of Excipients Each capsule contains magnesium stearate (I oxide [E172] and gelatin. Shell Life The expiry date is indicated on the packaging. (E127) i

outlet [17] and gelatin. The expiry date in statustion on the packaging. Special Proceedings of Storage Donot store above 2000 °C Add/XXI is a rosistmant of the GlaudomithKline group of cor Smithlice Beschuli Imitelt⁴⁴ Worthing, I Smithlice Reschul Imitelt⁴⁴ Worthing, I Glaus Velkioner Production⁴⁴ Mayrens, France Smithlice Beschul Group of companies Mayrens, France 2000 (subcombiting group of companies (GSV Version Humber 2010)

gsk GlaxoSmithKline

INSTRUCTORS TO THE PATIENT THIS IS A NEDCOMENT Medicament is a product within allers your health, and consumption contrary to instructions is dangerous for you. Followstrichly the docts's precription, the method of use and instructions of the pharmacist are experts in medicine. Declar and the Pharmacist are experts in medicine, benefits and risks. cine, it ed fo I here the same prescription warout consuming on or Keep medicament out of the reach of children Council of Arab Health Mit Using of Arab Health Mit