Levoflox® 5 mg/ml 5 mg/ml, solution for intravenous infusion Levofloxacin

Read all of this leaflet cavefully before you start taking this medicine

- Keep this leaflet, you may need to read it again.

If you have any further questions, sak your doctor or planmacist.

- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same asyours.

- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

Inthis leaflet:

Serious side effects
What Levoflox Smg/ml is and what it is used for
Before you take Levoflox Smg/ml
How you take Levoflox Smg/ml
Possible side effects
How to store Levoflox Smg/ml
Further information

6. How to store Levoflox Singiml
7. Further mitorianstics
1. Sections side effects
1. The committee of the c

Levolto. Sing/ml solution for intravenous infusion can be used to treat infections of the:

- Lungs, in people with pneumonia

- Urinary tried, including your kidneys or bladder

- Porstate gland, where you have a long lasting infection

- Prostate gland, where you have a long lasting infection

- Prostate gland, where you have a long lasting infection

In some special situations, Levolfox Sing/ml solution for intravenous infusion may be used to

lessen the chances of getting a pulmonary disease amend anthrax or worsening of the disease
after you are exposed to the bacteria causing anthrax.

- Before you fake "Levoltox Sing/ml"

- Donot take levoltox 5 ing/ml*

- Donot take levoltox 5 ing/ml*

- Donot take levoltox 5 ing/ml*

- One of the control of the control

*If you are called a length of eventions and any other quandons antibutic such as monthocan.

If Shan of an allergat root too include a rath a wallowing or breathing problems, swelling of your lips, face, throat or tongue.

If You have ever had epilegy with your tendines cach as tendonits that was related to treatment with a support of the problems. The problems is the cord that joins your muscle to your skeleton.

If You are a called to a gowing tecnages.

If You are a called to a gowing tecnages.

If You are pregnant, might become pregnant, or think you may be pregnant.

If You are pregnant, might become pregnant, or think you may be pregnant.

If You are pregnant, might become pregnant, or think you may be pregnant.

If You are pregnant, or the problems of the problems of your skeleton.

If you are a called to a gowing of the above applies to you.

If you are not made to the problems of your skeleton, must eer pharmacist before you are given Levollos (sing.ml.)

False to your doctor, musr or pharmacist before you have your medicine if.

You have ever had a fit (seizure)

You have ever had heart problems

You have ever had heart problems or glucose—6—phosphate dehydrogenuse deficiency.

You have ever had heart problems such your bed of medicine, if you were born with or have sain inbalance in the blood (seepschild you level of potassines such of medicine, if you were born with or have sain inbalance in the blood (seepschild you level of potassines is kind of medicine, if you were born with or have sain inbalance in the blood (seepschild you level of potassines is kind of medicine that result in abnormal ECG changes (see section. "Other medicines and Levollox Sing/ml").

You have ever had heart problems is kind of medicine, if you were form the history of heart attack (myocardial infarction), you are fenale or elderly or you are fating in the blood, have a very slow heart

1 On mave myastnenia gravis.
1 On mave myastnenia gravis.
2 On mave myastnenia gravis.
2 On mave myastnenia gravis.
3 On mave myastnenia gravis.
3 On myastnenia gravis.
3 On myastnenia gravis.
3 On myastnenia gravis.
4 On myastnenia gravis.
5 On myastnenia gravis.
5 On myastnenia gravis.
5 On myastnenia gravis.
6 On myastnenia gravis.
7 On myastnenia gravis.
8 On myastnenia gravis.
9 On

clambiromycinj.

Probenecid – used for gout. Your doctor may want to use a lower dose, if you have kiduey problems.

*Cimedmine – used for ulcers and hearthum. Your doctor may want to use a lower dose, if you have kiduey problems.

Fell your doctor if any of the above applies to you.

Unive tests may slow 'false-positive' results for storag painkillers called 'opates' in people having Levoflox 5 mg/ml.

Tuber cutosis tests any slow 'false positive' results for storag painkillers called 'opates' in people having Levoflox 5 mg/ml.

Tuber cutosis tests

Taberculous tests

It is medicine may cause "false negative" results for some laboratory tests that search for the bacteria that cause tuberculous.

Preganacy and breast-feeding

Do not have this medicine if

Do not five or carry out any work that requires a high level of attention.

Do not drive or carry out any work that requires a high level of attention.

Do not drive or carry out any work that requires a high level of attention.

Do not drive or carry out any work that requires a high level of attention.

This medicine contains \$40 mg of soddium per dose of 500 mg levofloxacin. This should be taken into consideration by patients on a controlled soddium diet.

How you take Levoflox Smgml

How Levoflox Sngml solution for mouse influsion is a medicine for use in hospitals.

It will be gaven to you by a doctor or muse as an injection. The injection will be into one of your veins and be given over a period of time (this is called an intervalous influsion) in should be 60 minutes or more your heart ride and blood pressure a spossible side effects that have been spen during the influsion of a similar antibiofic. If your blood pressure aroposible side effects that have been seen during the influsion of a similar antibiofic. If your blood pressure drops noticeably while you are being given the finds on it is given.

If you are not sure why you are being given Levoflox Smgml you should have the dose will depend on the type of the least of the properties of the least of the lea

intravenous infusion
Oxacin

a. If you take more Levoltox Smg/ml solution for intravenous infusion than you should
it is unlikely that your doctor or murse will give you too much medicine.

A third way oak if you are not sure will give you too much medicine.

A third way oak if you are not sure why you are gettin, and decide the medicine and the control of the con

Unusual first beatung of your heart (nethywarda) or low blood pressure (hypotension)

Muscle weslesses: This is important in people with mynathenia gravis (a run desease of the nervous system). Changes in the way your kiney works and occasional kindey failure which may be due to an allergic kindney resection called interstitual nephriss.

Not known (frequency cannot be estimated from the available data)

Not known (frequency cannot be estimated from the available data)

Not known (frequency cannot be estimated from the available data)

Not known (frequency cannot be estimated from the available data)

Not known (frequency cannot be estimated from the available data)

Not known (frequency cannot be estimated from the available data)

Not known (frequency cannot be estimated from the available data)

Not known (frequency cannot be estimated from the available data)

Not known (frequency cannot be estimated from the available data)

Not known (frequency cannot be estimated from the available data)

Not known (frequency cannot be estimated from the available data)

Loss of circulation (anaphylactic like shock)

Loss of circulation (anaphyl

Tel: 00 96 1 46 46 955 / Fax: 00 966 1 4634 362

d- This leaflet was last approved in "August 2014"; version number "00"

e- To report any side effect(s):

Said Arabia:

- Saija Arabus:
The National Pharmacovigilance and Drug Safety Cenwe (NPC)
Fax: +8667662-205-115-222-11, Exts: 23172340-2334-2354-2553-2556of Toll free phone: 800249000
G. E-majl: npc.drug@sfda.gov.sa
Website: www.sida.gov.sanpx

Other GCC states:
 Please contact the relevant competent authority.

f. Council of Arab Health Ministers

Medicament is a product which affects your beath and its consumption centrary to instructions is changerous for your beath and its consumption centrary to instructions is changerous for you search; the decire is precapitous for method of use and the unstructions of the plantancist who rold the medicament. Do not by yourself asterning the period of treatment prescribed for your decire.

Do not prefet the same prescription without consulting your decire.

ut of reach of children KEEPANYDRUG OUT OF THE REACH OF CHILDREN. Council of Arab Health Ministers

NAMES OF

2. This patient information leaflet is approved by the Saudi Food and Drug Authority