Oxaliplatin Concentrate for Solution for Infusion OXITAN

Anaphylactic reactions to Oxaliplatin have been reported, and may occur within minutes of Oxaliplatin administration. Epinephrine, corticosteroids, and antihistamine have been employed to alleviate symptoms of anaphylaxis.

Oxaliplatin Concentrate for Solution for Infusion (Oxitan) is an antineoplastic agent Oxaliplatin is an organoplatinum complex in which the platinum atom is complexed with 1,2-diaminocyclohexane (DACH) and with an oxalate ligand as a leaving group.

COMPOSITION Each mL contains: Oxaliplatin BP 5.0 mg Water for Injection BP

The molecular formula of oxaliplatin is C₈H₁₄N₂O₄Pt and the chemical name is cis-[(1R,2R)-1,2-cyclohexanediamine-N,N] [oxalato(2-)-O,O] platinum.

The molecular weight is 397.3. Oxaliplatin is sparingly soluble in water.

PHARMACOLOGY

Mechanism of Action

Oxaliplatin undergoes nonenzymatic conversion in physiologic solutions to active derivatives via displacement of the labile oxalate ligand. Several transient reactive species are formed, including monoaquo and diaquo DACH platinum, which covalently bind with macromolecules. Both inter- and intrastrand Pt-DNA crosslinks are formed Crosslinks are formed between the N7 positions of two adjacent guanines (GG), adjacent adenine-guanines (AG), and guanines separated by an intervening nucleotide (GNG). These crosslinks inhibit DNA replication and transcription. Cytotoxicity is cellcycle nonspecific.

In vivo studies have shown antitumor activity of oxaliplatin against colon carcinoma In combination with 5-fluorouracil (5-FU), oxaliplatin exhibits in vitro and in vivo antiproliferative activity greater than either compound alone in several tumor models [HT29 (colon), GR (mammary), and L1210 (leukemia)].

The reactive oxaliplatin derivatives are present as a fraction of the unbound platinum in plasma ultrafiltrate. The decline of ultrafilterable platinum levels following oxaliplatin administration is triphasic, characterized by two relatively short distribution phases (t 0.43 hours and t_{1/28}: 16.8 hours) and a long terminal elimination phase (t_{1/28}: 391 hours) Pharmacokinetic parameters obtained after a single 2-hour IV infusion of oxaliplatin at a dose of 85 mg/m 2 expressed as ultrafilterable platinum were C_{max} of 0.814 μ g/mL and volume of distribution of 440 I

Interpatient and intrapatient variability in ultrafilterable platinum exposure (AUC assessed over 3 cycles was moderate to low (23% and 6%, respectively). A pharmacodynamic relationship between platinum ultrafiltrate levels and clinical safety and effectiveness has not been established

At the end of a 2-hour infusion of oxaliplatin, approximately 15% of the administered platinum is present in the systemic circulation. The remaining 85% is rapidly distributed into tissues or eliminated in the urine. In patients, plasma protein binding of platinum is irreversible and is greater than 90%. The main binding proteins are albumin and gammaglobulins. Platinum also binds irreversibly and accumulates (approximately 2-fold) in erythrocytes, where it appears to have no relevant activity. No platinum accumulation was observed in plasma ultrafiltrate following 85 mg/m² every two weeks

Oxaliplatin undergoes rapid and extensive nonenzymatic biotransformation. There is no

evidence of cytochrome P450-mediated metabolism in vitro. Up to 17 platinum-containing derivatives have been observed in plasma ultrafiltrate samples from patients, including several cytotoxic species (monochloro DACH platinum, dichloro DACH platinum, and monoaquo and diaquo DACH platinum) and a

The major route of platinum elimination is renal excretion. At five days after a single 2-hour infusion of oxaliplatin, urinary elimination accounted for about 54% of the platinum eliminated, with fecal excretion accounting for only about 2%. Platinum was cleared from plasma at a rate (10 – 17 L/h) that was similar to or exceeded the average human glomerular filtration rate (GFR; 7.5 L/h). There was no significant effect of gender on the clearance of ultrafilterable platinum. The renal clearance of ultrafilterable platinum is significantly correlated with GFR.

Pharmacokinetics in Special Populations

number of noncytotoxic, conjugated species.

Renal Impairment

A study conducted in 38 patients with advanced GI cancer and varying degrees of renal impairment with patients in the normal (creatinine clearance (CrCL) > 80 mL/ min, N=11), mild (CrCL=50-80 mL/min, N=13), and moderate (CrCL=30-49 mL/min N=10) groups treated with 85 mg/m² Oxaliplatin and those in the severe (CrCL < 30 mL/min, N=4) group treated with 65 mg/m² Oxaliplatin. The mean AUC of unbound platinum was 40%, 95%, and 342% higher in the mild, moderate, and severe groups, respectively, than in the normal group. Mean $C_{\mbox{\scriptsize max}}$ of unbound platinum appeared to be similar among the normal, mild and moderate renal function groups, but was 38% higher in the severe group than in the normal group. Caution should be exercised in renally impaired patients. The starting dose of Oxaliplatin should be reduced in patients

Drug - Drug Interactions

No pharmacokinetic interaction between 85 mg/m² of oxaliplatin and 5-FU has been observed in patients treated every 2 weeks, but increase of 5-FU plasma concentrations by approximately 20% have been observed with doses of 130 mg/m² of oxaliplatin administered every 3 weeks. In vitro, platinum was not displaced from plasma proteins by the following medications: erythromycin, salicylate, sodium valproate, granisetron, and paclitaxel. In vitro, oxaliplatin is not metabolized by, nor does it inhibit, human cytochrome P450 isoenzymes. No P450-mediated drug-drug interactions are therefore anticipated in patients

Since platinum-containing species are eliminated primarily through the kidney, clearance of these products may be decreased by co-administration of potentially nephrotoxic compounds, although this has not been specifically studied.

CLINICAL STUDIES

Combination Adjuvant Therapy with Oxaliplatin and Infusional 5-FU/LV in

Patients with Colon Cancer An international, multicenter, randomized study compared the efficacy and evaluated the safety of oxaliplatin in combination with an infusional schedule of 5-FU/LV to infusional 5-FU/LV alone, in patients with stage II (Dukes' B2) or III (Dukes' C) colon cancer who had undergone complete resection of the primary tumor. The primary objective of the study was to compare the 3-year disease-free survival (DFS) in patients receiving exaliplatin and infusional 5-FU/LV to those receiving 5-FU/LV alone. Patients were treated for a total of 6 months (i.e., 12 cycles). A total of 2246 patients were randomized; 1123 patients per study arm. Patients in the study had to be between 18 and 75 years of age, have histologically proven stage II (T_3 - T_4 , N0, M0; Dukes' B2) or III (any T, N_{1.2} M0; Dukes' C) colon carcinoma (with the inferior pole of the tumor above the peritoneal reflection, i.e., >15 cm from the anal margin) and undergone (within 7 weeks prior to randomization) complete resection of the primary tumor without gross or microscopic evidence of residual disease. Patients had to have had no prior chemotherapy, immunotherapy, or radiotherapy, and have an ECOG performance status of 0, 1, or 2 (KPS ≥60%), absolute neutrophil count (ANC) >1.5x109/L, platelets

 \geq 100x10°/L, serum creatinine \leq 1.25 x ULN, total bilirubin <2 x ULN, AST/ALT <2 x ULN and carcino-embryogenic antigen (CEA) <10 ng/mL. Patients with preexisting peripheral neuropathy (NCI grade \geq 1) were ineligible for this trial.

The following table shows the dosing regimens for the two arms of the study.

Table 1 – Dosing Regimens in Adjuvant Therapy Study			
Dose	Regimen		
Day 1: Oxaliplatin: 85 mg/m² (2-hour infusion) + LV 200 mg/m² (2-hour infusion), followed by 5-FU: 400 mg/m² (bolus), 600 mg/m² (22-hour infusion)	q2w		
Day 2: LV 200 mg/m² (2-hour infusion), followed by 5-FU: 400 mg/m² (bolus), 600 mg/m² (22-hour infusion)	12 cycles		
Day 1: LV 200 mg/m² (2-hour infusion), followed by 5-FU: 400 mg/m² (bolus), 600 mg/m² (22-hour infusion) Day 2: LV 200 mg/m² (2-hour infusion), followed by 5-FU: 400 mg/m² (bolus), 600 mg/m² (22-hour	q2w 12 cycles		
	Dose Day 1: Oxaliplatin: 85 mg/m² (2-hour infusion) + LV 200 mg/m² (2-hour infusion), followed by 5-FU: 400 mg/m² (bolus), 600 mg/m² (22-hour infusion)) Day 2: LV 200 mg/m² (2-hour infusion), followed by 5-FU: 400 mg/m² (bolus), 600 mg/m² (22-hour infusion)) Day 1: LV 200 mg/m² (2-hour infusion), followed by 5-FU: 400 mg/m² (2-hour infusion) Day 2: LV 200 mg/m² (2-hour infusion), followed by 5-FU: 400 mg/m² (2-hour infusion)) Day 2: LV 200 mg/m² (2-hour infusion), followed		

The following tables show the baseline characteristics and dosing of the patient population entered into the study. The baseline characteristics were well balanced

Table 2 - Patient Characteristics in Adjuvant Therapy Study

	Oxaliplatin + infusional 5-FU/ LV N=1123	Infusional 5-FU/ LV N=1123
Sex: Male (%)	56.1	52.4
Female (%)	43.9	47.6
Median age (years)	61.0	60.0
<65 years of age (%)	64.4	66.2
≥65 years of age (%)	35.6	33.8
Karnofsky Performance Status (KPS) (%	b)	•
100	29.7	30.5
90	52.2	53.9
80	4.4	3.3
70	13.2	11.9
≤60	0.6	0.4
Primary site (%)	'	•
Colon including cecum	54.6	54.4
Sigmoid	31.9	33.8
Recto sigmoid	12.9	10.9
Other including rectum	0.6	0.9
Bowel obstruction (%)		•
Yes	17.9	19.3
Perforation (%)	<u> </u>	
Yes	6.9	6.9
Stage at Randomization (%)		
II (T=3,4 N=0, M=0)	40.1	39.9
III (T=any, N=1,2, M=0)	59.6	59.3
IV (T=any, N=any, M=1)	0.4	0.8
Staging – T (%)	·	•
T1	0.5	0.7
T2	4.5	4.8
T3	76.0	75.9
T4	19.0	18.5
Staging – N (%)		
N0	40.2	39.9
N1	39.4	39.4
N2	20.4	20.7
Staging – M (%)	<u> </u>	
M1	0.4	0.8

	Oxaliplatin + infusional 5-FU/ LV N=1108	Infusional 5-FU/LV N=1111
Median Relative Dose Intensity (%)		
5-FU	84.4	97.7
Oxaliplatin	80.5	N/A
Median Number of Cycles	12	12
Median Number of cycles with Oxaliplatin	11	N/A

The following table summarizes the disease-free survival (DFS) results in the overall randomized population and in patients with stage II and III disease based on an ITT analysis. The median duration of follow-up was approximately 77 months.

	Oxaliplatin + Infusional 5-FU/LV	Infusional 5-FU/ LV		
Parameter				
	Overall			
N	1123	1123		
Number of events – relapse or death (%)	304 (27.1)	360 (32.1)		
Disease-free survival % [95% CI] *	73.3 [70.7, 76.0]	67.4 [64.6, 70.2]		
Hazard ratio [95% CI] **	0.80 [0.68, 0.93]			
Stratified Log rank test	p=0.003			
Stage III (Dukes' C)				
N	672 675			
Number of events –relapse or death (%)	226 (33.6)	271 (40.1)		
Disease-free survival % [95% CI] *	66.4 [62.7, 70.0]	58.9 [55.2, 62.7]		
Hazard ratio [95% CI] **	0.78 [0.65,	0.93]		
Log rank test	p=0.00	5		
Stag	e II (Dukes' B2)			
N	451	448		
Number of events – relapse or death (%)	78 (17.3)	89 (19.9)		
Disease-free survival % [95% CI] *	83.7 [80.2, 87.1]	79.9 [76.2, 83.7]		
Hazard ratio [95% CI] **	0.84 [0.62,	1.14]		
Log rank test	p=0.25	8		

* Disease-free survival at 5 years **A hazard ratio of less than 1.00 favors Oxaliplatin + Infusional 5-fluorouracil/leucovorin In the overall and stage III colon cancer populations DFS was statistically significantly improved in the Oxaliplatin combination arm compared to infusional 5-fluorouracil/ leucovorin alone. However, a statistically significant improvement in DFS was not noted

The following table summarizes the overall survival (OS) results in the overall randomized population and in patients with stage II and III disease, based on the ITT

in Stage II patients.

Table 5 - Summary of OS analysis - ITT analysis					
Parameter Oxaliplatin + Infusional Infusional 5-FU/LV Infusional 5-FU/LV					
Overall					
N	1123	1123			

	1		
Number of death events (%)	245 (21.8)	283 (25.2)	
Hazard ratio* [95% CI]	0.84 [0.71 , 1.00]		
Stage III (Dukes' C)			
N	672	675	
Number of death events (%)	182 (27.1)	220 (32.6)	
Hazard ratio* [95% CI]	0.80 [0.65 , 0.97]		
Stage II (Dukes' B2)			
N	451	448	
Number of death events (%)	63 (14.0)	63 (14.1)	
Hazard ratio* [95% CI]	1.00 [0.70 , 1.41]		
*A hazard ratio of less than 1.00 favors Oxaliplatin + Infusional 5-fluorouracil/leucovorin			

Data cut off for overall survival 16 January 2007

Combination Therapy with Oxaliplatin and 5-fluorouracil/leucovorin in Patients Previously Untreated for Advanced Colorectal Cancer A North American, multicenter, open-label, randomized controlled study was sponsored

by the National Cancer Institute (NCI) as an intergroup study led by the North Central Cancer Treatment Group (NCCTG). The study had 7 arms at different times during its conduct, four of which were closed due to either changes in the standard of care, toxicity, or simplification. During the study, the control arm was changed to Irinotecan plus 5-fluorouracil/leucovorin. The results reported below compared the efficacy and safety of two experimental regimens, Oxaliplatin in combination with infusional 5-fluorouracil/ leucovorin and a combination of Oxaliplatin plus Irinotecan, to an approved control regimen of Irinotecan plus 5-fluorouracil/leucovorin in 795 concurrently randomized patients previously untreated for locally advanced or metastatic colorectal cancer. After completion of enrollment, the dose of Irinotecan plus 5-fluorouracil/leucovorin was decreased due to toxicity. Patients had to be at least 18 years of age, have known locally advanced, locally recurrent, or metastatic colorectal adenocarcinoma not curable by surgery or amenable to radiation therapy with curative intent, histologically proven colorectal adenocarcinoma, measurable or evaluable disease, with an ECOG performance status 0,1, or 2. Patients had to have granulocyte count ≥ 1.5 x 10⁹/L, platelets ≥ 100 x 10°/L, hemoglobin ≥9.0 gm/dL, creatinine ≤ 1.5 x ULN, total bilirubin \leq 1.5 mg/dL, AST \leq 5 x ULN, and alkaline phosphatase \leq 5 x ULN. Patients may have received adjuvant therapy for resected Stage II or III disease without recurrence within 12 months. The patients were stratified for ECOG performance status (0, 1 vs. 2), prior adjuvant chemotherapy (yes vs. no), prior immunotherapy (yes vs. no), and age (<65 vs. ≥65 years). Although no post study treatment was specified in the protocol, 65 to 72% of patients received additional post study chemotherapy after study treatment discontinuation on all arms. Fifty-eight percent of patients on the Oxaliplation plus 5-fluorouracil/leucovorin arm received an Irinotecan-containing regimen and 23% of patients on the Irinotecan plus 5-fluorouracil/leucovorin arm received Oxaliplatin containing regimens. Oxaliplatin was not commercially available during the trial.

The following table presents the dosing regimens of the three arms of the study.

Table 6 - Dosing Regimens in Patients Previously Untreated for Advanced

Treatment Arm	Dose	Regimen
Oxaliplatin + 5-FU/LV (FOL- FOX4) (N=267)	Day 1: Oxaliplatin: 85 mg/m² (2-hour infusion) + LV 200 mg/m² (2-hour infusion), followed by 5-FU: 400 mg/m² (bolus), 600 mg/m² (22-hour infusion). Day 2: LV 200 mg/m² (2-hour infusion),	every 2 weeks
1 0,4) (11-201)	followed by 5-FU: 400 mg/m² (bolus), 600 mg/m² (22-hour infusion)	
Irinotecan + 5-FU/LV (IFL) (N=264)	Day 1: irinotecan 125 mg/m² as a 90–min infusion + LV 20 mg/m² as a 15-min infusion or intravenous push, followed by 5-FU 500 mg/m² intravenous bolus weekly x 4	every 6 weeks
Oxaliplatin + Irinotecan (IROX) (N=264)	Day 1: Oxaliplatin: 85 mg/m² intravenous (2 hour infusion) + irinotecan 200 mg/m² intravenous over 30 minutes	every 3 weeks

Table 7 – Patient Demographics in Patients Previously Untreated for Advanced

	Oxaliplatin + 5-FU/LV N=267	Irinotecan + 5-FU/ LV N=264	Oxaliplatin + irinotecan N=264
Sex: Male (%)	58.8	65.2	61.0
Female (%)	41.2	34.8	39.0
Median age (years)	61.0	61.0	61.0
<65 years of age (%)	61	62	63
≥65 years of age (%)	39	38	37
ECOG (%)			
0.1	94.4	95.5	94.7
2	5.6	4.5	5.3
Involved organs (%)			
Colon only	0.7	0.8	0.4
Liver only	39.3	44.3	39.0
Liver + other	41.2	38.6	40.9
Lung only	6.4	3.8	5.3
Other (including lymph nodes)	11.6	11.0	12.9
Not reported	0.7	1.5	1.5
Prior radiation (%)	3.0	1.5	3.0
Prior surgery (%)	74.5	79.2	81.8
Prior adjuvant (%)	15.7	14.8	15.2

The length of a treatment cycle was 2 weeks for the Oxaliplatin and 5-fluorouracil/ leucovorin regimen; 6 weeks for the Irinotecan plus 5-fluorouracil/leucovorin regimen and 3 weeks for the Oxaliplatin plus Irinotecan regimen. The median number of cycles administered per patient was 10 (23.9 weeks) for the Oxaliplatin and 5-fluorouracil/ leucovorin regimen, 4 (23.6 weeks) for the Irinotecan plus 5-fluorouracil/leucovorin regimen, and 7 (21.0 weeks) for the Oxaliplatin plus Irinotecan regimen. Patients treated with the Oxaliplatin and 5-fluorouracil/leucovorin combination had a significantly longer time to tumor progression based on investigator assessment, longer overall surviva and a significantly higher confirmed response rate based on investigator assessment compared to patients given Irinotecan plus 5-fluorouracil/leucovorin. The following table summarizes the efficacy results.

Table 8 - Summary of Efficacy

	Oxaliplatin + 5-FU/LV N=267	rinotecan + 5-FU/LV N=264	Oxaliplatin + Irinotecan N=264
Survival (ITT)			
Number of deaths N (%)	155 (58.1)	192 (72.7)	175 (66.3)
Median survival (months)	19.4	14.6	17.6
Hazard Ratio and (95% confidence interval)	0.65 (0.53-0.80)*		
P-value	<0.0001*	-	-
TTP (ITT, investigator assessment)			
Percentage of progressors	82.8	81.8	89.4
Median TTP (months)	8.7	6.9	6.5
Hazard Ratio and (95% confidence interval) ***	0.74 (0.61-0.89)*		
P-value	0.0014*	-	-
Response Rate (investigator assessm	ent)**		
Patients with measurable disease	210	212	215
Complete response N (%)	13 (6.2)	5 (2.4)	7 (3.3)
Partial response N (%)	82 (39.0)	64 (30.2)	67 (31.2)
Complete and partial response N (%)	95 (45.2)	69 (32.5)	74 (34.4)

95% confidence interval	(38.5 – 52.0)	(26.2 – 38.9)	(28.1 – 40.8)
P-value	0.0080*	-	-

**Based on all patients with measurable disease at baseline The numbers in the response rate and TTP analysis are based on unblinded investigator assessment.

***A hazard ratio of less than 1.00 favors Oxaliplatin + Infusional 5-fluorouracil/

A descriptive subgroup analysis demonstrated that the improvement in survival for Oxaliplatin plus 5-fluorouracil/leucovorin compared to Irinotecan plus 5-fluorouracil/ leucovorin appeared to be maintained across age groups, prior adjuvant therapy, and number of organs involved. An estimated survival advantage in Oxaliplatin plus 5-fluorouracil/leucovorin versus Irinotecan plus 5-fluorouracil/leucovorin was seen in both genders; however it was greater among women than men. Insufficient subgroup

Combination Therapy with Oxaliplatin and 5-fluorouracil/leucovorin in Previously Treated Patients with Advanced Colorectal Cancer

A multicenter, open-label, randomized, three-arm controlled study was conducted in the US and Canada comparing the efficacy and safety of Oxaliplatin in combination with an infusional schedule of 5-fluorouracil/leucovorin to the same dose and schedule of 5-fluorouracil/leucovorin alone and to single agent Oxaliplatin in patients with advanced colorectal cancer who had relapsed/progressed during or within 6 months of first-line therapy with bolus 5-fluorouracil/leucovorin and Irinotecan. The study was analyzed for response rate after 450 patients were enrolled. Survival was assessed in all patients enrolled in the completed study. Accrual was completed, with 821 patients enrolled. Patients in the study had to be at least 18 years of age, have unresectable, measurable, histologically proven colorectal adenocarcinoma with a Karnofsky performance status >50%. Patients had to have SGOT(AST) and SGPT(ALT) ≤2x the institution's upper limit of normal (ULN), unless liver metastases were present and documented at baseline by CT or MRI scan, in which case ≤5x ULN was permitted. Patients had to have alkaline phosphatase ≤2x the institution's ULN, unless liver metastases were present and documented at baseline by CT or MRI scan, in which cases ≤5x ULN was permitted. Prior radiotherapy was permitted if it had been completed at least 3 weeks The dosing regimens of the three arms of the study are presented in the table below.

Table 9 – Dosing Regimens in Refractory and Relapsed Colorectal Cancer Clinical

Treatment Arm	Dose	Regimen
Oxaliplatin + 5-FU/LV (N =152)	Day 1: Oxaliplatin: 85 mg/m² (2-hour infusion) + LV 200 mg/m² (2-hour infusion), followed by 5-FU: 400 mg/m² (bolus), 600 mg/m² (22-hour infusion)	every 2 weeks
	Day 2: LV 200 mg/m² (2-hour infusion), followed by 5-FU: 400 mg/m² (bolus), 600 mg/m² (22-hour infusion)	
5-FU/LV (N=151)	Day 1: LV 200 mg/m 2 (2-hour infusion), followed by 5-FU: 400 mg/m 2 (bolus), 600 mg/ m 2 (22-hour infusion)	every 2 weeks
	Day 2: LV 200 mg/m² (2-hour infusion), followed by 5-FU: 400 mg/m² (bolus), 600 mg/ m² (22-hour infusion)	
Oxaliplatin (N=156)	Day 1: Oxaliplatin 85 mg/m² (2-hour infusion)	every 2 weeks
	ne study for evaluation of response must have h measuring ≥20mm using conventional CT or	

≥10mm using a spiral CT scan. Tumor response and progression were assessed every 3 cycles (6 weeks) using the Response Evaluation Criteria in Solid Tumors (RECIST) until radiological documentation of progression or for 13 months following the first dose of study drug(s), whichever came first. Confirmed responses were based on two tumor assessments separated by at least 4 weeks. The demographics of the patient copulation entered into the study are shown in the table below

Table 10 - Patient Demographics in Refractory and Relapsed Colorectal Cancer Clinical Trial

	5-FU/LV	Oxaliplatin	Oxaliplatin + 5-FU/
	(N = 151)	(N = 156)	LV
			(N = 152)
Sex: Male (%)	54.3	60.9	57.2
Female (%)	45.7	39.1	42.8
Median age (years)	60.0	61.0	59.0
Range	21-80	27-79	22-88
Race (%)			
Caucasian	87.4	84.6	88.8
Black	7.9	7.1	5.9
Asian	1.3	2.6	2.6
Other	3.3	5.8	2.6
KPS (%)			
70 – 100	94.7	92.3	95.4
50 - 60	2.6	4.5	2.0
Not reported	2.6	3.2	2.6
Prior radiotherapy (%)	25.2	19.2	25.0
Prior pelvic radiation (%)	18.5	13.5	21.1
Number of metastatic sites	(%)		
1	27.2	31.4	25.7
≥2	72.2	67.9	74.3
Liver involvement (%)		•	
Liver only	22.5	25.6	18.4
Liver + other	60.3	59.0	53.3

The median number of cycles administered per patient was 6 for the Oxaliplatin and 5-fluorouracil/leucovorin combination and 3 each for 5-fluorouracil/leucovorin alone and

Patients treated with the combination of Oxaliplatin and 5-fluorouracil/leucovorin had an increased response rate compared to patients given 5-fluorouracil/leucovorin or Oxaliplatin alone. The efficacy results are summarized in the tables below.

Table 11 - Response Rates (ITT Analysis)

Best Response	5-FU/LV (N=151)	Oxaliplatin (N=156)	Oxaliplatin + 5-FU/LV (N=152)
CR	0	0	0
PR	0	2 (1%)	13 (9%)
p-value	0.0002 for 5	5-FU/LV vs. Oxalipla	atin + 5-FU/LV
95%CI	0-2.4%	0.2-4.6%	4.6-14.2%
Table 12 - Summary of Radio	graphic Time to	Progression*	

Arm	5-FU/LV (N=151)	Oxaliplatin (N=156)	Oxaliplatin + 5-FU/LV (N=15
No. of Progressors	74	101	50
No. of patients with no radiological evaluation beyond baseline	22 (15%)	16 (10%)	17 (11%)
Median TTP (months)	2.7	1.6	4.6
95% CI	1.8-3.0	1.4-2.7	4.2-6.1

This is not an ITT analysis. Events were limited to radiographic disease progression documented by independent review of radiographs. Clinical progression was not included in this analysis, and 18% of patients were excluded from the analysis based on unavailability of the radiographs for independent review.

At the time of the interim analysis 49% of the radiographic progression events had occurred. In this interim analysis an estimated 2-month increase in median time to radiographic progression was observed compared to 5-fluorouracil/leucovorin alone.

Of the 13 patients who had tumor response to the combination of Oxaliplatin and

5-fluorouracil/leucovorin, 5 were female and 8 were male, and responders included patients <65 years old and ≥65 years old. The small number of non-Caucasian participants made efficacy analyses in these populations uninterpretable.

Oxaliplatin Concentrate for Solution for Infusion (Oxitan) used in combination with infusional 5-fluorouracil/leucovorin, is indicated for

· Adjuvant treatment of stage III colon cancer in patients who have undergone complete resection of the primary tumor.

Treatment of advanced colorectal cancer.

Clinical Trials Experience

CONTRAINDICATIONS

to patients with a history of known allergy to oxaliplatin or other platinum compounds.

Oxaliplatin Concentrate for Solution for Infusion (Oxitan) should not be administered

pulmonary toxicities and hepatotoxicities can occur Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the

Serious adverse reactions including anaphylaxis and allergic reactions, neuropathy,

clinical trials of another drug and may not reflect the rates observed in practice. More than 1100 patients with stage II or III colon cancer and more than 4000 patients with advanced colorectal cancer have been treated in clinical studies with oxaliplatin either as a single agent or in combination with other medications. The most common adverse reactions in patients with stage II or III colon cancer receiving adjuvant therapy were peripheral sensory neuropathy, neutropenia, thrombocytopenia, anemia, nausea, increase in transaminases and alkaline phosphatase, diarrhea, emesis, fatigue, and stomatitis. The most common adverse reactions in previously untreated and treated patients were peripheral sensory neuropathies, fatique, neutropenia, nausea, emesis,

Combination Adjuvant Therapy with Oxaliplatin and Infusional 5-FU/LV in Patients with Colon Cancer

One thousand one hundred eight patients with stage II or III colon cancer, who had undergone complete resection of the primary tumor, were treated in a clinical study with oxaliplatin in combination with infusional 5-FU/LV. The incidence of grade 3 or 4 adverse events was 70% on the oxaliplatin combination arm, and 31% on the infusional 5-FU/LV arm. The adverse reactions in this trial are shown in the tables below. Discontinuation of treatment due to adverse events occurred in 15% of the patients receiving exaliplating and infusional 5-FU/LV. Both 5-FU/LV and oxaliplatin are associated with gastrointestinal or hematologic adverse events. When oxaliplatin is administered in combination with infusional 5-FU/LV, the incidence of these events is increased.

The incidence of death within 28 days of last treatment, regardless of causality, was 0.5% (n=6) in both the oxaliplatin combination and infusional 5-FU/LV arms, respectively. Deaths within 60 days from initiation of therapy were 0.3% (n=3) in both the oxaliplatin combination and infusional 5-FU/LV arms, respectively. On the oxaliplatin combination arm, 3 deaths were due to sepsis/neutropenic sepsis, 2 from intracerebral bleeding, and one from eosinophilic pneumonia. On the 5-FU/LV arm, one death was due to suicide, 2 from Stevens-Johnson syndrome (1 patient also had sepsis), 1 unknown cause, 1 anoxic cerebral infarction, and 1 probable abdominal aorta rupture.

The following table provides adverse events reported in the adjuvant therapy colon cancer clinical trial by body system and decreasing order of frequency in the oxaliplatin and infusional 5-FU/LV arm for events with overall incidences ≥5% and for NCI grade 3/4 events with incidences >1%.

Table 13 – Adverse Experiences Reported in Patients with Colon Cancer Receiving Adjuvant Treatment (>5% of all patients and with >1% NCI grade 3/4 events)

	Oxaliplatin + N=11		5-FU/LV N=1111		
Adverse Event (WHO/Pref)	All Grades (%)	Grade 3/4 (%)	All Grades (%)	Grade 3/4 (%)	
Any Event	100	70	99	31	
	Allergy/Immu	ınology			
Allergic Reaction	10	3	2	<1	
Co	onstitutional Syr	nptoms/Pain			
Fatigue	44	4	38	1	
Abdominal Pain	18	1	17	2	
	Dermatolog	y/Skin			
Skin Disorder	32	2	36	2	
Injection Site Reaction ¹	11	3	10	3	
	Gastrointe	stinal			
Nausea	74	5	61	2	
Diarrhea	56	11	48	7	
Vomiting	47	6	24	1	
Stomatitis	42	3	40	2	
Anorexia	13	1	8	<1	
	Fever/Infe	ction			
Fever	27	1	12	1	
Infection	25	4	25	3	
	Neurolo	gy			
Overall Peripheral Sensory Neuropathy	92	12	16	<1	

The following table provides adverse events reported in the adjuvant therapy colon cancer clinical trial by body system and decreasing order of frequency in the oxaliplatin and infusional 5-FU/LV arm for events with overall incidences ≥5% but with incidences <1% NCI grade 3/4 events.

Table 14 - Adverse Experiences Reported in Patients with Colon Cancer Receiving Adjuvant Treatment (>5% of all patients, but with <1% NCI grade 3/4 events) Oxaliplatin 5-FU/LV

	+ 5-FU/LV N=1108	N=1111
Adverse Event (WHO/Pref)	All Grades (%)	All Grades (%)
Allerg	y/lmmunology	
Rhinitis	6	8
Constitutional Syn	nptoms/Pain/Ocular/Vis	sual
Epistaxis	16	12
Weight Increase	10	10
Conjunctivitis	9	15
Headache	7	5
Dyspnea	5	3
Pain	5	5
Lacrimation Abnormal	4	12
Derm	natology/Skin	
Alopecia	30	28
Gas	trointestinal	
Constipation	22	19
Taste Perversion	12	8
Dyspepsia	8	5
N	Metabolic	
Phosphate Alkaline Increased	42	20
N	leurology	
Sensory Disturbance	8	1
Although specific events can vary, to	he overall frequency of	f adverse events w

similar in men and women and in patients <65 and >65 years. However, the following grade 3/4 events were more common in females: diarrhea, fatigue, granulocytopenia nausea, and vomiting. In patients >65 years old, the incidence of grade 3/4 diarrhea and granulocytopenia was higher than in younger patients. Insufficient subgroup sizes prevented analysis of safety by race. The following additional adverse events were reported in >2% and <5% of the patients in the oxaliplatin and infusional 5-FU/ LV combination arm (listed in decreasing order of frequency): pain, leukopenia, weight decrease, coughing.

The number of patients who developed secondary malignancies was similar; 62 in the Oxaliplatin combination arm and 68 in the infusional 5-fluorouracil/leucovorin arm. An exploratory analysis showed that the number of deaths due to secondary malignancies was 1.96% in the Oxaliplatin combination arm and 0.98% in infusional 5-fluorouracil/ leucovorin arm. In addition, the number of cardiovascular deaths was 1.4% in the Oxaliplatin combination arm as compared to 0.7% in the infusional 5-fluorouracil/leucovorin arm. Clinical significance of these findings is unknown

Patients Previously Untreated for Advanced Colorectal Cancer

Two hundred and fifty-nine patients were treated in the oxaliplatin and 5-FU/LV combination arm of the randomized trial in patients previously untreated for advanced colorectal cancer. The adverse event profile in this study was similar to that seen in other studies and the adverse reactions in this trial are shown in the tables

Both 5-FU and oxaliplatin are associated with gastrointestinal and hematologic adverse events. When oxaliplatin is administered in combination with 5-FU, the incidence of these events is increased

The incidence of death within 30 days of treatment in the previously untreated for advanced colorectal cancer study, regardless of causality, was 3% with the oxaliplatin and 5-FU/LV combination, 5% with irinotecan plus 5-FU/LV, and 3% with oxaliplatin plus irinotecan. Deaths within 60 days from initiation of therapy were 2.3% with the exaliplatin and 5-FU/LV combination, 5.1% with irinotecan plus 5-FU/LV, and 3.1% with oxaliplatin plus irinotecan.

The following table provides adverse events reported in the previously untreated for advanced colorectal cancer study by body system and decreasing order of frequency in the oxaliplatin and 5-FU/LV combination arm for events with overall incidences ≥5% and for grade 3/4 events with incidences $\geq 1\%$.

Table 15 – Adverse Experiences Reported in Patients with Colon Cancer Receiving Adjuvant Treatment (>5% of all patients, but with <1% NCl grade 3/4 events)

	Oxalipi 5-FU N=2	I/LV	Irinote 5-FU N=2	/LV	Oxaliplatin + irinotecan N=258	
	All	.55	All	.50	All	1
Adverse Event (WHO/Pref)	Grades	Grade 3/4 (%)	Grades (%)	Grade 3/4 (%)	Grades	Grade 3/4 (%
Λ Γt	(%) 99	00	(, ,	70	(%)	70
Any Event		82 ergy/lmm	98	70	99	76
Hypersensitivity	12	2	5	0	6	1
Пурегоепошицу		∟ ∠ Cardiova	_	0	0	
Thrombosis	6	5	6	6	3	3
Hypotension	5	3	6	3	4	3
	stitutional	_	-	-	al .	
Fatigue	70	7	58	11	66	16
Abdominal Pain	29	8	31	7	39	10
Myalgia	14	2	6	0	9	2
Pain	7	1	5	1	6	1
Vision Abnormal	5	0	2	1	6	1
Neuralgia	5	0	0	0	2	1
rveuraigia		ermatolo				<u> </u>
Skin Reaction - hand/ foot	7	1	2	1	1	0
Injection site reaction	6	0	1	0	4	1
joodaan alta raadaan		Gastroint	estinal			
Nausea	71	6	67	15	83	19
Diarrhea	56	12	65	29	76	25
Vomiting	41	4	43	13	64	23
Stomatitis	38	0	25	1	19	1
Anorexia	35	2	25	4	27	5
Constipation	32	4	27	2	21	2
Diarrhea - colostomy	13	2	16	7	16	3
Gastrointestinal NOS*	5	2	4	2	3	2
Gastrointestinal NOS			Infection		3	
Infection no ANC**	10	4	5	1	7	2
Infection low ANC**	8	8	12	11	9	8
Lymphopenia	6	2	4	1	5	2
Febrile Neutropenia	4	4	15	14	12	11
rebrile Neutropenia	Hepatic/Me				12	- 11
I li maneli rasmia					10	1 2
Hyperglycemia	14	2	11	3	12	3
Hypokalemia	11	3	7		6	2
Dehydration	9	5	16	11	14	7
Hypoalbuminemia	8	0	5	2	9	1
Hyponatremia	8	2	7	4	4	1
Urinary Frequency	5	1	2	1	3	1
0 "11 "	1 00	Neurol		_	- 00	_
Overall Neuropathy	82	19	18	2	69	7
Paresthesias	77	18	16	2	62	6
Pharyngo-laryngeal Dysesthesias	38	2	1	0	28	1
Neuro-sensory	12	1	2	0	9	1
Neuro NOS*	1	0	1	0	1	0
		Pulmo				
Cough	35	1	25	2	17	1
Dyspnea	18	7	14	3	11	2
Hiccups	5	1	2	0	3	2
Not otherwise specifie	d					
* Not otherwise specifie ** Absolute neutrophil co The following table pro advanced colorectal car	ount vides advei					

with incidences < 1% NCI grade 3/4 events. Table 16 - Adverse Experiences Reported in Patients Previously Untreated for Advanced Colorectal Cancer Clinical Trial (>5% of all patients and with >1% NCI

Adverse Event (WHO/	Oxaliplatin + 5-FU/LV N=259	Irinotecan + 5-FU/LV N=256	Oxaliplatin + irinotecan N=258
Pref)	All Grades (%)	All Grades (%)	All Grades (%)
	Allergy/Immun	ology	`
Rash	11	4	7
Rhinitis Allergic	10	6	6
	Cardiovascu	lar	
Edema	15	13	10
Constitu	utional Symptoms/F	ain/Ocular/Visual	
Headache	13	6	9
Weight Loss	11	9	11
Epistaxis	10	2	2
Tearing	9	1	2
Rigors	8	2	7
Dysphasia	5	3	3
Sweating	5	6	12
Arthralgia	5	5	8
	Dermatology/	Skin	
Alopecia	38	44	67
Flushing	7	2	5
Pruritus	6	4	2
Dry Skin	6	2	5

Reason for the Artwork*: Change in text matter Dimension: 500 x 400 mm Artwork No.: 7220131933 Supersede No.: 7220131484 Checked Prepard Approved By $\mathbf{B}\mathbf{y}$ By Country PDD PDD **PDD Production** QA I&D RA $\mathbf{Q}\mathbf{M}$ RA

Form No.: SOP/PDD/003-03 Rev. 05

CONFIDENTIAL

* Attached separate sheet if required

NA

Size: 500 x 400 mm

	Gastrointesti	nal	
Taste Perversion	14	6	8
Dyspepsia	12	7	5
Flatulence	9	6	5
Mouth Dryness	5	2	3
	Hematology/Infe	ection	
Fever normal ANC*	16	9	9
Нера	tic/Metabolic/Labo	oratory/Renal	
Hypocalcemia	7	5	4
Elevated Creatinine	4	4	5
	Neurology		
Insomnia	13	9	11
Depression	9	5	7
Dizziness	8	6	10
Anxiety	5	2	6

* Absolute neutrophil coun

Adverse reactions were similar in men and women and in patients <65 and ≥65 years but older patients may have been more susceptible to diarrhea, dehydration, hypokalemia, leukopenia, fatigue, and syncope. The following additional adverse events, at least possibly related to treatment and potentially important, were reported in >2% and <5% of the patients in the oxaliplatin and 5-FU/LV combination arm (listed in decreasing order of frequency): metabolic, pneumonitis, catheter infection, vertigo, prothrombin time, pulmonary, rectal bleeding, dysuria, nail changes, chest pain, rectal pain, syncope, hypertension, hypoxia, unknown infection, bone pain, pigmentation changes, and

Previously Treated Patients with Advanced Colorectal Cancer

Four hundred and fifty patients (about 150 receiving the combination of oxaliplatin and 5-FU/LV) were studied in a randomized trial in patients with refractory and relapsed colorectal cancer. The adverse event profile in this study was similar to that seen in other studies and the adverse reactions in this trial are shown in the tables below.

Thirteen percent of patients in the oxaliplatin and 5-FU/LV combination arm and 18% in the 5-FU/LV arm of the previously treated study had to discontinue treatment because of adverse effects related to gastrointestinal, or hematologic adverse events, or neuropathies. Both 5-FU and oxaliplatin are associated with gastrointestinal and hematologic adverse events. When oxaliplatin is administered in combination with 5-FU, the incidence of these events is increased.

The incidence of death within 30 days of treatment in the previously treated study regardless of causality, was 5% with the oxaliplatin and 5-FU/LV combination, 8% with oxaliplatin alone, and 7% with 5-FU/LV. Of the 7 deaths that occurred on the oxaliplatin and 5-FU/LV combination arm within 30 days of stopping treatment, 3 may have been treatment related, associated with gastrointestinal bleeding or dehydration.

The following table provides adverse events reported in the previously treated study by body system and in decreasing order of frequency in the oxaliplatin and 5-FU/LV combination arm for events with overall incidences >5% and for grade 3/4 events with incidences ≥ 1%. This table does not include hematologic and blood chemistry abnormalities; these are shown separately below

Table 17 - Adverse Experiences Reported in Patients Previously Untreated for Advanced Colorectal Cancer Clinical Trial (> 5% of all patients but with < 1% NCI

	5-FU/LV (N=142) Oxalipla (N=15)					
Adverse Event (WHO/Pref)	All Grades (%)	Grade 3/4 (%)	All Grades (%)	Grade 3/4 (%)	All Grades (%)	Grade 3/4 (%)
Any Event	98	41	100	46	99	73
		Cardiova	scular			
Dyspnea	11	2	13	7	20	4
Coughing	9	0	11	0	19	1
Edema	13	1	10	1	15	1
Thromboembolism	4	2	2	1	9	8
Chest Pain	4	1	5	1	8	1
	Constitu	ıtional Sy	mptoms/P	Pain		
Fatigue	52	6	61	9	68	7
Back Pain	16	4	11	0	19	3
Pain	9	3	14	3	15	2
	D	ermatolo	gy/Skin			
Injection Site Reaction	5	1	9	0	10	3
	(estinal			
Diarrhea	44	3	46	4	67	11
Nausea	59	4	64	4	65	11
Vomiting	27	4	37	4	40	9
Stomatitis	32	3	14	0	37	3
Abdominal Pain	31	5	31	7	33	4
Anorexia	20	1	20	2	29	3
Gastroesophageal Reflux	3	0	1	0	5	2
	Her	natology/	Infection			
Fever	23	1	25	1	29	1
Febrile Neutropenia	1	1	0	0	6	6
	Hepatic/Me	etabolic/L	aboratory	Renal		
Hypokalemia	3	1	3	2	9	4
Dehydration	6	4	5	3	8	3
		Neurol	ogy			
Neuropathy	17	0	76	7	74	7
Acute	10	0	65	5	56	2
Persistent	9	0	43	3	48	6

grade 3/4 events. Table 18 - Adverse Experiences Reported in Previously Treated Colorectal Can-

cer Clinical Trial (>5% of all patients but with < 1% NCI grade 3/4 events)

Adverse Event	5-FU/LV (N=142)	Oxaliplatin (N=153)	Oxaliplatin+ 5-FU/LV (N=150)
(WHO/Pref)	All Grades (%) All Grades (%)		All Grades (%)
	Allergy/Imm	nunology	
Rhinitis	4	6	15
Allergic Reaction	1	3	10
Rash	5	5	9
	Cardiova	scular	
Peripheral Edema	11	5	10
Cons	stitutional Sympton	ns/Pain/Ocular/Visu	al
Headache	8	13	17
Arthralgia	10	7	10
Epistaxis	1	2	9
Abnormal Lacrimation	6	1	7
Rigors	6	9	7
	Dermatolo	gy/Skin	
Hand-Foot Syndrome	13	1	11
Flushing	2	3	10
Alopecia	3	3	7
	Gastroint	estinal	
Constipation	23	31	32
Dyspepsia	10	7	14
Taste Perversion	1	5	13

Reason for the Artwork*: Change in text matter

Checked

By

PDD

Artwork No.: 7220131933

Prepard

 $\mathbf{B}\mathbf{y}$

PDD

		_	
Mucositis	10	2	7
Flatulence	6	3	5
	Hepatic/Metabolic/L	aboratory/Renal	
Hematuria	4	0	6
Dysuria	1	1	6
	Neurol	ogy	
Dizziness	8	7	13
Insomnia	4	11	9
	Pulmoi	nary	
Upper Resp Tract			
Infection	4	7	10
Pharyngitis	10	2	9
Hiccup	0	2	5

Adverse reactions were similar in men and women and in patients <65 and >65 years, but older patients may have been more susceptible to dehydration, diarrhea, hypokalemia and fatigue. The following additional adverse events, at least possibly related to treatment and potentially important, were reported in \geq 2% and <5% of the patients in the oxaliplatin and 5-FU/LV combination arm (listed in decreasing order of frequency): anxiety, myalgia, erythematous rash, increased sweating, conjunctivitis, weight decrease. dry mouth, rectal hemorrhage, depression, ataxia, ascites, hemorrhoids, muscle weakness, nervousness, tachycardia, abnormal micturition frequency, dry skin, pruritus hemoptysis, purpura, vaginal hemorrhage, melena, somnolence, pneumonia, proctitis, involuntary muscle contractions, intestinal obstruction, gingivitis, tenesmus, hot flashes, enlarged abdomen, urinary incontinence.

matologic changes

The following tables list the hematologic changes occurring in ≥5% of patients, based on laboratory values and NCI grade, with the exception of those events occurring in adjuvant patients and anemia in the patients previously untreated for advanced colorectal cancer, which are based on AE reporting and NCI grade alone.

19- Adverse Hematologic Experiences in Patients with Colon Cancer Receiving Adjuvant Therapy (>5% of patients)

Hematology	Oxaliı + 5-FU/LV		5-FU/LV (N=1111)		
Parameter	All Grades (%)	Grade 3/4 (%)	All Grades (%)	Grade 3/4 (%)	
Anemia	76	1	67	<1	
Neutropenia	79	41	40	5	
Thrombocytopenia	77	2	19	<1	

	Oxaliplatin + 5-FU/LV N=259		Irinotecan + 5-FU/ LV N=256		Oxaliplatin + irinotecan N=258	
Hematology Parameter	All Grades (%)	Grade 3/4 (%)	All Grades (%)	Grade 3/4 (%)	All Grades (%)	Grade 3/4 (%)
Anemia	27	3	28	4	25	3
Leukopenia	85	20	84	23	76	24
Neutropenia	81	53	77	44	71	36
Thrombocytopenia	71	5	26	2	44	4

Table 21 – Adverse Hematologic Experiences in Previously Treated Patients (>5%

or patients)						
	5-FU/LV (N=142)		Oxalip (N=1		Oxaliplatin + 5-FU/LV (N=150)	
Hematology Parameter	All Grades (%)	Grade 3/4 (%)	All Grades (%)	Grade 3/4 (%)	All Grades (%)	Grade 3/4 (%)
Anemia	68	2	64	1	81	2
Leukopenia	34	1	13	0	76	19
Neutropenia	25	5	7	0	73	44
Thrombocytopenia	20	0	30	3	64	4

Thrombocytopenia and Bleeding

Thrombocytopenia was frequently reported with the combination of oxaliplatin and infusional 5-FU/LV. The incidence of all hemorrhagic events in the adjuvant and previously treated patients was higher on the oxaliplatin combination arm compared to the infusional 5-FU/LV arm. These events included gastrointestinal bleeding, hematuria, and epistaxis. In the adjuvant trial, two patients died from intracerebral hemorrhages.

The incidence of grade 3/4 thrombocytopenia was 2% in adjuvant patients with colon cancer. In patients treated for advanced colorectal cancer the incidence of grade 3/4 thrombocytopenia was 3-5%, and the incidence of these events was greater for the combination of oxaliplatin and 5-FU/LV over the irinotecan plus 5-FU/LV or 5-FU/LV control groups. Grade 3/4 gastrointestinal bleeding was reported in 0.2% of adjuvant patients receiving oxaliplatin and 5-FU/LV. In the previously untreated patients, the incidence of epistaxis was 10% in the oxaliplatin and 5-FU/LV arm, and 2% and 1%, respectively, in the irinotecan plus 5-FU/LV or irinotecan plus oxaliplatin arms.

Neutropenia was frequently observed with the combination of oxaliplatin and 5-FU/LV, with grade 3 and 4 events reported in 29% and 12% of adjuvant patients with colon cancer, respectively. In the adjuvant trial, 3 patients died from sepsis/neutropenic sepsis. Grade 3 and 4 events were reported in 35% and 18% of the patients previously untreated for advanced colorectal cancer, respectively. Grade 3 and 4 events were reported in 27% and 17% of previously treated patients, respectively. In adjuvant patients the incidence of either febrile neutropenia (0.7%) or documented infection with concomitant grade 3/4 neutropenia (1.1%) was 1.8% in the oxaliplatin and 5-FU/LV arm. The incidence of febrile neutropenia in the patients previously untreated for advanced colorectal cancer was 15% (3% of cycles) in the irinotecan plus 5-FU/LV arm and 4% (less than 1% of cycles) in the oxaliplatin and 5-FU/LV combination arm. Additionally, in this same population, infection with grade 3 or 4 neutropenia was 12% in the irinotecan plus 5-FU/LV and 8% in the oxaliplatin and 5-FU/LV combination. The incidence of febrile neutropenia in the previously treated patients was 1% in the 5-FU/LV arm and 6% (less than 1% of cycles) in the oxaliplatin and 5-FU/LV combination arm.

In patients receiving the combination of oxaliplatin plus infusional 5-FU/LV for adjuvant treatment for colon cancer the incidence of grade 3/4 nausea and vomiting was greater than in those receiving infusional 5-FU/LV alone. In patients previously untreated for advanced colorectal cancer receiving the combination of oxaliplatin and 5-FU/LV, the incidence of grade 3 and 4 vomiting and diarrhea was less compared to irinotecan plus 5-FU/LV controls. In previously treated patients receiving the combination of oxaliplatin and 5-FU/LV, the incidence of grade 3 and 4 nausea, vomiting, diarrhea, and mucositis/ stomatitis increased compared to 5-FU/LV controls.

The incidence of gastrointestinal adverse events in the previously untreated and previously treated patients appears to be similar across cycles. Premedication with antiemetics, including 5-HT₃ blockers, is recommended. Diarrhea and mucositis may be exacerbated by the addition of oxaliplatin to 5-FU/LV, and should be managed with appropriate supportive care. Since cold temperature can exacerbate acute neurological symptoms, ice (mucositis prophylaxis) should be avoided during the infusion of

Oxaliplatin did not increase the incidence of alopecia compared to 5-FU/LV alone. No complete alopecia was reported. The incidence of grade 3/4 skin disorders was 2% in both the oxaliplatin plus infusional 5-FU/LV and the infusional 5-FU/LV alone arms in the adjuvant colon cancer patients. The incidence of hand-foot syndrome in patients previously untreated for advanced colorectal cancer was 2% in the irinotecan plus 5-FU/ LV arm and 7% in the oxaliplatin and 5-FU/LV combination arm. The incidence of handfoot syndrome in previously treated patients was 13% in the 5-FU/LV arm and 11% in the oxaliplatin and 5-FU/LV combination arm.

Intravenous Site Reactions

Extravasation, in some cases including necrosis, has been reported. Injection site reaction, including redness, swelling, and pain, has been reported.

Dimension: 500 x 400 mm

Supersede No.: 7220131484

I&D RA

Anticoagulation and Hemorrhage

There have been reports while on study and from post-marketing surveillance of prolonged prothrombin time and INR occasionally associated with hemorrhage in patients who received oxaliplatin plus 5-fluorouracil/leucovorin while on anticoagulants. Patients receiving oxaliplatin plus 5-fluorouracil/leucovorin and requiring oral anticoagulants may require closer monitoring.

About 5-10% of patients in all groups had some degree of elevation of serum creatinine. The incidence of grade 3/4 elevations in serum creatinine in the oxaliplatin and 5-FU/ LV combination arm was 1% in the previously treated patients. Serum creatinine measurements were not reported in the adjuvant trial.

Hepatotoxicity (defined as elevation of liver enzymes) appears to be related to oxaliplatin combination therapy. The following tables list the clinical chemistry changes associated with hepatic toxicity occurring in ≥5% of patients, based on adverse events reported and NCI CTC grade for patients previously untreated for advanced colorectal cancer, laboratory values, and NCI CTC grade for previously treated patients.

Table 22 - Adverse Hepatic Experiences in Patients with Stage II or III Colon Cancer Receiving Adjuvant Therapy (>5% of patients)

Hanatia Daramatar	Oxaliı + 5-FU/LV		5-FU/LV (N=1111)		
Hepatic Parameter	All Grades (%)	Grade 3/4 (%)	All Grades (%)	Grade 3/4 (%)	
Increase in Transaminases	57	2	34	1	
ALP Increased	42	<1	20	<1	
Bilirubinaemia	20	4	20	5	

Untreated for Advanced Colorectal Cancer (>5% of patients)

	Oxaliplatin + 5-FU/LV N=259 All Grade Grades 3/4		Irinotecan + 5-FU/LV N=256		Oxaliplatin + irinotecan N=258	
			All Grades	Grade 3/4	All Grades	Grade 3/4
Clinical Chemistry	(%)	(%)	(%)	(%)	(%)	(%)
ALT (SGPT-ALAT)	6	1	2	0	5	2
AST (SGOT-ASAT)	17	1	2	1	11	1
Alkaline Phosphatase	16	0	8	0	14	2
Total Bilirubin	6	1	3	1	3	2

Table 24 – Adverse Hepatic - Clinical Chemistry Experience in Previously Treated

Olisiaal Ohamiataa	5-FU/LV (N=142)		Oxaliplatin (N=153)		Oxaliplatin + 5-FU/LV (N=150)	
Clinical Chemistry	All Grades (%)	Grade 3/4 (%)	All Grades (%)	Grade 3/4 (%)	All Grades (%)	Grade 3/4 (%)
ALT (SGPT-ALAT)	28	3	36	1	31	0
AST (SGOT-ASAT)	39	2	54	4	47	0
Total Bilirubin	22	6	13	5	13	1

The incidence of thromboembolic events in adjuvant patients with colon cancer was 6% (1.8% grade 3/4) in the infusional 5-FU/LV arm and 6% (1.2% grade 3/4) in the

oxaliplatin and infusional 5-FU/LV combined arm, respectively. The incidence was 6 and 9% of the patients previously untreated for advanced colorectal cancer and previously treated patients in the oxaliplatin and 5-FU/LV combination arm, respectively.

Postmarketing Experience

The following adverse reactions have been identified during post-approval use of Oxaliplatin. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Body as a whole:

angioedema, anaphylactic shock Central and peripheral nervous system disorders:

Liver and Gastrointestinal system disorders:

loss of deep tendon reflexes, dysarthria, Lhermitte's sign, cranial nerve palsies, fasciculations, convulsion

severe diarrhea/vomiting resulting in hypokalemia, colitis (including Clostridium difficile diarrhea), metabolic acidosis: ileus: intestinal obstruction, pancreatitis: veno-occlusive disease of liver also known as sinusoidal obstruction syndrome, and perisinusoidal fibrosis which rarely may progress.

Hearing and vestibular system disorders:

Platelet, bleeding, and clotting disorders:

immuno-allergic thrombocytopenia prolongation of prothrombin time and of INR in

Red Blood Cell disorders:

hemolytic uremic syndrome, immuno-allergic hemolytic anemia Renal disorders:

Acute tubular necrosis, acute interstitial nephritis and acute renal failure.

pulmonary fibrosis, and other interstitial lung diseases (sometimes fatal)

Vision disorders: decrease of visual acuity, visual field disturbance, optic neuritis and transient vision loss (reversible following therapy discontinuation)

DRUG INTERACTIONS

No specific cytochrome P-450-based drug interaction studies have been conducted. No pharmacokinetic interaction between 85 mg/m² Oxaliplatin and 5-fluorouracil/leucovorin has been observed in patients treated every 2 weeks. Increases of 5-fluorouracil plasma concentrations by approximately 20% have been observed with doses of 130 mg/m² Oxaliplatin dosed every 3 weeks. Because platinum-containing species are eliminated primarily through the kidney, clearance of these products may be decreased by co-administration of potentially nephrotoxic compounds; although, this has not been specifically studied.

PRECAUTIONS AND WARNINGS

Allergic Reactions

Grade 3/4 hypersensitivity, including anaphylactic/anaphylactoid reactions, to Oxaliplatin has been observed in 2-3% of colon cancer patients. These allergic reactions which can be fatal, can occur within minutes of administration and at any cycle, and were similar in nature and severity to those reported with other platinum-containing compounds, such as rash, urticaria, erythema, pruritus, and, rarely, bronchospasm and hypotension. The symptoms associated with hypersensitivity reactions reported in the previously untreated patients were urticaria, pruritus, flushing of the face, diarrhea associated with oxaliplatin infusion, shortness of breath, bronchospasm, diaphoresis, chest pains, hypotension, disorientation and syncope. These reactions are usually managed with standard epinephrine, corticosteroid, antihistamine therapy, and require discontinuation of therapy. Rechallenge is contraindicated in these. Drug-related deaths associated with platinum compounds from anaphylaxis have been reported.

 $\mathbf{Q}\mathbf{M}$

Oxalinlatin is associated with two types of neuropathy:

Country

* Attached separate sheet if required

An acute, reversible, primarily peripheral, sensory neuropathy that is of early onset, occurring within hours or one to two days of dosing, that resolves within 14 days, and that frequently recurs with further dosing. The symptoms may be precipitated or exacerbated by exposure to cold temperature or cold objects and they usually present as transient paresthesia, dysesthesia and hypoesthesia in the hands, feet, perioral area. or throat. Jaw spasm, abnormal tongue sensation, dysarthria, eye pain, and a feeling of chest pressure have also been observed. The acute, reversible pattern of sensory

neuropathy was observed in about 56% of study patients who received Oxaliplatin with 5-FU/LV. In any individual cycle acute neurotoxicity was observed in approximately 30% of patients. In adjuvant patients the median cycle of onset for grade 3 peripheral sensory neuropathy was 9 in the previously treated patients the median number of cycles administered on the oxaliplatin with 5 FU/LV combination arm was 6.

An acute syndrome of pharyngolaryngeal dysesthesia seen in 1-2% (grade 3/4) of patients previously untreated for advanced colorectal cancer, and the previously treated patients, is characterized by subjective sensations of dysphagia or dyspnea, without any larvngospasm or bronchospasm (no stridor or wheezing), Ice (mucositis prophylaxis) should be avoided during the infusion of Oxaliplatin because cold temperature can exacerbate acute neurological symptoms.

A persistent (>14 days), primarily peripheral, sensory neuropathy that is usually characterized by paresthesias, dysesthesias, hypoesthesias, but may also include deficits in proprioception that can interfere with daily activities (e.g., writing, buttoning, swallowing, and difficulty walking from impaired proprioception). These forms of neuropathy occurred in 48% of the study patients receiving Oxaliplatin with 5-FU/ LV. Persistent neuropathy can occur without any prior acute neuropathy event. The majority of the patients (80%) who developed grade 3 persistent neuropathy progressed from prior Grade 1 or 2 events. These symptoms may improve in some patients upon discontinuation of Oxaliplatin

In the adjuvant colon cancer trial, neuropathy was graded using a prelisted module derived from the Neuro-Sensory section of the National Cancer Institute Common Toxicity Criteria (NCI CTC) scale, Version 1, as follows:

Table 25: NCI CTC Grading for Neuropathy in Adjuvant Patients

NCI Grade	Definition			
Grade 0 No change or none				
Grade 1	d paresthesias, loss of deep tendon reflexes			
Grade 2	Mild or moderate objective sensory loss, moderate paresthesias			
Grade 3	Severe objective sensory loss or paresthesias that interfere with function			
Grade 4 Not applicable				

Peripheral sensory neuropathy was reported in adjuvant patients treated with the Oxaliplatin combination with a frequency of 92% (all grades) and 13% (grade 3). At the 28-day follow-up after the last treatment cycle, 60% of all patients had any grade (Grade 1=40%, Grade 2=16%, Grade 3=5%) peripheral sensory neuropathy decreasing to 39% at 6 months follow-up (Grade 1=31%, Grade 2=7%, Grade 3=1%) and 21% at 18 months of follow-up (Grade 1=17%, Grade 2=3%, Grade 3=1%).

In the advanced colorectal cancer studies, neuropathy was graded using a studyspecific neurotoxicity scale, which was different from the NCI CTC scale, Version 2.0.

Grading Scale for Paresthesias/Dysesthesias in Advanced Colorectal Cancer

Grade	Definition			
Grade 1	Resolved and did not interfere with functioning			
Grade 2	Interfered with function but not daily activities			
Grade 3 Pain or functional impairment that interfered with daily activities				
Grade 4 Persistent impairment that is disabling or life-threatening				
Overall, neuropathy was reported in patients previously untreated for advanced				

colorectal cancer in 82% (all grades) and 19% (grade 3/4), and in the previously treated patients in 74% (all grades) and 7% (grade 3/4) events. Information regarding reversibility of neuropathy was not available from the trial for patients who had not been previously treated for colorectal cancer.

Reversible Posterior Leukoencephalopathy Syndrome

Reversible Posterior Leukoencephalopathy Syndrome (RPLS, also known as PRES, Posterior Reversible Encephalopathy Syndrome) has been observed in clinical trials < 0.1%) and postmarketing experience. Signs and symptoms of RPLS could be headache, altered mental functioning, seizures, abnormal vision from blurriness to blindness, associated or not with hypertension. Diagnosis of RPLS is based upon confirmation by brain imaging.

Oxaliplatin has been associated with pulmonary fibrosis (<1% of study patients), which may be fatal. The combined incidence of cough and dyspnea was 7.4% (any grade) and <1% (grade 3) with no grade 4 events in the Oxaliplatin plus infusional 5-FU/LV arm compared to 4.5% (any grade) and no grade 3 and 0.1% grade 4 events in the infusional 5-FU/LV alone arm in adjuvant colon cancer patients. In the study, one patient died from eosinophilic pneumonia in the Oxaliplatin combination arm. The combined incidence of cough, dyspnea, and hypoxia was 43% (any grade) and 7% (grade 3 and 4) in the Oxaliplatin plus 5-FU/LV arm compared to 32% (any grade) and 5% (grade 3 and 4) in the Irinotecan plus 5-FU/LV arm of unknown duration for patients with previously untreated colorectal cancer. In case of unexplained respiratory symptoms such as nonproductive cough, dyspnea, crackles, or radiological pulmonary infiltrates, Oxaliplatin should be discontinued until further pulmonary investigation excludes interstitial lung disease or pulmonary fibrosis.

Hepatotoxicity as evidenced in the adjuvant study, by increase in transaminases (57% vs. 34%) and alkaline phosphatase (42% vs. 20%) was observed more commonly in the Oxaliplatin combination arm. The incidence of increased bilirubin was similar on both arms. Changes noted on liver biopsies include: peliosis, nodular regenerative hyperplasia or sinusoidal alterations, perisinusoidal fibrosis, and veno-occlusive lesions. Hepatic vascular disorders should be considered and if appropriate, should be investigated in case of abnormal liver function test results or portal hypertension, which cannot be explained by liver metastases.

Oxaliplatin may cause fetal harm when administered to a pregnant woman. There are no adequate and well-controlled studies of Oxaliplatin in pregnant women. Women of childbearing potential should be advised to avoid becoming pregnant while receiving treatment with Oxaliplatin

Recommended Laboratory Tests

Standard monitoring of the white blood cell count with differential, hemoglobin, platelet count, and blood chemistries (including ALT, AST, bilirubin and creatinine) is recommended before each Oxaliplatin cycle.

There have been reports while on study and from post-marketing surveillance of prolonged prothrombin time and INR occasionally associated with hemorrhage in patients who received Oxaliplatin plus 5-fluorouracil/leucovorin while on anticoagulants. Patients receiving Oxaliplatin plus 5-fluorouracil/leucovorin and requiring oral DOSAGE AND ADMINISTRATION anticoagulants may require closer monitoring.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term animal studies have not been performed to evaluate the carcinogenic potential of oxaliplatin. Oxaliplatin was not mutagenic to bacteria (Ames test) but was mutagenic to mammalian cells in vitro (L5178Y mouse lymphoma assay). Oxaliplatin was clastogenic both in vitro (chromosome aberration in human lymphocytes) and in vivo (mouse bone marrow micronucleus assay).

In a fertility study, male rats were given Oxaliplatin at 0, 0.5, 1, or 2 mg/kg/day for five days every 21 days for a total of three cycles prior to mating with females that received two cycles of Oxaliplatin on the same schedule. A dose of 2 mg/kg/day (less than one-seventh the recommended human dose on a body surface area basis) did not affect pregnancy rate, but caused developmental mortality (increased early resorptions, decreased live fetuses, decreased live births) and delayed growth (decreased feta Testicular damage, characterized by degeneration, hypoplasia, and atrophy, was

observed in dogs administered Oxaliplatin at 0.75 mg/kg/day x 5 days every 28 days for three cycles. A no effect level was not identified. This daily dose is approximately onesixth of the recommended human dose on a body surface area basis

Use in Specific Populations

Based on direct interaction with DNA, Oxaliplatin may cause fetal harm when administered to a pregnant woman. There are no adequate and well-controlled studies of Oxaliplatin in pregnant women. Reproductive toxicity studies in rats demonstrated adverse effects on fertility and embryo-fetal development at maternal doses that were below the recommended human dose based on body surface area. If this drug is used

during pregnancy or if the patient becomes pregnant while taking this drug, the patient should be apprised of the potential hazard to the fetus. Women of childbearing potential should be advised to avoid becoming pregnant and use effective contraception while receiving treatment with Oxaliplatin.

Pregnant rats were administered Oxaliplatin at less than one-tenth the recommended human dose based on body surface area during gestation days 1-5 (pre-implantation), 6-10, or 11-16 (during organogenesis). Oxaliplatin caused developmental mortality (increased early resorptions) when administered on days 6-10 and 11-16 and adversely affected fetal growth (decreased fetal weight, delayed ossification) when administered on days 6-10. Administration of Oxaliplatin to male and female rats prior to mating resulted in 97% post-implantation loss in animals that received approximately oneseventh the recommended human dose based on the body surface area.

It is not known whether Oxaliplatin or its derivatives are excreted in human milk. Because many drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants from Oxaliplatin, a decision should be made whether to discontinue nursing or delay the use of the drug, taking into account the importance of the drug to the mother.

Pediatric Use

The effectiveness of Oxaliplatin in children has not been established. Oxaliplatin has been tested in 2 Phase I and 2 Phase II trials in 159 patients ages 7 months to 22 years with solid tumors (see below) and no significant activity observe

In a Phase I/II study. Oxaliplatin was administered as a 2-hour intravenous infusion on days 1, 8 and 15 every 4 weeks (1 cycle), for a maximum of 6 cycles, to 43 patients with refractory or relapsed malignant solid tumors, mainly neuroblastoma and osteosarcoma. Twenty eight pediatric patients in the Phase I study received Oxaliplatin at 6 dose levels starting at 40 mg/m 2 with escalation to 110 mg/m 2 . The dose limiting toxicity (DLT) was sensory neuropathy at the 110 mg/m² dose. Fifteen patients received Oxaliplatin at a dose of 90 mg/m² intravenous in the Phase II portion of the study. At this dose, paresthesia (60%, G3/4: 7%), fever (40%, G3/4: 7%) and thrombocytopenia (40%, G3/4: 27%) were the main adverse reactions. No responses were observed.

In a second Phase I study. Oxaliplatin was administered to 26 pediatric patients as a 2-hour intravenous infusion on day 1 every 3 weeks (1 cycle) at 5 dose levels starting at 100 mg/m² with escalation to 160 mg/m², for a maximum of 6 cycles. In a separate cohort, Oxaliplatin 85 mg/m² was administered on day 1 every 2 weeks, for a maximum of 9 doses. Patients had metastatic or unresectable solid tumors mainly neuroblastoma and ganglioneuroblastoma. No responses were observed. The DLT was sensory neuropathy at the 160 mg/m² dose. Based on these studies, Oxaliplatin 130 mg/m² as a 2-hour intravenous infusion on day 1 every 3 weeks (1 cycle) was used in subsequent Phase II studies. A dose of 85 mg/m 2 on day 1 every 2 weeks was also found to be

In one Phase II study, 43 pediatric patients with recurrent or refractory embryonal CNS tumors received Oxaliplatin 130 mg/m² every 3 weeks for a maximum of 12 months in absence of progressive disease or unacceptable toxicity. In patients < 10 kg the Oxaliplatin dose used was 4.3 mg/kg. The most common adverse reactions reported were leukopenia (67%, G3/4: 12%), anemia (65%, G3/4: 5%), thrombocytopenia (65%, G3/4: 26%), vomiting (65%, G3/4: 7%), neutropenia (58%, G3/4: 16%) and sensory neuropathy (40%, G3/4: 5%). One partial response was observed

In a second Phase II study, 47 pediatric patients with recurrent solid tumors, including Ewing sarcoma or peripheral PNET, osteosarcoma, rhabdomyosarcoma and neuroblastoma, received Oxaliplatin 130 mg/m² every 3 weeks for a maximum of 12 months or 17 cycles. In patients < 12 months old the Oxaliplatin dose used was 4.3 mg/kg. The most common adverse reactions reported were sensory neuropathy (53%. G3/4: 15%), thrombocytopenia (40%, G3/4: 26%), anemia (40%, G3/4: 15%), vomiting (32%, G3/4: 0%), nausea (30%, G3/4: 2%) and AST increased (26%, G3/4: 4%). No responses were observed.

The pharmacokinetic parameters of ultrafiltrable platinum have been evaluated in 105 pediatric patients during the first cycle. The mean clearance in pediatric patients estimated by the population pharmacokinetic analysis was 4.7 L/h. The inter-patient variability of platinum clearance in pediatric cancer patients was 41%. Mean platinum pharmacokinetic parameters in ultrafiltrate were C_{max} of 0.75 ±0.24 mcg/mL, $AUC_{0.48}$ of 7.52 ±5.07 mcgxh/mL and AUC, of 8.83 ±1.57 mcgxh/mL at 85 mg/m² of Oxaliplatin and C_{max} of 1.10 ±0.43 mcg/mL, $AUC_{0.48}$ of 9.74 ±2.52 mcgxh/mL and AUC_{inf} of 17.3 ±5.34 mcgxh/mL at 130 mg/m² of Oxalipla

No significant effect of age on the clearance of ultrafilterable platinum has been

In a adjuvant therapy colon cancer randomized clinical trial, 723 patients treated with Oxaliplatin and infusional 5-fluorouracil/leucovorin were <65 years and 400 patients A descriptive subgroup analysis demonstrated that the improvement in DFS for the Oxaliplatin combination arm compared to the infusional 5-fluorouracil/leucovorin alone

arm appeared to be maintained across genders. The effect of Oxaliplatin in patients \geq 65

years of age was not conclusive. Insufficient subgroup sizes prevented analysis by race. Patients ≥ 65 years of age receiving the Oxaliplatin combination therapy experienced more grade 3-4 granulocytopenia than patients < 65 years of age (45% versus 39%). In a previously untreated for advanced colorectal cancer randomized clinical trial of Oxaliplatin, 160 patients treated with Oxaliplatin and 5-fluorouracil/leucovorin were < 65 years and 99 patients were ≥65 years. The same efficacy improvements in response

In a previously treated for advanced colorectal cancer randomized clinical trial of Oxaliplatin, 95 patients treated with Oxaliplatin and 5-fluorouracil/leucovorin were <65 years and 55 patients were ≥65 years. The rates of overall adverse reactions, including grade 3 and 4 events, were similar across and within arms in the different age groups in all studies. The incidence of diarrhea, dehydration, hypokalemia, leukopenia, fatigue and syncope were higher in patients ≥65 years old. No adjustment to starting dose was required in patients ≥65 years old.

rate, time to tumor progression, and overall survival were observed in the ≥65 year old

Patients with Renal Impairment The exposure (AUC) of unbound platinum in plasma ultrafiltrate tends to increase in

patients as in the overall study population.

renally impaired patients. Caution and close monitoring should be exercised when Oxaliplatin is administered to patients with renal impairment. The starting Oxaliplatin dose does not need to be reduced in patients with mild (creatinine clearance=50-80 mL/ min) or moderate (creatinine clearance=30-49 mL/min) renal impairment. However, the starting dose of Oxaliplatin should be reduced in patients with severe renal impairment (creatinine clearance < 30 mL/min).

Oxaliplatin Concentrate for Solution for Infusion (Oxitan) should be administered under the supervision of a qualified physician experienced in the use of cancer chemotherapeutic agents. Appropriate management of therapy and complications is possible only when adequate diagnostic and treatment facilities are readily available.

Administer Oxaliplatin in combination with 5-fluorouracil/leucovorin every 2 weeks. For advanced disease, treatment is recommended until disease progression or unacceptable toxicity. For adjuvant use, treatment is recommended for a total of 6

Day 1:	Oxaliplatin 85 mg/m² IV infusion in 250-500 mL 5% Dextrose injection USP and leucovorin 200 mg/m² IV infusion in 5% Dextrose injection USP both given over 120 minutes at the same time in separate bag using a Y-line, followed by 5-FU 400 mg/m² IV bolus given over 2-minutes, followed by 5-FU 600 mg/m² IV infusion in 500 mL 5% Dex trose injection, USP (recommended) as a 22-hour continuous infusion
Day 2:	Leucovorin 200 mg/m² IV infusion over 120 minutes, followed by 5-FU 400 mg/m² IV bolus given over 2-4 minutes, followed by 5-FU 600 mg m² IV infusion in 500 mL 5% Dextrose injection, USP (recommended as a 22-hour continuous infusion.

Day 1	5-FU bolus 400 mg/m² over 2 minutes to 4 minutes		5-FU bolus 400 mg/r over 2 minutes to 4 minutes	
Leucovorin	5-FU infusion	Leucovorin	5-FU infusion	
200 mg/m ²	600 mg/m ²	200 mg/m ²	600 mg/m ²	

Concentrate for Solution for Infusion 85 mg/m² 2 h ← 22 hrs → 0 h 2 h ← 22 hrs — ←2 hrs → \leftarrow 2 hrs \rightarrow

The administration of Oxaliplatin does not require prehydration. Premedication with antiemetics, including 5-HT, blockers with or without dexamethasone, is recommended

Prior to subsequent therapy cycles, patients should be evaluated for clinical toxicities and laboratory tests. Prolongation of infusion time for oxaliplatin from 2 hours to 6 hours decreases the $\mathrm{C}_{\mathrm{max}}$ by an estimated 32% and may mitigate acute toxicities. The infusion times for 5-FU and leucovorin do not need to be changed.

Adjuvant Therapy in Patients with Stage III Colon Cancer

Neuropathy and other toxicities were graded using the NCI CTC scale version 1.

For natients who experience persistent grade 2 neurosensory events that do not resolve, a dose reduction of oxaliplatin to 75 mg/m2 should be considered. For patients with persistent grade 3 neurosensory events, discontinuing therapy should be considered. The infusional 5-FU/LV regimen need not be altered

A dose reduction of oxaliplatin to 75 mg/m² and infusional 5-FU to 300 mg/m² bolus and 500 mg/m² 22-hour infusion is recommended for patients after recovery from grade 3/4 gastrointestinal (despite prophylactic treatment) or grade 4 neutropenia or grade 3/4 thrombocytopenia. The next dose should be delayed until: neutrophils ≥1.5 x 10⁹/L and

Dose Modifications in Therapy in Previously Untreated and Previously Treated Patients with Advanced Colorectal Cancer

Neuropathy was graded using a study-specific neurotoxicity scale. Other toxicities were graded by the NCI CTC, version 2.0.

For patients who experience persistent Grade 2 neurosensory events that do not resolve, a dose reduction of oxaliplatin to 65 mg/m² should be considered. For patients with persistent grade 3 neurosensory events, discontinuing therapy should be considered. The 5-FU/LV regimen need not be altered.

A dose reduction of oxaliplatin to 65 mg/m² and 5-FU by 20% (300 mg/m² bolus and 500 mg/m² 22-hour infusion) is recommended for patients after recovery from grade 3/4 gastrointestinal (despite prophylactic treatment) or grade 4 neutropenia or grade 3/4 thrombocytopenia. The next dose should be delayed until: neutrophils >1.5 x 109/L

Preparation of Infusion solution

Oxaliplatin is incompatible in solution with alkaline medications or media hence it is advised that dilution must never be performed with sodium chloride solution or other chloride containing solutions.

To be diluted and given as slow I.V. infusion. The preparation for infusion can be stored for 24 hours at room temperature.

Handling and Disposal

As with other potentially toxic anticancer agents, care should be exercised in the handling and preparation of infusion solutions prepared from oxaliplatin. The use of gloves is recommended. If a solution of oxaliplatin contacts the skin, wash the skin immediately and thoroughly with soap and water. If oxaliplatin contacts the mucous membranes, flush thoroughly with water.

Procedures for the handling and disposal of anticancer drugs should be considered. Several guidelines on the subject have been published. There is no general agreement that all of the procedures recommended in the guidelines are necessary or appropriate

OVERDOSE

There is no known antidote for Oxaliplatin overdose. In addition to thrombocytopenia. the anticipated complications of an Oxaliplatin overdose include hypersensitivity reaction, myelosuppression, nausea, vomiting, diarrhea and neurotoxicity. Several cases of overdoses have been reported with Oxaliplatin. Adverse reactions observed were Grade 4 thrombocytopenia (<25,000/mm³) without any bleeding, anemia, sensory neuropathy such as paresthesia, dysesthesia, laryngospasm and facial muscle spasms, gastrointestinal disorders such as nausea, vomiting, stomatitis, flatulence, abdomen enlarged and Grade 4 intestinal obstruction, Grade 4 dehydration, dyspnea, wheezing, chest pain, respiratory failure, severe bradycardia and death.

Patients suspected of receiving an overdose should be monitored, and supportive treatment should be administered. The maximum dose of Oxaliplatin that has been administered in a single infusion is 825 mg.

Store at a temperature below 25°C. Protect from light. Do not freeze. 10 mL USP type I tubular clear glass vial with 10mL fill.

20 mL USP type I tubular clear glass vial with 20mL fill.

US Prescribing Information of ELOXATIN, sanofi-aventis U.S. LLC Bridgewater, NJ

INSTRUCTIONS TO THE PATIENT

Medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you. Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament. The doctor and pharmacists are expert in medicine, its benefits and risks. Do not by yourself interrupt the period of treatment prescribed for you. Do not repeat the same escription without consulting your doctor.

KEEP MEDICAMENT OUT OF THE REACH OF CHILDREN

Version number: MGE/00/2013 Date of Release: 09/2013

Size: 500 x 400 mm

NA

Form No.: SOP/PDD/003-03 Rev. 05

Approved By

PDD Production

CONFIDENTIAL

able 20 – Adverse I dvanced Colorecta				ients Prev	iously Untr	eated for
	Oxaliplatin + 5-FU/LV N=259		Irinotecan + 5-FU/ LV N=256		Oxaliplatin + irinotecan N=258	
Hematology Parameter	All Grades (%)	Grade 3/4 (%)	All Grades (%)	Grade 3/4 (%)	All Grades (%)	Grade 3/4 (%)
Anemia	27	3	28	4	25	3
_eukopenia	85	20	84	23	76	24
	0.4			4.4	-4	