

Azonit D[®]

(Isoconazole nitrate, Diflucortolone valerate)

Pharmacological properties

Azonit D[®] is a topical antimycotic drug, the presence of diflucortolone valerate makes it particularly indicated for the treatment of mycoses with symptoms of inflammation.

Indications

- Superficial mycoses of hairless and hairy skin e.g. Dermatophytes, pityriasis versicolor, candidiasis.
- Mycoses with cutaneous symptoms of inflammatory or eczematous nature.

Dosage and administration

Usual adult dose:

Azonit D[®] is applied twice daily on the affected area. For treatment of mycoses between toes, a cotton soaked with **Azonit D[®]** should be placed between them.

Usually, after one week of treatment with **Azonit D[®]**, or when the inflammatory and eczematous skin symptoms disappear, it is recommended to discontinue **Azonit D[®]**, and if necessary continue treatment with antimycotic agent alone, or the corticosteroid only.

Usual children dose:

Azonit D[®] should be used by infants only if absolutely necessary and under medical supervision.

Use in pregnancy

Azonit D[®] should be used by pregnant women only if absolutely necessary and under medical supervision.

Side effects

Azonit D[®] is well tolerated. As with other products containing corticosteroids, when treatment takes longer than 4 weeks or when large body areas are involved, side effects related to corticosteroid absorption starts to appear, such as cutaneous atrophy, hypertrichosis, hypopigmentation, striae, telangiectasia, burning sensation, irritation, folliculitis.

Precautions

- Long term treatment may enhance the growth of microorganism not susceptible to the anti-microbial agent present in this product.
- Care should be exercised when applied on the face, as the product should not come near the eyes.

Drug interactions

There are no reported drug interaction.

Contraindications

Hypersensitivity to any of the ingredients, presence of tubercular, luetic or viral infections in the area to be treated.

Warnings

Long term use of the topical products may cause sensitization phenomena, in case it happened, the treatment should be discontinued and appropriate therapy should be employed.

Presentation

Azonit D[®] cream: Isoconazole nitrate 10mg and diflucortolone valerate 1mg/gm of cream.

(Available in different pack sizes).

(This is a medicament - keep medicaments out of reach of children)



Pharma International

- Medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's proscription, method for use and the instructions of the pharmacist who sold the medicament.
- The doctor and the pharmacist are experts in medicine, its benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.