

IBUPROFEN, PARACETAMOL AND FAMOTIDINE TABLETS  
LABORATE **GENTLE RELIEF**  
TABLETS

**Composition:**

Each uncoated tablet contains:

Ibuprofen	BP	400mg
Paracetamol	BP	325mg
Famotidine	BP	20mg
Excipients		q.s.

**Description**

Ibuprofen belongs to a group of medicines called Non-Steroidal Anti-inflammatory Drugs (NSAIDs) that work by relieving pain, reducing inflammation, swelling and fever. Paracetamol may act predominantly by inhibiting prostaglandin synthesis in the central nervous system (CNS) and, to a lesser extent, through a peripheral action by blocking pain-impulse generation. Famotidine is a histamine H<sub>2</sub>-receptor antagonist that inhibits stomach acid production, and it is commonly used in the treatment of peptic ulcer disease and gastroesophageal reflux disease.

**Indication**

For the treatment of mild to moderate pain including headache, migraine, neuralgia, toothache, sore throat, symptomatic relief of sprains, strains, rheumatic pain, sciatica, lumbago, fibrositis, muscular aches and pains, joint swelling and stiffness, influenza, gout pain, feverishness, feverish colds and short-term treatment of gastro esophageal reflux disease.

**Contraindication**

It is not advocated for those allergic to its ingredients, history of hypersensitivity to aspirin or other NSAIDs, aspirin-induced nasal polyps or bronchospasm, active peptic ulceration and in severe renal dysfunction.

**Adverse Reaction**

Epigastric discomfort, hypersensitivity reactions, hematological disturbances, nervousness, dry skin, rash, insomnia, nausea, vomiting, diarrhoea, constipation, muscle pain, agitation and rash.

**Drug Interactions**

Corticosteroids, Anticoagulants, Heparin, Oral Antidiabetics, Insulin, Digoxin, Cyclosporine, Lithium, Methotrexate, Metoclopramide, Probenecid, Cimetidine, Ranitidine and Nizatidine.

**Warning and Precautions**

Patients with moderate or severe renal insufficiency, pregnancy and nursing mothers. Overdose of Paracetamol may be injurious to liver. Bronchospasm may be precipitated in patients suffering from or with a previous history of bronchial asthma. Ibuprofen should not be given to patients in whom aspirin and other non-steroidal anti-inflammatory drugs induce the symptoms of asthma, rhinitis and urticaria.

**Pregnancy and Lactation**

Not Recommended.

**Dosage**

As directed by the Physician.

**Presentation**

2x10 tablets in a unit carton.

**Storage**

Store in cool, dry and dark place.

**Keep all medicines out of the reach of children.**

MANUFACTURED IN INDIA BY :



**LABORATE**  
PHARMACEUTICAL (INDIA)  
(A Unit of Laborate Pharmaceuticals India Ltd.)  
E-11, IND. AREA, PANIPAT - 132 103