

Information for use. Please read carefully!

Trachisan[®]

SORE THROAT GARGLE

Rapid relief of sore throat

**For the treatment of inflammation
in the oral and pharyngeal cavities**

SUGAR-FREE



ready-to-use, always use undiluted

Composition: 100 ml gargle contains 0.100 g lignocaine hydrochloride 1 H₂O and 0.200 g chlorhexidine gluconate.

Note: The preparation contains 7.9% (V/V) alcohol!

Indications: Inflammatory and infectious diseases of the oral and pharyngeal cavities such as tonsillitis (angina tonsillar), inflammation of the gums, the oral mucosa, the tongue and the pharynx; in common colds and to prevent infections after surgical interventions in the oral and pharyngeal cavities (e.g. tooth extraction, tonsillectomy); bad breath. To prevent the formation of soft dental plaque.

Contra-indications: The preparation should not be used in the case of

- hypersensitivity to the active ingredients contained in Trachisan[®] Sore Throat Gargle,
- non-bleeding desquamation of the oral mucosa (erosive and desquamative changes),
- larger, fresh wounds in the oral and pharyngeal cavities.

Side effects: In general, Trachisan[®] Sore Throat Gargle is well tolerated. However, the preparation may cause gustatory disturbances as well as numbness of the tongue. These side effects disappear automatically, however, once treatment has been discontinued. When used over a longer period of time, yellow to brownish discolouration of the teeth, fillings, dentures or tongue may occur rarely, which can be removed by thorough oral hygiene. However, this also disappears spontaneously when use of the preparation is discontinued.

Dosage guidelines and method of use: Unless otherwise prescribed, rinse in the case of inflammation of the oral cavity, or gargle in inflammation of the pharyngeal cavity with 1 tablespoonful of Trachisan[®] Sore Throat Gargle, 3 – 4 times daily, for 1/2 minute. Do not subsequently rinse with water. The gargle should not be swallowed; if swallowed accidentally, however, it is not harmful.

In order to guarantee a long-lasting effect on the mucosa, the gargle should be used after meals.

Duration of use: Trachisan[®] Sore Throat Gargle should not be used for more than 6 weeks without interruption, unless otherwise prescribed by a doctor or dentist.

Storage instructions and shelf life: Trachisan[®] Sore Throat Gargle should be stored below 25°C. Protect from light! The product should not be used after the expiry date.

How supplied: Bottles of 200 ml gargle.

Keep this and all medicine out of children's reach!



**ENGELHARD
ARZNEIMITTEL**

Karl Engelhard
Fabrik pharm. Präparate GmbH & Co. KG
60316 Frankfurt/Main (F.R. Germany)

Stand 08.96/6421/0/Di