

## Betmiga™ 25 mg prolonged-release tablets

## Betmiga™ 50 mg prolonged-release tablets

Mirabegron

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
  - If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours. If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

### What is in this leaflet:

1. What Betmiga is and what it is used for
2. What you need to know before you use Betmiga
3. How to use Betmiga
4. Possible side effects
5. How to store Betmiga
6. Contents of the pack and other information

### 1. What Betmiga is and what it is used for

Betmiga contains the active substance mirabegron. It is a bladder muscle relaxant (a so called beta 3-adrenoceptor agonist), which reduces the activity of an overactive bladder and treats the related symptoms.

Betmiga is used to treat the symptoms of an overactive bladder in adults such as:

- suddenly needing to empty your bladder (called urgency)
- having to empty your bladder more than usual (called increased urinary frequency)
- not being able to control when to empty your bladder (called urgency incontinence)

### 2. What you need to know before you use Betmiga

#### Do not use Betmiga:

- if you are allergic to mirabegron or any of the other ingredients of this medicine (listed in section 6).

#### Warnings and precautions

Talk to your doctor or pharmacist before using Betmiga.

- if you have kidney or liver problems. Your doctor may need to reduce your dose or may tell you not to use Betmiga, especially if you are taking other medicines such as itraconazole, ketoconazole, ritonavir or clarithromycin. Tell your doctor about the medicines that you take if you have very high uncontrolled blood pressure.

if you have an ECG (heart tracing) abnormality known as QT prolongation or you are taking any medicine known to cause this such as:

- medicines used for abnormal heart rhythm such as quinidine, sotalol, procainamide, ibutilide, flecainide, dofetilide, and amiodarone
- medicines used for allergic rhinitis:
- antipsychotic medicines (medicines for mental illness) such as thioridazine, mesoridazine, haloperidol, and chlorpromazine;
- anti-infectives such as pentamidine, moxifloxacin, erythromycin, and clarithromycin.

### Children and adolescents

Do not give this medicine to children and adolescents under the age of 18 years because the safety and efficacy of Betmiga in this age group has not been established.

### Other medicines and Betmiga

Tell your doctor or pharmacist if you are using, have recently used or might use any other medicines.

Betmiga may affect the way other medicines work, and other medicines may affect how this medicine works.

- Tell your doctor if you use thioridazine (a medicine for mental illness), propafenone or flecainide (medicines for abnormal heart rhythm), Imipramine or desipramine (medicines used for depression). These specific medicines may require dose adjustment by your doctor.
- Tell your doctor if you use digoxin (a medicine for heart failure or abnormal heart rhythm). Blood levels of this medicine are measured by your doctor. If the blood level is out of range, your doctor may adjust the dose of digoxin.

Tell your doctor if you use dabigatran etexilate (a medicine to prevent the formation of blood clots in the veins after knee or hip replacement surgery). This medicine may require dose adjustment by your doctor.

### Pregnancy and breast-feeding

If you are pregnant, think you may be pregnant or are planning to have a baby you should not use Betmiga.

If you are breast feeding, ask your doctor or pharmacist for advice before using this medicine. It is likely that this medicine passes into your breast milk. You and your doctor should decide if you should use Betmiga or breastfeed. You should not do both.

### Driving and using machines

There is no information to suggest that this medicine affects your ability to drive or use machines.

### 3. How to use Betmiga

Always use this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose is one 50 mg tablet by mouth once daily. If you have kidney or liver problems, your doctor may need to reduce your dose to one 25 mg tablet by mouth once daily. You should take this medicine with liquids and swallow the tablet whole. Do not crush or chew the tablet. Betmiga can be taken with or without food.

#### If you take more Betmiga than you should

If you have taken more tablets than you have been told to take, or if someone else accidentally takes your tablets, contact your doctor, pharmacist or hospital for advice immediately.

Symptoms of overdose may include a forceful beating of the heart, an increased pulse rate or an increased blood pressure.

#### If you forget to take Betmiga

If you forget to take your medicine, take the missed dose as soon as you remember. If it is less than 6 hours before your next scheduled dose, skip the dose and continue to take your medicine at the usual time.

Do not take a double dose to make up for a forgotten dose. If you miss several doses, tell your doctor and follow the advice given to you.

#### If you stop using Betmiga

Do not stop treatment with Betmiga early. If you do not see an immediate effect. Your bladder might need some time to adapt. You should continue taking your tablets. Do not stop taking them when your bladder condition improves. Stopping treatment may result in recurrence of symptoms of overactive bladder.

Do not stop taking Betmiga without talking to your doctor first, as your overactive bladder symptoms may come back.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

### 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The most serious side effects may include irregular heart beat (atrial fibrillation). This is an uncommon side effect (may affect up to 1 in 100 people), but if this side effect occurs, immediately stop taking the medicine and seek urgent medical advice.

Other side effects include:

#### Common side effects (may affect up to 1 in 10 people)

- Increased heart rate (tachycardia)
- Infection of the structures that carry urine (urinary tract infections)

#### Uncommon side effects (may affect up to 1 in 100 people)

- Bladder infection (cystitis)
- Feeling your heartbeat (palpitations)
- Vaginal infection
- Indigestion (dyspepsia)
- Infection of the stomach (gastritis)
- Swelling of the joints
- Itching of the vulva or vagina (vulvovaginal pruritus)
- Increased blood pressure
- Increase in liver enzymes (GGT, AST and ALT)
- Itching, rash or hives (urticaria, rash, rash macular, rash papular, pruritus)

#### Rare side effects (may affect up to 1 in 1,000 people)

- Swelling of the eyelid (eyelid oedema)
- Swelling of the lip (lip oedema)
- Small purple spots on the skin (purpura)
- Inflammation of small blood vessels mainly affecting the skin (leukocytoclastic vasculitis).

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

#### 5. How to store Betmiga

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton, blister or bottle after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

After first opening of the bottle, the tablets can be stored for 6 months.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

#### 6. Contents of the pack and other information

##### What Betmiga contains

The active substance is mirabegron. Each tablet contains 25 mg or 50 mg of mirabegron.

The other ingredients are:

Tablet core: Macrogols, hydroxypropylcellulose, butylhydroxytoluene, magnesium stearate

Film-coating: Hypromellose, macrogol, iron oxide yellow (E 172), iron oxide red (E 172) (25 mg tablet only).

##### What Betmiga looks like and contents of the pack

Betmiga 25 mg prolonged release film-coated tablets are oval, brown film-coated tablets, debossed with the company logo and "325" on the same side.

Betmiga 50 mg prolonged release film-coated tablets are oval, yellow film-coated tablets, debossed with the company logo and "355" on the same side.

Betmiga is available in aluminium-aluminium blister in packs containing 10, 20, 30, 60, 90 or 200 tablets and in high density polyethylene (HDPE) bottles with child-resistant closures containing 90 tablets.

Not all pack sizes may be available in your country. The bottle may not be available in your country.

##### Marketing Authorisation Holder and Manufacturer

Astellas Pharma Europe B.V.

Sylviusweg 62

2333 BE Leiden

The Netherlands

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

##### België/Belgique/Belgien

Astellas Pharma B.V. Branch

Tel/Tel.: +32 (0)2 5580710

##### България

Астелас Фарма ЕООД

Тел.: +359 2 862 53 72

##### Česká republika

Astellas Pharma s.r.o.

Tel.: +420 236 080300

##### Danmark

Astellas Pharma a/s

Tlf.: +45 43 430355

##### Deutschland

Astellas Pharma GmbH

Tel.: +49 (0)89 454401

##### Eesti

Astellas Pharma Europe B.V.

Holland

Tel.: +31 (0)71 5455745

##### Ελλάδα

Astellas Pharmaceuticals AEBE

Τηλ: +30 210 8189900

##### España

Astellas Pharma S.A.

Tel.: +34 91 4952700

##### France

Astellas Pharma S.A.S.

Tel.: +33 (0)1 55917500

##### Irland

Astellas Pharma Co. Ltd.

Tel.: +353 (0)1 4671555

##### Ísland

Vistor hf

Sím: +354 535 7000

##### Italia

Astellas Pharma S.p.A.

Tel.: +39 (0)2 921381

##### Κύπρος

Astellas Pharmaceuticals AEBE

Τηλ: +30210 8189900

##### Latvija

Astellas Pharma Europe B.V.

Niederlande

Tel.: +31 (0)71 5455745

##### Lietuva

Astellas Pharma Europe B.V.

Nyderlandai

Tel.: +31 (0)71 5455745

##### Luxembourg/Luxemburg

Astellas Pharma B.V. Branch

Belgique/Belgien

Tel/Tel.: +32 (0)2 5580710

##### Magyarország

Astellas Pharma Kft.

Tel.: +36 1 5778200

##### Malta

E.J. Busuttil Ltd.

Tel.: +356 214 47184

##### Nederland

Astellas Pharma B.V.

Tel.: +31 (0)71 5455745

##### Norge

Astellas Pharma

Tlf.: +47 66 76 46 00

##### Österreich

Astellas Pharma Ges.m.b.H.

Tel.: +43 (0)1 8772668

##### Polşa

Astellas Pharma Sp.z o.o.

Tel.: +48 225451 111

##### Portugal

Astellas Pharma, Lda.

Tel.: +351 21 4401320

##### România

S.C. Astellas Pharma SRL

Tel.: +40 (0)21 361 04 95/96/92

##### Slovenija

Astellas Pharma d.o.o.

Tel.: +386 1401 1400

##### Slovenská republika

Astellas Pharma s.r.o.

Tel.: +421 2 4444 2157

##### Suomi/Finland

Astellas Pharma

Puh/Tel.: +358 (0)9 85606000

##### Sverige

Astellas Pharma AB

Tel.: +46 (0)40-650 15 00

##### United Kingdom

Astellas Pharma Ltd.

Tel.: +44 (0)203 379 8700

This leaflet was last revised in January 2013.

Detailed information on this medicine is available on the European Medicines Agency website <http://www.ema.europa.eu>.