

PACKAGE LEAFLET: INFORMATION FOR THE USER



DICLODENT

DICLOFENAC 0.074%



DICLOFENAC 0.074% DOSED AT 0.74 mg/ml AS OROMUCOSAL SOLUTION

Read all of this leaflet carefully because it contains important information for you.

This medicine is available without prescription. However, you still need to use Diclodent carefully to get the best results from it.

Keep this leaflet. You may need to read it again.

Ask your pharmacist if you need more information or advice.

You must contact a doctor if your symptoms worsen or do not improve.

Composition

1 ml of oromucosal solution contains: 0.74 mg of diclofenac.

Pharmaceutical form and presentation

Oromucosal solution in a 200 ml bottle.

Oromucosal solution in 15 ml sachets. Each package contains 12 sachets.

ATC Classification

A01AD11 – Various, non-steroidal anti-inflammatory drug.

Marketing Authorization Holder

Laboratórios Atral, S.A.

Rua da Estação, n.º 42 – Vala do Carregado

2600-726 Castanheira do Ribatejo – Portugal

Tel.: 263 856 800

Fax.: 263 855 020

email: info@atralcipan.pt

WHAT DICLODENT IS USED FOR

Symptomatic treatment of irritation-inflammatory states of the oropharyngeal cavity, also associated with pain (e.g. gingivitis, stomatitis, pharyngitis), also as a consequence of conservative dental treatment or dental extraction.

DO NOT TAKE DICLODENT

If you are hypersensitive (allergic) to the components of the product or to any other substances that are chemically closely correlated to them, to acetylsalicylic acid or other non-steroidal anti-inflammatory drugs.

POSSIBLE SIDE EFFECTS

Although no side effects associated to Diclodent were reported during clinical trials, some signs of oral mucosa irritation (generally mild) and cough may possibly be observed, particularly in supervised treatments.

The use of topical preparations may lead to the development of sensitization phenomena, especially when used for a long period of time. If this occurs, treatment with Diclodent should be discontinued and appropriate therapy instituted, if necessary.

USING OTHER MEDICINES

No interactions with other medicinal products generally used in concomitant treatment have been reported.

PREGNANCY AND BREAST-FEEDING

During pregnancy and breast-feeding, the product should be used only in those cases of absolute necessity and under direct medical control.

EFFECTS ON THE ABILITY TO DRIVE AND USE MACHINES

None.

HOW TO TAKE DICLODENT

Gargle or swish around the mouth with 15 ml of the solution (1 measure or 1 sachet), or diluted with a little water, 2 to 3 times a day.

OVERDOSE

No cases of overdose with Diclodent have been reported.

ATTENTION

If you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

Do not use Diclodent after the expiry date which is stated on the label.

Keep out of the reach and sight of children.

HOW TO STORE DICLODENT

Do not store above 25°C.

This leaflet was last approved in December 2009.

Ref.: FI-01-MAI-10

34822500

