

blockers and calcium channel blockers).

Pressor amines (e.g. noradrenaline, adrenaline): Possible decreased response to pressor amines but not sufficient to preclude their use.

Non-steroidal anti-inflammatory medicines (NSAIDs), including selective COX-2 inhibitors, acetylsalicylic acid >3 g/day), and non-selective NSAIDs: NSAIDs can attenuate the antihypertensive effect of both angiotensin II antagonists and Hydrochlorothiazide when administered simultaneously. Furthermore, concomitant use of Valsartan/Hydrochlorothiazide and NSAIDs may lead to worsening of renal function and an increase in serum potassium. Therefore, monitoring of renal function at the beginning of the treatment is recommended, as well as adequate hydration of the patient.

Interactions related to Valsartan

- Concomitant use not recommended: Potassium-sparing diuretics, potassium supplements, salt substitutes containing potassium and other substances that may increase potassium levels: If a medicinal product that affects potassium levels is considered necessary in combination with Valsartan, monitoring of potassium plasma levels is advised.

- No interaction: In drug interaction studies with Valsartan, no interactions of clinical significance have been found with Valsartan or any of the following substances: Cimetidine, warfarin, furosemide, digoxin, atenolol, indomethacin, Hydrochlorothiazide, amlodipine, glibenclamide. Digoxin and indomethacin could interact with the Hydrochlorothiazide component of Valsartan/Hydrochlorothiazide.

Interactions related to Hydrochlorothiazide

- Concomitant use requiring caution: Medicinal products associated with potassium loss and hypokalemia (e.g. kaliuretic diuretics, corticosteroids, laxatives, ACTH, amphotericin, carbamazepine, penicillin G, salicylic acid and derivatives): If these medicinal products are to be prescribed with the Valsartan/Hydrochlorothiazide combination, monitoring of potassium plasma levels is advised. These medicinal products may potentiate the effect of Hydrochlorothiazide on serum potassium.

Medicinal products that could induce torsades de pointes: Class Ia antiarrhythmics (e.g. quinidine, hydroquinidine, disopyramide); Class III antiarrhythmics (e.g. amiodarone, sotalol, dofetilide, ibutilide); some antipsychotics (e.g. thioridazine, chlorpromazine, levomepromazine, trifluoperazine, cyamemazine, sulpiride, sultopride, amisulpride, tiapride, pimozide, haloperidol, droperidol); others (e.g. bepridil, cisapride, diphenamil, erythromycin i.v., halofantrin, ketanserin, mizolastin, pentamidine, sparflaxacin, terfenadine, vincamine i.v.). Due to the risk of hypokalemia, Hydrochlorothiazide should be administered with caution when associated with medicinal products that could induce torsades de pointes.

Digitalis glycosides: Thiazide-induced hypokalemia or hypomagnesemia may occur as unwanted effects favoring the onset of digitalis-induced cardiac arrhythmias.

Calcium salts and vitamin D: Administration of thiazide diuretics, including Hydrochlorothiazide, with vitamin D or with calcium salts may potentiate the rise in serum calcium.

Antidiabetic agents (oral agents and insulin): The treatment with a thiazide may influence the glucose tolerance. Dose adjustment of the antidiabetic medicinal product may be necessary. Metformin should be used with caution because of the risk of lactic acidosis induced by possible functional renal failure linked to Hydrochlorothiazide.

Beta blockers and diazoxide: Concomitant use of thiazide diuretics, including Hydrochlorothiazide, with beta blockers may increase the risk of hyperglycemia. Thiazide diuretics, including Hydrochlorothiazide, may enhance the hyperglycemic effect of diazoxide.

Medicinal products used in the treatment of gout (probenecid, sulfapyrazone and allopurinol): Dose adjustment of uricosuric medications may be necessary as Hydrochlorothiazide may raise the level of serum uric acid. Increase of dosage of probenecid or sulfapyrazone may be necessary. Co-administration of thiazide diuretics, including Hydrochlorothiazide, may increase the incidence of hypersensitivity reactions to allopurinol.

Anticholinergic agents (e.g. atropine, biperiden): The bioavailability of thiazide-type diuretics may be increased by anticholinergic agents, apparently due to a decrease in gastrointestinal motility and the stomach emptying rate.

Amantadine: Thiazides, including Hydrochlorothiazide, may increase the risk of adverse effects caused by amantadine.

Cholestyramine and cholestipol resins: Absorption of thiazide diuretics, including Hydrochlorothiazide, is impaired in the presence of anionic exchange resins.

Cytotoxic agents (e.g. cyclophosphamide, methotrexate): Thiazides, including Hydrochlorothiazide, may reduce renal excretion of cytotoxic agents and potentiate their myelosuppressive effects.

Non-depolarizing skeletal muscle relaxants (e.g. tubocurarine): Thiazides, including Hydrochlorothiazide, potentiate the action of curare derivatives.

Ciclosporin: Concomitant treatment with ciclosporin may increase the risk of hyperkalemia and gout-type complications.

Alcohol, anesthetics and sedatives: Potentiation of orthostatic hypotension may occur.

Methyldopa: There have been isolated reports of hemolytic anemia in patients receiving concomitant treatment with methyldopa and Hydrochlorothiazide.

Carbamazepine: Patients receiving Hydrochlorothiazide concomitantly with carbamazepine may develop hyponatremia. Such patients should therefore be advised about the possibility of hyponatremic reactions, and should be monitored accordingly.

Iodine contrast media: In case of diuretic-induced dehydration, there is an increased risk of acute renal failure, especially with high doses of the iodine product. Patients should be rehydrated before the administration.

ADVERSE EFFECTS

Adverse drug reactions are ranked by frequency, the most frequent first, using the following convention: Very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$), not known (cannot be estimated from the available data). Within each frequency grouping, adverse reactions are ranked in order of decreasing seriousness.

Valsartan/Hydrochlorothiazide

- Metabolism and nutrition disorders: Dehydration (uncommon).
- Nervous system disorders: Dizziness (very rare); paresthesia (uncommon); syncope (not known).
- Eye disorders: Blurred vision (uncommon).
- Ear and labyrinth disorders: Tinnitus (uncommon).
- Vascular disorders: Hypotension (uncommon).
- Respiratory, thoracic and mediastinal disorders: Cough (uncommon); non cardiogenic pulmonary edema (not known).
- Gastrointestinal disorders: Diarrhea (very rare).
- Musculoskeletal and connective tissue disorders: Myalgia (uncommon); arthralgia (very rare).
- Renal and urinary disorders: Impaired renal function (not known).
- General disorders and administration site conditions: Fatigue (uncommon).
- Investigations: Increased serum uric acid, increased serum bilirubin and serum creatinine, hypokalemia, hyponatremia, elevation of blood urea nitrogen, neutropenia (not known).

Valsartan

- Blood and lymphatic system disorders: Decrease in hemoglobin, decrease in hematocrit, thrombocytopenia (not known).
- Immune system disorders: Other hypersensitivity/allergic reactions including serum sickness (not known).
- Metabolism and nutrition disorders: Increase of serum potassium (not known).
- Ear and labyrinth disorders: Vertigo (uncommon).
- Vascular disorders: Vasculitis (not known).
- Gastrointestinal disorders: Abdominal pain (uncommon).
- Hepatobiliary disorders: Elevation of liver function values (not known).
- Skin and subcutaneous tissue disorders: Angioedema, rash, pruritus (not known).
- Renal and urinary disorders: Renal failure (not known).

Hydrochlorothiazide

- Non-melanoma skin cancer (Basal cell carcinoma and Squamous cell carcinoma) - (not known)
- Blood and lymphatic system disorders: Thrombocytopenia sometimes with purpura (rare); agranulocytosis, leucopenia, hemolytic anemia, bone marrow depression (very rare).
- Immune system disorders: Hypersensitivity reactions (very rare).
- Psychiatric disorders: Depression, sleep disturbances (rare).
- Nervous system disorders: Headache (rare).
- Cardiac disorders: Cardiac arrhythmias (rare).
- Vascular disorders: Postural hypotension (common).
- Respiratory, thoracic and mediastinal disorders: Respiratory distress including pneumonitis and pulmonary edema (very rare).

- Gastrointestinal disorders: Loss of appetite, mild nausea and vomiting (common); constipation, gastrointestinal discomfort (rare); pancreatitis (very rare).

- Hepatobiliary disorders: Intrahepatic cholestasis or jaundice (rare).

- Skin and subcutaneous tissue disorders: Urticaria and other forms of rash (common); photosensitization (rare); necrotising vasculitis and toxic epidermal necrolysis, cutaneous lupus erythematosus-like reactions, reactivation of cutaneous lupus erythematosus (very rare).

- Reproductive system and breast disorders: Impotence (common).

DOSAGE AND ADMINISTRATION

The recommended dose of Viostan® Plus 80mg/12.5mg, Viostan® Plus 160mg/12.5mg and Viostan® Plus 160mg/25mg is one film coated tablet once daily. Dose titration with the individual components is recommended. In each case, up-titration of individual components to the next dose should be followed in order to reduce the risk of hypotension and other adverse events.

When clinically appropriate, direct change from monotherapy to the fixed combination may be considered in patients whose blood pressure is not adequately controlled on Valsartan or Hydrochlorothiazide monotherapy, provided the recommended dose titration sequence for the individual components is followed.

The clinical response to Viostan® Plus film coated tablets should be evaluated after initiating therapy and if blood pressure remains uncontrolled, the dose may be increased by increasing either one of the components to a maximum dose of Viostan® Plus 320mg/25mg.

The antihypertensive effect is substantially present within 2 weeks.

In most patients, maximal effects are observed within 4 weeks. However, in some patients, 4-8 weeks treatment may be required. This should be taken into account during dose-titration.

Method of administration

Viostan® Plus film coated tablets can be taken with or without food and should be administered with water.

Special populations

- Renal impairment: No dose adjustment is required for patients with mild to moderate renal impairment (creatinine clearance ≥ 30 ml/min). Due to the Hydrochlorothiazide component, Viostan® Plus is contraindicated in patients with severe renal impairment.

- Hepatic impairment: In patients with mild to moderate hepatic impairment without cholestasis the dose of Valsartan should not exceed 80 mg. Viostan® Plus is contraindicated in patients with severe hepatic impairment.

- Elderly: No dose adjustment is required in elderly patients.

- Pediatric patients: Viostan® Plus is not recommended for use in children below the age of 18 years due to a lack of data on safety and efficacy.

OVERDOSAGE

Overdose with Valsartan may result in marked hypotension, which could lead to depressed level of consciousness, circulatory collapse and/or shock. In addition, the following signs and symptoms may occur due to an overdose of the Hydrochlorothiazide component: Nausea, somnolence, hypovolemia, and electrolyte disturbances associated with cardiac arrhythmias and muscle spasms.

The therapeutic measures depend on the time of ingestion and the type and severity of the symptoms, stabilization of the circulatory condition being of prime importance.

If hypotension occurs, the patient should be placed in the supine position and salt and volume supplementation should be given rapidly.

Valsartan cannot be eliminated by means of hemodialysis because of its strong plasma binding behavior whereas clearance of Hydrochlorothiazide will be achieved by dialysis.

STORAGE CONDITIONS

Store below 25°C.

Keep in original pack in intact conditions.

Date of revision: November 2018.

L00000

This is a medication

- A medication is a product which affects your health, and its consumption contrary to instructions is dangerous for you
- Follow strictly the doctor's prescription, the method of use, and the instructions of the pharmacist who sold the medication
- The doctor and the pharmacist are experts in medicine, its benefits and risks
- Do not by yourself interrupt the period of treatment prescribed for you
- Do not repeat the same prescription without consulting your doctor
- Medication: keep out of reach of children

Council of Arab Health Ministers
Union of Arab Pharmacists

Benta S.A.L.
Dbayeh - Lebanon