NovoRapid® FlexPen®

100 U/ml solution for injection in pre-filled pen

(equivalent to 3.5 mg).

1 pre-filled pen contains 3 ml equivalent to 300 U.

*Insulin aspart is produced by recombinant DNA technology in Saccharomyces cerevisiae.

Pharmaceutical form

Clear, colourless, aqueous solution for injection in pre-filled pen. FlexPen®.

Therapeutic indications

Treatment of diabetes mellitus in adults, adolescents and children aged 2 years and above.

NovoRapid® is a rapid-acting insulin analogue. NovoRapid® dosage is individual and determined in accordance with the needs of the patient. It should normally be used in combination with intermediate-acting or long-acting insulin given at least once a day. Blood glucose monitoring and insulin dose adjustment are recommended to achieve optimal glycaemic control.

The individual insulin requirement in adults and children is usually between 0.5 and 1.0 U/kg/day. In a basal-bolus treatment regimen, 50 - 70% of this requirement may be provided by NovoRapid® and the remainder by intermediate-acting or long-acting insulin. Adjustment of dosage may be necessary if patients undertake increased physical activity, change their usual diet or during concomitant illness. NovoRapid® has a faster onset and a shorter duration of action than soluble human insulin.

Due to the faster onset of action, NovoRapid® should generally be given immediately before a meal. When necessary NovoRapid® can be given soon after a meal. Due to the shorter duration, NovoRapid® has a lower risk of causing nocturnal hypoglycaemic episodes.

Special populations

As with all insulin products, in elderly patients and patients with renal or hepatic impairment, glucose monitoring should be intensified and the insulin aspart dosage adjusted on an individual basis.

Paediatric population

NovoRapid® can be used in children in preference to soluble human insulin when a rapid onset of action might be beneficial. For example, in the timing of the injections in relation to meals.

Transfer from other insulin products

When transferring from other insulin products, adjustment of the NovoRapid® dose and the dose of the basal insulin may be necessary.

Method of administration

8-9670-00-011-1

NovoRapid® is administered subcutaneously by injection in the abdominal wall, the thigh, the upper arm, the deltoid region or the gluteal region. Injection sites should always be rotated within the same region in order to reduce the risk of lipodystrophy. As with all insulin products, subcutaneous injection in the abdominal wall ensures a faster absorption than other injection sites.

The duration of action will vary according to the dose. injection site, blood flow, temperature and level of physical activity. However, the faster onset of action compared to soluble human insulin is maintained regardless of injection

NovoRapid® FlexPen® is a pre-filled pen designed to be used with NovoFine® or NovoTwist® disposable needles up to a length of 8 mm.

NovoRapid® FlexPen® is colour-coded and accompanied by a package leaflet with detailed instructions for use to be followed

Qualitative and quantitative composition 1 ml of the solution contains 100 U of insulin aspart*

Continuous Subcutaneous Insulin Infusion (CSII): NovoRapid® may be used for Continuous Subcutaneous Insulin Infusion (CSII) in pump systems suitable for insulin infusion. CSII should be administered in the abdominal wall. Infusion sites should be rotated

When used with an insulin infusion pump, NovoRapid® should not be mixed with any other insulin products.

Patients using CSII should be comprehensively instructed in the use of the pump system and use the correct reservoir and tubing for the pump. The infusion set (tubing and cannula) should be changed in accordance with the instructions in the product information supplied with the infusion set. Patients administering NovoRapid® by CSII must have alternative insulin available in case of pump system failure.

Intravenous use:

If necessary, NovoRapid® can be administered intravenously only by healthcare professionals. For intravenous use, infusion systems with NovoRapid® 100 U/ml at concentrations from 0.05 U/ml to 1.0 U/ml insulin

aspart in the infusion fluids 0.9% sodium chloride. 5% dextrose or 10% dextrose inclusive 40 mmol/l potassium chloride using polypropylene infusion bags, are stable at room temperature for 24 hours.

Although stable over time, a certain amount of insulin will be initially adsorbed to the infusion bag. Monitoring of blood glucose is necessary during insulin infusion.

Contraindications

Hypersensitivity to the active substance or any of the excinients

Special warnings and precautions for use

Before travelling between different time zones, the patient should seek the doctor's advice since this may mean that the patient has to take the insulin and meals at different times.

Hyperglycaemia

Inadequate dosing or discontinuation of treatment, especially in type 1 diabetes, may lead to hyperglycaemia and diabetic ketoacidosis.

Hypoglycaemia

Omission of a meal or unplanned, strenuous physical exercise may lead to hypoglycaemia.

Hypoglycaemia may occur if the insulin dose is too high in relation to the insulin requirement.

Patients whose blood glucose control is greatly improved, e.g by intensified insulin therapy, may experience a change in their usual warning symptoms of hypoglycaemia, and should be advised accordingly. Usual warning symptoms may disappear in patients with longstanding diabetes.

A consequence of the pharmacodynamics of rapid-acting insulin analogues is that if hypoglycaemia occurs, it may occur earlier after an injection when compared to soluble human

Since NovoRapid® should be administered in immediate relation to a meal, the rapid onset of action should be considered in patients with concomitant diseases or medication where a delayed absorption of food might be expected.

Concomitant illness, especially infections and feverish conditions, usually increases the patient's insulin requirement Concomitant diseases of the kidney, liver or affecting the adrenal, pituitary or thyroid gland can require changes in the insulin dose

When patients are transferred between different types of insulin products, the early warning symptoms of hypoglycaemia may become less pronounced than those experienced with their previous insulin.

Transfer from other insulin products

Transferring a patient to another type or brand (e.g. as strength or manufacturer) of insulin should be done under strict medical supervision and may require a change in dosage or number of daily injections from that used with their usual insulin products. If an adjustment is needed, it may occur with the first dose or during the first few weeks or months.

Injection site reactions

As with any insulin therapy, injection site reactions may occur and include pain, redness, hives, inflammation, bruising. swelling and itching. Continuous rotation of the injection site within a given area may help to reduce or prevent these reactions. Reactions usually resolve in a few days to a few weeks. On rare occasions, injection site reactions may require discontinuation of NovoRapid®.

Combination of thiazolidinediones and insulin medicinal products

Cases of congestive heart failure have been reported when thiazolidinediones were used in combination with insulin. especially in patients with risk factors for development of congestive heart failure. This should be kept in mind if treatment with the combination of thiazolidinediones and insulin medicinal products is considered. If the combination is used, patients should be observed for signs and symptoms of congestive heart failure, weight gain and oedema. Thiazolidinediones should be discontinued if any deterioration in cardiac symptoms occurs.

Insulin antibodies

Insulin administration may cause insulin antibodies to form. In rare cases, the presence of such insulin antibodies may necessitate adjustment of the insulin dose in order to correct a tendency to hyper- or hypoglycaemia.

Interaction with other medicinal products and other forms of interaction

A number of medicinal products are known to interact with the glucose metabolism.

The following substances may reduce the patient's insulin requirements:

Oral antidiabetic products, monoamine oxidase inhibitors (MAOIs), beta-blockers, angiotensin converting enzyme (ACE) inhibitors, salicylates, anabolic steroids and sulphonamides.

The following substances may increase the patient's insulin requirements:

Oral contraceptives, thiazides, glucocorticoids, thyroid hormones, sympathomimetics, growth hormone and danazol. Beta-blocking agents may mask the symptoms of

hypoglycaemia. Octreotide/lanreotide may either increase or decrease the insulin requirements

Alcohol may intensify or reduce the hypoglycaemic effect of insulin.

Pregnancy

NovoRapid® (insulin aspart) can be used in pregnancy. Data from two randomised controlled clinical trials do not indicate any adverse effect of insulin aspart on pregnancy or on the health of the foetus/newborn when compared to soluble human insulin (see *Pharmacodynamic properties*).

Intensified blood glucose control and monitoring of pregnant women with diabetes are recommended throughout pregnancy and when contemplating pregnancy. Insulin requirements usually fall in the first trimester and increase subsequently during the second and third trimesters. After delivery, insulin requirements normally return rapidly to pre-pregnancy values.

Lactation

There are no restrictions on treatment with NovoRapid® during breast-feeding. Insulin treatment of the nursing mother presents no risk to the baby. However, the NovoRapid® dosage may need to be adjusted.

Effects on ability to drive and use machines

The patient's ability to concentrate and react may be impaired as a result of hypoglycaemia. This may constitute a risk in situations where these abilities are of special importance (e.g. driving a car or operating machinery). Patients should be advised to take precautions to avoid

hypoglycaemia while driving. This is particularly important in those who have reduced or absent awareness of the warning signs of hypoglycaemia or have frequent episodes of hypoglycaemia.

Undesirable effects

a. Summary of the safety profile

Adverse reactions observed in patients using NovoRapid® are mainly due to the pharmacologic effect of insulin. The most frequently reported adverse reaction during treatment is hypoglycaemia. The frequencies of hypoglycaemia vary with patient population, dose regimens and level of glycaemic control, please see section c below. At the beginning of the insulin treatment, refraction anomalies, oedema and injection site reactions (pain, redness, hives, inflammation, bruising, swelling and itching at the injection site) may occur. These reactions are usually of transitory nature. Fast improvement in blood glucose control may be associated with acute painful neuropathy, which is usually reversible. Intensification of insulin therapy with abrupt improvement in glycaemic control may be associated with temporary worsening of diabetic retinopathy, while long-term improved glycaemic control decreases the risk of progression of diabetic retinopathy.

b. Tabulated list of adverse reactions

Adverse reactions listed below are based on clinical trial data and classified according to MedDRA System Organ Class. Frequency categories are defined according to the following convention: Very common (≥ 1/10); common (≥ 1/100 to < 1/10): uncommon (> 1/1.000 to < 1/100): rare (≥ 1/10,000 to < 1/1,000); very rare (< 1/10,000); not known (cannot be estimated from the available data).

Immune system disorders	Uncommon – Urticaria, rash, eruptions
	Very rare – Anaphylactic reactions*
Metabolism and nutrition disorders	Very common – Hypoglycaemia*
Nervous system disorders	Rare – Peripheral neuropathy (painful neuropathy)
Eye disorders	Uncommon – Refraction disorders
	Uncommon – Diabetic retinopathy
Skin and subcutaneous tissue disorders	Uncommon – Lipodystrophy*
General disorders and administration site conditions	Uncommon – Injection site reactions
	Uncommon – Oedema

* see section c

c. Description of selected adverse reactions

Anaphylactic reactions

The occurrence of generalised hypersensitivity reactions (including generalised skin rash, itching, sweating, gastrointestinal upset, angioneurotic oedema, difficulties in breathing, palpitation and reduction in blood pressure) is very rare but can potentially be life threatening.

Hypoglycaemia

The most frequently reported adverse reaction is hypoglycaemia. It may occur if the insulin dose is too high in relation to the insulin requirement. Severe hypoglycaemia may lead to unconsciousness and/or convulsions and may result in temporary or permanent impairment of brain function or even death. The symptoms of hypoglycaemia usually occur suddenly. They may include cold sweats, cool pale skin, fatique, nervousness or tremor, anxiousness, unusual tiredness or weakness, confusion, difficulty in concentration, drowsiness, excessive hunger, vision changes, headache, nausea and palpitation.

In clinical trials, the frequency of hypoglycaemia varied with patient population, dose regimens and level of glycaemic control. During clinical trials the overall rates of hypoglycaemia did not differ between patients treated with insulin aspart compared to human insulin.

Lipodystrophy

Lipodystrophy is reported as uncommon. Lipodystrophy may occur at the injection site.

Overdose

A specific overdose for insulin cannot be defined, however. hypoglycaemia may develop over seguential stages if too high doses relative to the patient's requirements are administered

- Mild hypoglycaemic episodes can be treated by oral administration of glucose or sugary products. It is therefore recommended that the diabetic patient always carries sugar containing products.
- Severe hypoglycaemic episodes, where the patient has become unconscious, can be treated with glucagon (0.5 to 1 mg) given intramuscularly or subcutaneously by a trained person, or with glucose given intravenously by a healthcare professional. Glucose must be given intravenously if the patient does not respond to glucagon within 10 to 15 minutes.

Upon regaining consciousness, administration of oral carbohydrate is recommended for the patient in order to prevent a relapse.

Pharmacodynamic properties

Pharmacotherapeutic group: Drugs used in diabetes. Insulins and analogues for injection, fast-acting. ATC code A10AB05

Mechanism of action

NovoRapid® produces a more rapid onset of action compared to soluble human insulin, together with a lower glucose concentration, as assessed within the first four hours after a meal, NovoRapid® has a shorter duration of action compared to soluble human insulin after subcutaneous injection. When NovoRapid® is injected subcutaneously, onset of action will occur within 10 to 20 minutes of injection. The maximum effect is exerted between 1 and 3 hours after injection. The duration of action is 3 to 5 hours.

Insulin aspart is equipotent to soluble human insulin on a molar basis.

Adults: clinical trials in patients with type 1 diabetes have demonstrated a lower postprandial blood glucose with NovoRapid® compared to soluble human insulin. In two long-term open label trials in patients with type 1 diabetes comprising 1070 and 884 patients, respectively. NovoRapid® reduced glycosylated haemoglobin by 0.12 percentage points and by 0.15 percentage points compared to soluble human insulin; a difference of doubtful clinical significance.

Elderly: in a PK/PD trial the relative differences in the PD properties between insulin aspart and soluble human insulin in the elderly patients with type 2 diabetes were similar to those seen in healthy subjects and younger patients with diabetes.

Children and adolescents: when given to children.

NovoRapid® showed similar long-term glucose control compared to soluble human insulin. In clinical trials for children and adolescents aged 2 to 17 the pharmacodynamic profile of insulin aspart in children was similar to that seen in adults.

Clinical trials in patients with type 1 diabetes have demonstrated a reduced risk of nocturnal hypoglycaemia with insulin aspart compared to soluble human insulin. The risk of daytime hypoglycaemia was not significantly increased.

Pregnancy: a clinical trial comparing safety and efficacy of insulin aspart vs. soluble human insulin in the treatment of pregnant women with type 1 diabetes (322 exposed pregnancies) did not indicate any adverse effect of insulin aspart on pregnancy or on the health of the foetus/newborn. In addition, the data from a clinical trial including 27 women with destational diabetes randomised to treatment with insulin aspart vs. soluble human insulin showed similar safety profiles between treatments as well as a significant improvement in postprandial glucose control in the insulin aspart treated group.

Pharmacokinetic properties

In NovoRapid® substitution of amino acid proline with aspartic acid at position B28 reduces the tendency to form hexamers as observed with soluble human insulin. NovoRapid® is therefore more rapidly absorbed from the subcutaneous layer compared to soluble human insulin.

The time to maximum concentration is, on average, half of that for soluble human insulin. A mean maximum plasma concentration of 492 pmol/l was reached 40 minutes after

a subcutaneous dose of 0.15 U/kg bodyweight in type 1 diabetic patients. The insulin concentrations returned

to baseline about 4 to 6 hours after dose. The absorption rate was somewhat slower in type 2 diabetic patients, resulting in a lower C_{max} (352 ± 240 pmol/l) and later t_{max} (60 minutes). The intra-individual variability in time to maximum concentration is significantly less for NovoRapid® than for soluble human insulin, whereas the intra-individual variability in C_{max} for NovoRapid[®] is larger.

Children and adolescents: the pharmacokinetic and pharmacodynamic properties of NovoRapid® were investigated in children and adolescents with type 1 diabetes. Insulin aspart was rapidly absorbed in both age groups, with similar t_{max} as in adults. However, C_{max} differed between the age groups, stressing the importance of the individual titration of NovoRapid®.

Elderly: the relative differences in pharmacokinetic properties between insulin aspart and soluble human insulin in elderly patients with type 2 diabetes were similar to those observed in healthy subjects and in younger patients with diabetes. A decreased absorption rate was observed in elderly subjects. resulting in a later t_{max} (82 minutes), whereas C_{max} was similar to that observed in younger patients with type 2 diabetes and slightly lower than in patients with type 1 diabetes.

Hepatic impairment: in subjects with hepatic impairment t_{ma} was delayed to about 85 min. (50 min. in subjects with normal hepatic function) while AUC, C_{max} and CL/F were

Renal impairment: a single dose pharmacokinetic study of insulin aspart in 18 subjects with normal to severely impaired renal function was performed. No apparent effect of creatinine clearance values on AUC, C_{max}, CL/F and t_{max} of insulin aspart was found. Data were limited in subjects with moderate and severe renal impairment. Subjects with renal failure necessitating dialysis treatment were not investigated

Preclinical safety data

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity and toxicity to reproduction. In in vitro tests, including binding to insulin and IGF-1 receptor sites and effects on cell growth, insulin aspart behaved in a manner that closely resembled human insulin. Studies also demonstrate that the dissociation of binding to the insulin receptor of insulin aspart is equivalent to human insulin

List of excipients

Glycerol, phenol, metacresol, zinc chloride, disodium phosphate dihydrate, sodium chloride, hydrochloric acid/sodium hydroxide (for pH adjustment) and water for injections.

Incompatibilities

Substances added to NovoRapid® may cause degradation of insulin aspart, e.g. if the product contains thiol or sulphites. This product must not be mixed with other products. Exceptions are NPH (Neutral Protamine Hagedorn) insulin and infusion fluids as described in section Posology.

Special precautions for storage

Store in a refrigerator ($2^{\circ}C - 8^{\circ}C$). Keep away from the cooling element. Do not freeze. Keep the pen cap on NovoRapid® FlexPen® in order to protect from light.

NovoRapid® must be protected from excessive heat and light. After first opening or carried as a spare: Do not refrigerate. Store below 30°C

The in-use shelf life is 4 weeks.

Nature and contents of container

3 ml solution in cartridge (type 1 glass) with a plunger (bromobutyl) and a stopper (bromobutyl/polyisoprene) contained in a pre-filled multidose disposable pen made of polypropylene in a carton.

Pack sizes of 1, 5 and 10 pre-filled pens. Not all pack sizes may be marketed.

Special precautions for disposal and other handling Needles and NovoRapid® FlexPen® must not be shared. The

cartridge must not be refilled NovoRapid® must not be used if it does not appear clear and

colourless or if it has been frozen. The patient should be advised to discard the needle after each

injection. NovoRapid® may be used in an infusion pump system (CSII) as described in section Method of administration. Tubings in which the inner surface materials are made of polyethylene or

polyolefin have been evaluated and found compatible with pump use. In case of emergency in current NovoRapid® users (hospitalisation or insulin pen malfunction), NovoRapid® can be withdrawn with an U100 insulin syringe from the FlexPen®.

Produced by

Novo Nordisk A/S, Novo Allé, DK-2880 Bagsværd, Denmark

INSTRUCTIONS FOR USE FOR THE PATIENT

Do not use NovoRapid®

- ▶ If you are allergic (hypersensitive) to insulin aspart or any of the other ingredients in NovoRapid®.
- ► If you suspect hypoglycaemia (low blood sugar) is ► If FlexPen® is dropped, damaged or crushed.
- ▶ If it has not been stored correctly or if it has been ► If the insulin does not appear clear and colourless.

► Always use a new needle for each injection to prevent

Before using NovoRapid®

- ► Check the label to make sure it is the right type of
- contamination ► Needles and NovoRapid® FlexPen® must not be

Method of administration

shared

NovoRapid® is for injection under the skin (subcutaneously) or for continuous infusion in a pump system. NovoRapid® may also be given directly into a vein (intravenously) by healthcare professionals. Never inject your

insulin directly into a muscle (intramuscular). Always vary the sites you inject within the same region to reduce the risk of developing lumps or skin pitting. The best places to give yourself an injection are: the front of your waist (abdomen); the upper arm or the front of your thighs. The insulin will work more quickly if injected around the

How to handle NovoRapid® FlexPen®

Read the included NovoRapid® FlexPen® INSTRUCTIONS FOR USE carefully. You must use the pen as described in the Instructions for Use

waist. You should measure your blood sugar regularly.

NovoRapid®, FlexPen®, NovoFine® and NovoTwist® are trademarks owned by Novo Nordisk A/S, Denmark.

© 2014 Novo Nordisk A/S

NovoRapid® solution for injection in pre-filled pen. FlexPen® INSTRUCTIONS FOR USE FOR THE PATIENT

Please read the following instructions carefully before using your NovoRapid®


Your FlexPen® is a unique dial-a-dose insulin pen. You can select doses from 1 to 60 units in increments of 1 unit. FlexPen® is designed to be used with NovoFine® or NovoTwist® disposable needles up to a length of 8 mm. As a precautionary measure, always carry a spare insulin delivery device in case your FlexPen® is lost or damaged.

Your NovoRapid® FlexPen® Needle (example) selector Pen cap Residual scale Push-— Big outer needle cap Cartridge -Pointer | button Inner needle cap —

Preparing your NovoRapid® FlexPen®

Check the label to make sure that your FlexPen® contains the correct type of insulin.

A Pull off the pen cap.



Remove the protective tab from a new disposable needle. Screw the needle straight and tightly onto your FlexPen®.



Pull off the big outer needle cap and keep it for later.



Pull off the inner needle cap and dispose of it.



- ▲ Always use a new needle for each injection to prevent contamination.
- \triangle Be careful not to bend or damage the needle before use.
- △ To reduce the risk of unexpected needle sticks, never put the inner needle cap back on when you have removed it from the needle.

Checking the insulin flow

Prior to each injection small amounts of air may collect in the cartridge during normal use. To avoid injection of air and ensure proper dosing:

Turn the dose selector to select 2 units.



F Hold your FlexPen® with the needle pointing upwards and tap the cartridge gently with your finger a few times to make any air bubbles collect at the top of the cartridge.



G Keeping the needle upwards, press the push-button all the way in. The dose selector returns to 0.

A drop of insulin should appear at the needle tip. If not, change the needle and repeat the procedure no more

If a drop of insulin still does not appear, the pen is defective, and you must use a new one.



Selecting your dose

Check that the dose selector is set at 0.

H Turn the dose selector to select the number of units you need to inject.

The dose can be corrected either up or down by turning the dose selector in either direction until the correct dose lines up with the pointer. When turning the dose selector, be careful not to push the push-button as insulin will

You cannot select a dose larger than the number of units left in the cartridge



Needle

Protective tab

Making the injection

Maintenance

Do not refill your FlexPen®.

Insert the needle into your skin. Use the injection technique shown by your doctor

Your FlexPen® is designed to work accurately and safely. It must be handled with care.

If it is dropped or crushed, there is a risk of damage and leakage of insulin.

You can clean the exterior of your FlexPen® by wiping it with a medicinal swab.

Inject the dose by pressing the push-button all the way in until 0 lines up with the pointer. Be careful only to push the push-button when injecting.

Do not soak it, wash or lubricate it as it may damage the pen.

Turning the dose selector will not inject insulin.




Meep the push-button fully depressed after the injection until the needle has been withdrawn from the skin.

The needle must remain under the skin for at least 6 seconds. This will ensure that the full dose has been iniected.



K Lead the needle into the big outer needle cap without touching the big outer needle cap. When the needle is covered, carefully push the big outer needle cap completely on and then unscrew the needle. Dispose of it carefully and put the pen cap back on.



△ Do not use the residual scale to measure your dose of insulin

- ⚠ Always remove the needle after each injection and store your FlexPen® without the needle attached. Otherwise the liquid may leak out which can cause inaccurate dosing.
- △ Caregivers should be most careful when handling used needles to avoid needle sticks.
- △ Dispose of the used FlexPen® carefully without the needle attached.
- ⚠ Needles and NovoRapid® FlexPen® must not be shared.

