

Fucithalmic®

Viscous eye-drops

Fucithalmic® is a sterile 1% aqueous sustained-release formulation of fusidic acid, preserved with benzalkonium chloride.

Properties:

Fucithalmic® is active against Staphylococci, Streptococci, Neisseria, Haemophilus, Moraxella and Corynebacteria.

The sustained-release formulation of Fucithalmic® ensures a prolonged contact with the conjunctival sac. A twice-daily application provides sufficient concentrations in all relevant tissues of the eye. Fucithalmic® penetrates well into the aqueous humour.

Indications:

Bacterial eye infections caused by susceptible organisms in conjunctivitis, blepharitis, sty, keratitis, dacryocystitis and in connection with removal of foreign bodies.

Dosage:

One drop to be instilled into the eye twice daily. Treatment should be continued for at least two days after the eye appears normal.

Contra-indications:

Hypersensitivity to any component.

Precautions:

Contact lenses should not be worn when Fucithalmic® viscous eye-drops are used.

Side-effects:

Transient stinging may occur. Fucithalmic® has a very low allergenic potential.

Discard 30 days after opening.

Do not store above 30°C.

Date of last revision: June 2009.

THIS IS A MEDICAMENT

- Medicament is a product, but not like other products.

- Medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.

- Follow strictly the Doctor's prescription, the method of use and the instructions of the Pharmacist who sold you the medicament. The Doctor and Pharmacist are experts in medicaments, their benefits and their risks.

- Do not by yourself interrupt the period of treatment prescribed to you.

- Do not repeat the same medicament and do not increase doses without consulting your Doctor.

Do not leave medicament within reach of children.

Fucithalmic®

Gotas oftálmicas viscosas

Fucithalmic® es una formulación acuosa estéril al 1% de cesión sostenida del ácido fusídico, preservada con cloruro de benzalconio.

Propiedades:

Fucithalmic® es activo frente a los estafilococos, estreptococos, Neisseria, Haemophilus, Moraxella y Corynebacteria. La formulación de cesión sostenida de Fucithalmic® asegura un contacto prolongado con el saco conjuntival. Una aplicación dos veces al día brinda concentraciones suficientes de ácido fusídico en todos los tejidos involucrados del ojo. Fucithalmic® tiene gran poder para penetrar en el humor acuoso.

Indicaciones:

Infecciones bacterianas del ojo causadas por microorganismos sensibles en la conjuntivitis, blefaritis, orzuelo, queratitis, dacriocistitis y al retirar cuerpos extraños.

Posología:

Se aplica una gota al ojo dos veces al día. El tratamiento debe continuarse dos días, como mínimo, después de recuperar el ojo su aspecto normal.