

antibiophilus®

CAPSULES

COMPOSITION

Lyophilized culture of *Lactobacillus casei* variety *rhamnosus* containing at least 8×10^8 cells per gram 250 mg
for one capsule No. 2 of 250 mg.

Composition of capsule tunic: gelatin, sodium bisulfite, titanium dioxide (E 171).

PRESENTATION

Bottle (glass) containing 20 capsules.

THERAPEUTIC INDICATIONS

This drug is indicated for the symptomatic treatment of certain diarrhoeas. The method of rehydration will be determined by your physician.

PRECAUTIONS FOR USE

IN ORDER TO AVOID POSSIBLE DRUG INTERACTIONS, YOU MUST SYSTEMATICALLY INFORM YOUR PHYSICIAN OR YOUR PHARMACIST ABOUT ANY OTHER DRUGS YOU ARE TAKING.
KEEP OUT OF THE REACH OF CHILDREN.

In adults:

Add sweetened drinks and salted bouillon to this treatment to compensate for liquid loss due to diarrhoea.

If the diarrhoea lasts longer than 2 days, CONSULT YOUR PHYSICIAN.

In the case of severe diarrhoea, or if blood or mucus is present in the stools, as well as in the case of fever, vomiting or other associated gastrointestinal symptoms, CONSULT YOUR PHYSICIAN.

In children:

Infants (under 2 years): consult your physician. In the meantime, stop milk and milk-product diet: administer sweetened drinks (fruit juice) and salted drinks (bouillon) in quantities sufficient to compensate for liquid loss caused by diarrhoea.

Check bodyweight at 6-hour intervals.

The infant should remain lively and energetic; if this is not the case, it would mean that the infant is dehydrated and should be transferred immediately to hospital.

Children over 2 years: this treatment does not exempt from rehydration: sweetened drinks, salted bouillon, discontinuation of milk and milk-product diet on the first day.

If the diarrhoea lasts longer than 2 days, CONSULT YOUR PHYSICIAN.

In the case of severe diarrhoea, or if blood or mucus is present in the stools, as well as in the case of fever, vomiting or other associated gastrointestinal symptoms, CONSULT YOUR PHYSICIAN.

OTHER POSSIBLE EFFECTS OF THE DRUG

LIKE ANY ACTIVE DRUG, THIS MEDICATION MAY HAVE SOME UNDESIRABLE EFFECTS IN CERTAIN INDIVIDUALS. DO NOT HESITATE TO SEEK ADVICE FROM YOUR PHYSICIAN OR PHARMACIST.

DOSAGE AND METHOD OF ADMINISTRATION

2 to 8 capsules per day.

Swallow the capsule with liquid, preferably not during meals.

IN ALL CASES, FOLLOW YOUR PHYSICIAN'S PRESCRIPTION TO THE LETTER.

DELIVERY CONDITIONS

THIS DRUG WAS PRESCRIBED TO YOU IN A SET OF SPECIFIC CIRCUMSTANCES:

- IT SHOULD NOT BE USED IN ANOTHER SET OF CIRCUMSTANCES.
- TREATMENT WITH IT SHOULD NOT BE RESUMED WITHOUT MEDICAL ADVICE
- DO NOT ADVISE SOMEBODY ELSE TO TAKE IT.

SHELF-LIFE

DO NOT EXCEED THE EXPIRY DATE INDICATED ON THE PACKAGE.

Laboratoires LYOCENTRE

24, avenue Georges Pompidou

15000 AURILLAC - FRANCE

Tel. (33) (0)4.71.46.80.00