

Synjardy 5 mg/850 mg film-coated tablets
Synjardy 5 mg/1,000 mg film-coated tablets
Synjardy 12.5 mg/850 mg film-coated tablets
Synjardy 12.5 mg/1,000 mg film-coated tablets
Empagliflozin/metformin hydrochloride

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Synjardy is and what it is used for
2. What you need to know before you take Synjardy
3. How to take Synjardy
4. Possible side effects
5. How to store Synjardy
6. Contents of the pack and other information

1. What Synjardy is and what it is used for

Synjardy is a diabetes medicine that contains two active substances called empagliflozin and metformin.

- Empagliflozin works by blocking a protein in the kidneys called sodium glucose co-transporter 2 (SGLT2). SGLT2 prevents glucose (blood sugar) from being lost in urine by absorbing glucose back into the bloodstream as blood is being filtered in the kidneys. By blocking this protein, the medicine causes blood sugar, sodium (salt) and water to be removed via the urine. Blood sugar levels, which are too high because of your type 2 diabetes, are thereby reduced. This medicine can also help prevent heart disease.
- Metformin works in a different way to lower blood sugar levels, mainly by blocking glucose production in the liver.

Synjardy is added to diet and exercise to treat type 2 diabetes in adult patients (aged 18 years and older) whose diabetes cannot be controlled by adding metformin alone or metformin with other medicines for diabetes.

Synjardy can also be combined with other medicines for the treatment of diabetes. These may be medicines taken by mouth or given by injection such as insulin.

In addition, Synjardy can be used as an alternative to taking both empagliflozin and metformin as single tablets. To avoid overdose, do not continue taking empagliflozin and metformin tablets separately, if you are taking this medicine.

It is important that you continue with your diet and exercise plan as told by your doctor, pharmacist or nurse.

What is type 2 diabetes?

Type 2 diabetes is a disease that comes from both your genes and your lifestyle. If you have type 2 diabetes, your pancreas does not make enough insulin to control the level of glucose in your blood, and your body is unable to use its own insulin effectively. This results in high levels of glucose in your blood which can lead to medical problems like heart disease, kidney disease, blindness, and poor circulation in your limbs.

2. What you need to know before you take Synjardy

Do not take Synjardy:

- if you are allergic to empagliflozin, metformin or any of the other ingredients of this medicine (listed in section 6);
- if you have severely reduced kidney function;
- if you have uncontrolled diabetes, with, for example, severe hyperglycaemia (high blood glucose), nausea, vomiting, diarrhoea, rapid weight loss, lactic acidosis (see 'Risk of lactic acidosis' below) or ketoacidosis. Ketoacidosis is a condition in which substances called 'ketone bodies' accumulate in the blood and which can lead to diabetic pre-coma. Symptoms include stomach pain, fast and deep breathing, sleepiness or your breath developing an unusual fruity smell;
- if you have had a diabetic pre-coma;
- if you have a severe infection such as an infection affecting your lung or bronchial system or your kidney. Severe infections may lead to kidney problems, which can put you at risk for lactic acidosis (see 'Warnings and precautions');
- if you have lost a lot of water from your body (dehydration), e.g. due to long-lasting or severe diarrhoea, or if you have vomited several times in a row. Dehydration may lead to kidney problems, which can put you at risk for lactic acidosis (see 'Warnings and precautions');
- if you are treated for acute heart failure or have recently had a heart attack, have severe problems with your circulation (such as shock) or have breathing difficulties. This may lead to a lack in oxygen supply to tissue which can put you at risk for lactic acidosis (see section 'Warnings and precautions');
- if you have problems with your liver;
- if you drink alcohol to excess, either every day or only from time to time (see section "Synjardy with alcohol").

Warnings and precautions

Talk to your doctor, pharmacist or nurse before taking this medicine, and during treatment:

- about what you can do to prevent dehydration;
- if you have "type 1 diabetes" – this type usually starts when you are young and your body does not produce any insulin;
- if you experience rapid weight loss, feeling sick or being sick, stomach pain, excessive thirst, fast and deep breathing, confusion, unusual sleepiness or tiredness, a sweet smell to your breath, a sweet or metallic taste in your mouth, or a different odour to your urine or sweat,

contact a doctor or the nearest hospital straight away. These symptoms could be a sign of “diabetic ketoacidosis” – a rare, but serious, sometimes life-threatening problem you can get with diabetes because of increased levels of “ketone bodies” in your urine or blood, seen in tests. The risk of developing diabetic ketoacidosis may be increased with prolonged fasting, excessive alcohol consumption, dehydration, sudden reductions in insulin dose, or a higher need of insulin due to major surgery or serious illness;

- if you are 75 years old or older, as increased passing of urine due to the medicine may affect fluid balance in your body and increase your risk of dehydration. Possible signs are listed in section 4, ‘Possible side effects’, dehydration;
- if you are 85 years old or older as you should not start taking Synjardy;
- if you have a serious infection of the kidney or the urinary tract with fever. Your doctor may ask you to stop taking Synjardy until you have recovered.

Risk of lactic acidosis

Synjardy may cause a very rare, but very serious side effect called lactic acidosis, particularly if your kidneys are not working properly. The risk of developing lactic acidosis is also increased with uncontrolled diabetes, serious infections, prolonged fasting or alcohol intake, dehydration (see further information below), liver problems and any medical conditions in which a part of the body has a reduced supply of oxygen (such as acute severe heart diseases).

If any of the above apply to you, talk to your doctor for further instructions.

Stop taking Synjardy for a short time if you have a condition that may be associated with dehydration (significant loss of body fluids) such as severe vomiting, diarrhoea, fever, exposure to heat or if you drink less fluid than normal. Talk to your doctor for further instructions.

Stop taking Synjardy and contact a doctor or the nearest hospital immediately if you experience some of the symptoms of lactic acidosis, as this condition may lead to coma.

Symptoms of lactic acidosis include:

- vomiting
- stomach ache (abdominal pain)
- muscle cramps
- a general feeling of not being well with severe tiredness
- difficulty in breathing
- reduced body temperature and heartbeat

Lactic acidosis is a medical emergency and must be treated in a hospital.

If you need to have major surgery you must stop taking Synjardy during and for some time after the procedure. Your doctor will decide when you must stop and when to restart your treatment with Synjardy.

During treatment with Synjardy, your doctor will check your kidney function at least once a year or more frequently if you are elderly and/or if you have worsening kidney function.

Talk to your doctor immediately if you develop a combination of symptoms of pain, tenderness, redness, or swelling of the genitals or the area between the genitals and the anus with fever or feeling generally unwell. These symptoms could be a sign of a rare but serious or even life-threatening infection, called necrotising fasciitis of the perineum or Fournier’s gangrene which destroys the tissue under the skin. Fournier’s gangrene has to be treated immediately

Foot care

Like for all diabetic patients it is important to check your feet regularly and adhere to any other advice regarding foot care given by your health care professional.

Urine glucose

Because of how this medicine works, your urine will test positive for sugar while you are taking this medicine.

Children and adolescents

This medicine is not recommended for use in children and adolescents under 18 years, because it has not been studied in these patients.

Other medicines and Synjardy

If you need to have an injection of a contrast medium that contains iodine into your bloodstream, for example in the context of an X-ray or scan, you must stop taking Synjardy before or at the time of the injection. Your doctor will decide when you must stop and when to restart your treatment with Synjardy.

Tell your doctor if you are taking, have recently taken or might take any other medicines. You may need more frequent blood glucose and kidney function tests, or your doctor may need to adjust the dosage of Synjardy. It is especially important to mention the following:

- medicines which increase urine production (diuretic), as Synjardy may increase the risk of losing too much fluid. Your doctor may ask you to stop taking Synjardy. Possible signs of losing too much fluid from your body are listed in section 4 ‘Possible side effects’.
- other medicines that lower the amount of sugar in your blood such as insulin or a “sulphonylurea” medicine. Your doctor may want to lower the dose of these other medicines, to prevent your blood sugar levels from getting too low (hypoglycaemia).
- medicines that may change the amount of metformin in your blood, especially if you have reduced kidney function (such as verapamil, rifampicin, cimetidine, dolutegravir, ranolazine, trimethoprim, vandetanib, isavuconazole, crizotinib, olaparib).
- bronchodilators (beta-2 agonists) which are used to treat asthma.
- corticosteroids (given by mouth, as an injection, or inhaled), which are used to treat inflammation in diseases like asthma and arthritis.
- medicines used to treat pain and inflammation (NSAID and COX-2-inhibitors, such as ibuprofen and celecoxib).
- certain medicines for the treatment of high blood pressure (ACE inhibitors and angiotensin II receptor antagonists).
- medicines that contain alcohol (see section ‘Synjardy with alcohol’).
- iodinated contrast agents (medicines used during an X-ray, see section ‘Warnings and precautions’).

Synjardy with alcohol

Avoid excessive alcohol intake while taking Synjardy since this may increase the risk of lactic acidosis (see section ‘Warnings and precautions’).

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Do not take Synjardy if you are pregnant. It is unknown if this medicine is harmful to the unborn child.

Metformin passes into human milk in small amounts. It is not known whether empagliflozin passes into human breast milk. Do not take Synjardy if you are breast-feeding.

Driving and using machines

Synjardy has minor influence on the ability to drive and use machines.

Taking this medicine in combination with medicines called sulphonylureas or with insulin can cause blood sugar levels to drop too low (hypoglycaemia), which may cause symptoms such as shaking, sweating and change in vision, and may affect your ability to drive and use machines. Do not drive or use any tools or machines if you feel dizzy while taking Synjardy.

3. How to take Synjardy

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

How much to take

The dose of Synjardy varies depending on your condition and the doses of diabetes medicines you currently take. Your doctor will adjust your dose as necessary and tell you exactly which strength of the medicine to take.

The recommended dose is one tablet twice a day. Your doctor will normally start Synjardy treatment by prescribing the strength of tablet that supplies the same dose of metformin you are already taking (850 mg or 1,000 mg twice a day), and the lowest dose of empagliflozin (5 mg twice a day). If you are already taking both medicines separately, your doctor will start treatment with tablets of Synjardy that will supply the same amount of both. If you have reduced kidney function, your doctor may prescribe a lower dose.

Taking this medicine

- Swallow the tablet whole with water.
- Take the tablets with meals to lower your chance of an upset stomach.
- Take the tablet twice daily by mouth.

Your doctor may prescribe Synjardy together with another diabetes medicine. Remember to take all medicines as directed by your doctor to achieve the best results for your health. Your doctor may need to adjust your doses to control your blood sugar.

Diet and exercise can help your body use its blood sugar better. It is important to stay on the diet and exercise program recommended by your doctor while taking Synjardy.

If you take more Synjardy than you should

If you take more Synjardy tablets than you should have, you may experience lactic acidosis. Symptoms of lactic acidosis are non-specific such as feeling or being very sick, vomiting, stomach ache with muscle cramps, a general feeling of not being well with severe tiredness, and difficulty in breathing. Further symptoms are reduced body temperature and heartbeat. **If this happens to you, you may need immediate hospital treatment, as lactic acidosis can lead to**

coma. Stop taking this medicine immediately and contact a doctor or the nearest hospital straight away (see section 2). Take the medicine pack with you.

If you forget to take Synjardy

If you miss a dose, take it as soon as you remember. If you do not remember until it is time for your next dose, skip the missed dose and go back to your regular schedule. Do not take a double dose of this medicine.

If you stop taking Synjardy

Do not stop taking Synjardy without first consulting your doctor. Your blood sugar levels may increase when you stop taking Synjardy.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Contact a doctor or the nearest hospital straight away if you have any of the following side effects:

Lactic acidosis, seen very rarely (may affect up to 1 in 10,000 people)

Synjardy may cause a very rare but very serious side effect called lactic acidosis (see section ‘Warnings and precautions’). If this happens you must **stop taking Synjardy and contact a doctor or the nearest hospital immediately**, as lactic acidosis may lead to coma.

Diabetic ketoacidosis, seen rarely (may affect up to 1 in 1,000 people)

These are the signs of diabetic ketoacidosis (see section 2, ‘Warnings and precautions’):

- increased levels of “ketone bodies” in your urine or blood
- rapid weight loss
- feeling sick or being sick
- stomach pain
- excessive thirst
- fast and deep breathing
- confusion
- unusual sleepiness or tiredness
- a sweet smell to your breath, a sweet or metallic taste in your mouth or a different odour to your urine or sweat.

This may occur regardless of blood glucose level. Your doctor may decide to temporarily or permanently stop your treatment with Synjardy.

Contact your doctor as soon as possible if you notice the following side effects:

Low blood sugar (hypoglycaemia), seen very commonly (may affect more than 1 in 10 people)

If you take Synjardy with another medicine that can cause low blood sugar, such as a sulfonylurea or insulin, your risk of getting low blood sugar is increased. The signs of low blood sugar may include:

- shaking, sweating, feeling very anxious or confused, fast heart beat
- excessive hunger, headache

Your doctor will tell you how to treat low blood sugar levels and what to do if you get any of the signs above. If you have symptoms of low blood sugar, eat glucose tablets, a high sugar snack or drink fruit juice. Measure your blood sugar if possible and rest.

Urinary tract infection, seen commonly (may affect up to 1 in 10 people)

The signs of urinary tract infection are:

- burning sensation when passing urine
- urine that appears cloudy
- pain in the pelvis, or mid-back pain (when kidneys are infected)

An urge to pass urine or more frequent urination may be due to the way Synjardy works, but they can also be signs of urinary tract infection. If you note an increase in such symptoms, you should also contact your doctor.

Dehydration, seen uncommonly (may affect up to 1 in 100 people)

The signs of dehydration are not specific, but may include:

- unusual thirst
- lightheadedness or dizziness upon standing
- fainting or loss of consciousness

Other side effects while taking Synjardy:

Very common

- feeling sick (nausea), vomiting
- diarrhoea or stomach ache
- loss of appetite

Common

- genital yeast infection (thrush)
- passing more urine than usual or needing to pass urine more often
- itching
- rash or red skin– this may be itchy and include raised bumps, oozing fluid or blisters
- changes to the way things taste
- thirst
- blood tests may show changes in blood fat (cholesterol) levels in your blood

Uncommon

- hives
- straining or pain when emptying the bladder
- blood tests may show changes related to kidney function (creatinine or urea)

- blood tests may show increases in the amount of red blood cells in your blood (haematocrit)

Very rare

- decreased vitamin B12 levels in the blood
- abnormalities in liver function tests, inflammation of the liver (hepatitis)
- redness of the skin (erythema)

Not known

- severe allergic reaction (may include swelling of the face, lips, mouth, tongue, or throat that may lead to difficulty breathing or swallowing)
- necrotising fasciitis of the perineum or Fournier's gangrene, a serious soft tissue infection of the genitals or the area between the genitals and the anus

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet.

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Synjardy

Store below 30°C

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the blister and the carton after 'EXP'. The expiry date refers to the last day of that month.

Do not use this medicine if you notice that the packaging is damaged or shows signs of tampering.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Synjardy contains

The active substances are empagliflozin and metformin.

Each Synjardy 5 mg/850 mg film-coated tablet (tablet) contains 5 mg empagliflozin and 850 mg metformin hydrochloride.

Each Synjardy 5 mg/1,000 mg film-coated tablet (tablet) contains 5 mg empagliflozin and 1,000 mg metformin hydrochloride.

Each Synjardy 12.5 mg/850 mg film-coated tablet (tablet) contains 12.5 mg empagliflozin and 850 mg metformin hydrochloride.

Each Synjardy 12.5 mg/1,000 mg film-coated tablet (tablet) contains 12.5 mg empagliflozin and 1,000 mg metformin hydrochloride.

The other ingredient(s) are:

- **Tablet core:** maize starch, copovidone, colloidal anhydrous silica, magnesium stearate.
- **Film coating:** hypromellose, macrogol 400, titanium dioxide (E171), talc.
Synjardy 5 mg/850 mg and Synjardy 5 mg/1,000 mg tablets also contain iron oxide yellow (E172). Synjardy 12.5 mg/850 mg and Synjardy 12.5 mg/1,000 mg tablets also contain iron oxide black (E172) and iron oxide red (E172).

What Synjardy looks like and contents of the pack

Synjardy 5 mg/850 mg film-coated tablets are yellowish white, oval, biconvex. They have “S5” and the Boehringer Ingelheim logo on one side and “850” on the other.

Synjardy 5 mg/1,000 mg film-coated tablets are brownish yellow, oval, biconvex. They have “S5” and the Boehringer Ingelheim logo on one side and “1000” on the other.

Synjardy 12.5 mg/850 mg film-coated tablets are pinkish white, oval, biconvex. They have “S12” and the Boehringer Ingelheim logo on one side and “850” on the other.

Synjardy 12.5 mg/1,000 mg film-coated tablets are dark brownish purple, oval, biconvex. They have “S12” and the Boehringer Ingelheim logo on one side and “1000” on the other.

The tablets are available in PVC/ PCTFE /aluminum unit dose blisters. The pack sizes are 10 (10's blister x 1), 30 (10's blister x 3), 60 (10's blister x 6) , and 100 film-coated tablets and multipacks containing 120 (2 packs of 60), 180 (2 packs of 90) and 200 (2 packs of 100) film-coated tablets.

Not all pack sizes may be marketed in your country.

Marketing Authorisation Holder

Boehringer Ingelheim International GmbH
Binger Strasse 173
55216 Ingelheim am Rhein
Germany

Manufacturer

Boehringer Ingelheim Pharma GmbH & Co. KG
Binger Strasse 173
55216 Ingelheim am Rhein
Germany

This leaflet was last revised in February 2019 (EU SPC)

For reporting side effects please contact Ministry of Health in your respective country:

Egypt

Egyptian Pharmacovigilance center
Email: PV.Center@Eda.mohealth.gov.eg
Telephone: (+2) 02 23684288 / 23648769 /23640368, ext. 1303
Fax: (+2) 02 23684194

UAE

Pharmacovigilance & Medical Device section
P.O.Box: 1853
Tel: 80011111
Email : pv@moh.gov.ae
Drug Department
Ministry of Health & Prevention
Dubai

Oman

Department of Pharmacovigilance & Drug Information
Directorate General of Pharmaceutical Affairs & Drug Control, MOH-Oman
Email: dg-padc@moh.gov.om
Fax: +968 22358489
Tel No: +968 22357686 / 22357687

Jordan

Jordan Food and Drug Administration (JFDA)
Tel: +962-6-5632000 Ext. 2209 / +962-799038998
Fax: +962-6-5105893
Email: jpc@jfda.jo

Lebanon

Lebanese Ministry of Public Health
Tel: +9611830300
Email: info@moph.gov.lb

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

Boehringer Ingelheim
Middle East & North Africa (Scientific Office)
Dubai, UAE.
Tel: +971 (4) 423 0400
Fax: +971 (4) 423 3637

This is a Medicament

- Medicament is a product which affects your health and its consumption contrary to an instruction is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the Pharmacist who sold the medicament.
- The doctor and the pharmacist are the experts in medicines, their benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.
- Keep all medicaments out of reach of children.