1. NAME OF THE MEDICINAL PRODUCT

Tykerb® 250 mg film-coated tablets

2. QUALITATIVE AND QUANTITATIVE COMPOSITION Each film-coated tablet contains lapatinib ditosylate monohydrate, equivalent to 250 mg lapatinib. For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Film-coated tablet (tablet).

Oval, biconvex, yellow film-coated tablets, with "GS XJG" debossed on one side.

4. CLINICAL PARTICULARS 4.1 Therapeutic indications

kerb is indicated for the treatment of adult patients with breast cancer, whose tumours overexpress HER2 Tykerb is indicated for the treatment of adult patients with breast cancer, whose tunions overexpressing (ErbB2);
- combination with capecitabine for patients with advanced or metastatic disease with progression following

prior therapy, which must have included anthracyclines and taxanes and therapy with trastuzumab in the metastatic setting (see section 5.1). metastatic setting (see Section 5.1).
in combination with trastuzumab for patients with hormone receptor-negative metastatic disease that has progressed on prior trastuzumab therapy(ies) in combination with chemotherapy (see section 5.1).
in combination with an aromatase inhibitor for postmenopausal women with hormone receptor positive metastatic disease, not currently intended for chemotherapy.

4.2 Posology and method of administration Tykerb treatment should only be initiated by a physician experienced in the administration of anti-cancer medicinal products.

HER2 (ErbB2) overexpressing tumours are defined by IHC3+, or IHC2+ with gene amplification or gene amplification alone. HER2 status should be determined using accurate and validated methods.

Tykerb / Capecitabine combination postology
The recommended dose of Tykerb is 1250 mg (i.e. five tablets) once daily continuously.
The recommended dose of capecitabine is 2000 mg/m²/day taken in 2 doses 12 hours apart on days 1-14 in a 21 day cycle (see section 5.1). Capecitabine should be taken with food or within 30 minutes after food.
Please refer to the full prescribing information of capecitabine.

Tykerb / trastuzumab combination posology
The recommended dose of Tykerb is 1000 mg (i.e. four tablets) once daily continuously.
The recommended dose of trastuzumab is 4 mg/kg administered as an intravenous (IV) loading dose, followed by 2 mg/kg IV weekly (see section 5.1). Please refer to the full prescribing information of trastuzumab.

 $\label{thm:combination} \textbf{Tykerb / aromatase inhibitor combination posology} \\ \textbf{The recommended dose of Tykerb is } 1500 \ mg \ (i.e. \ six \ tablets) \ once daily continuously. \\ \textbf{Please refer to the full prescribing information of the co-administered aromatase inhibitor for dosing details.} \\$

Dose delay and dose reduction

Tykerb should be discontinued in patients with symptoms associated with decreased left ventricular eje Tyker snould be discontinued in patients with symptoms associated with decreased left ventricular ejection fraction (LVEF) that are National Cancer Institute Common Terminology Criteria for Adverse Events (NCI CTCAE) grade 3 or greater or if their LVEF drops below the institutions lower limit of normal (see section 4.4). Tykerb may be restarted at a reduced dose (750 mg/day when administered with trastuzumab, 1000 mg/day when administered with an aromatase inhibitor) after a minimum of Queela and if the LVEF. minimum of 2 weeks and if the LVEF recovers to normal and the patient is asymptomatic

Interstitial lung disease / pneumonitis Tykerb should be discontinued in patients who experience pulmonary symptoms which are NCI CTCAE grade 3 or greater (see section 4.4).

Tykerb dosing should be interrupted in patients with diarrhoea which is NCI CTCAE grade 3 or grade 1 or 2 with complicating features (moderate to severe abdominal cramping, nausea or voniting greater than or equal to NCI CTCAE grade 2, decreased performance status, fever, sepsis, neutropenia, frank bleeding or dehydration) (see sections 4.4 and 4.8). Tykerb may be reintroduced at a lower dose (reduced from 1000 mg/ day to 750 mg/day, from 1250 mg/day to 1000 mg/day or from 1500 mg/day to 1250 mg/day) when diarrhoea resolves to grade 1 or less. Tykerb dosing should be permanently discontinued in patients with diarrhoea which is NCI CTCAE grade 4.

Other toxicities Uner toxicities Discontinuation or interruption of dosing with Tykerb may be considered when a patient develops toxicity greater than or equal to grade 2 on the NCI CTCAE. Dosing can be restarted, when the toxicity improves to grade 1 or less, at 1000 mg/day when administered with trastuzumab, 1250 mg/day when administered with dapecitabine or 1500 mg/day when administered with an aromatase inhibitor. If toxicity recurs, then Tykerb should be restarted at a lower dose (750 mg/day when administered with trastuzumab, 1000 mg/day when administered with capecitabine or 1250 mg/day when administered with an aromatase inhibitor).

No dose adjustment is necessary in patients with mild to moderate renal impairment. Caution is advised in patients with severe renal impairment as there is no experience of Tykerb in this population (see section 5.2). Hepatic impairmen

Tykerb should be discontinued if changes in liver function are severe and patients should not be retreated (see section 4.4). Administration of Tykerb to patients with moderate to severe hepatic impairment should be undertaken with caution due to increased exposure to the medicinal product. Insufficient data are available in patients with hepatic impairment to provide a dose adjustment recommendation (see section 5.2).

There are limited data on the use of Tykerb / capecitabine and Tykerb / trastuzumab in patients aged ≥ 65 years.

In the phase III clinical study of Tykerb in combination with letrozole, of the total number of hormone receptor positive metastatic breast cancer patients (Intent to treat population N=642), 4% were ≥ 65 years of age. No overall differences in efficacy and safety of the combination of Tykerb and letrozole were observed

Paediatric population The safety and efficacy of Tykerb in children below the age of 18 years have not yet been established. No

Method of administrationThe daily dose of Tykerb should not be divided. Tykerb should be taken either at least one hour before, or at least one hour after food. To minimise variability in the individual patient, administration of Tykerb should be standardised in relation to food intake, for example always to be taken one hour before a meal (see sec tions 4.5 and 5.2 for information on absorpt

Missed doses should not be replaced and the dosing should resume with the next scheduled daily dose (see Consult the full prescribing information of the co-administered medicinal product for relevant details of their posology including any dose reductions, contraindications and safety information

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

ween these patients and patients < 65 years of age.

crossover study in subjects with advanced solid tumours.

4.4 Special warnings and precautions for useData have shown that Tykerb combined with chemotherapy is less effective than trastuzumab when combined with chemotherapy.

Lapatinib has been associated with reports of decreases in LVEF (see section 4.8). Lapatinib has not been evaluated in patients with symptomatic cardiac failure. Caution should be taken if Tyker is to be administered to patients with conditions that could impair left ventricular function (including co-administration with potentially cardiotoxic medicinal products). Evaluation of cardiac function, including LVEF determination, should be conducted for all patients prior to initiation of treatment with Tykerb to ensure that the patient has a baseline LVEF that is within the institutions normal limits. LVEF should continue to be evaluated during treatment with Tykerb to ensure that LVEF does not decline to an unacceptable level (see section 4.2). In some cases, LVEF decrease may be severe and lead to cardiac failure. Fatal cases have been reported, causality of the deaths is uncertain. In studies across the clinical development programme for lapatinib, cardiac events including LVEF decreases were reported in approximately 1% of patients. Symptomatic LVEF decreases were observed in approximately 1% of patients. Symptomatic LVEF decreases were observed in approximately 0.3% of patients who received lapatinib. However, when lapatinib was administered in combination with trastuzumab in the metastatic setting, the incidence of cardiac events including LVEF decreases was higher (7%) versus the lapatinib alone arm (2%) in the pivotal trial. The cardiac events observed in this study were comparable in nature and severity to those previously seen with lapatinib. A concentration-dependent increase of the QTc interval was demonstrated in a dedicated placebo-controlled

Caution should be taken if Tykerb is administered to patients with conditions that could result in prolongation Caution should be taken if Tykerb is administered to patients with conditions that could result in prolongation of QTc (including hypokalemia, hypomagnesemia, congenital long QT syndrome), co-administration of of ther medicinal product known to cause QT prolongation, or conditions that increase the exposure of lapatinib, such as co-administration of strong CYP3A4 inhibitors. Hypokalemia or hypomagnesemia should be corrected prior to treatment. Electrocardiograms with QT measurement should be performed prior to and one to two weeks after the start of Tykerb therapy. When clinically indicated, e.g. after initiation of a concomitant treatment that might affect QT or that may interact with lapatinib, ECG measurement should also be considered.

Interstitial lung disease and pneumonitis
Lapatinib has been associated with reports of pulmonary toxicity including interstitial lung disease and pneumonitis (see section 4.8). Patients should be monitored for symptoms of pulmonary toxicity (dyspnoea, cough, fever) and treatment discontinued in patients who experience symptoms which are NCI CTCAE grade 3 or greater. Pulmonary toxicity may be severe and lead to respiratory failure. Fatal cases have been rep causality of the deaths is uncertain.

Hepatotoxicity
Hepatotoxicity has occurred with Tykerb use and may in rare cases be fatal. The hepatotoxicity may occur
Hepatotoxicity has occurred with Tykerb use and may in rare cases be fatal. The hepatotoxicity may occur
Hepatotoxicity
Hepato days to several months after initiation of treatment. At the initiation of treatment, patients should be advised of the potential for hepatotoxicity. Liver function (transaminases, bilirubin and alkaline phosphatase) should be monitored before the initiation of treatment and monthly thereafter, or as clinically indicated. Tykerb dosing should be discontinued if changes in liver function are severe and patients should not be retreated. Patients who carry the HLA alleles DQA1*02:01 and DRB1*07:01 have increased risk of Tykerb-associated hepatotoxicity. In a large, randomised clinical trial of Tykerb monotherapy (n=1,194), the cumulative frequency of severe liver injury (ALT >5 times the upper limit of normal, NCI CTCAE grade 3) at 1 year of treatment was 2.8% overall. The cumulative frequency in DQA1 *02:01 and DRB1 *07:01 allele carriers was 10.3% and in non-carriers was 0.5%. Carriage of the HLA risk alleles is common (15 to 25%) in Caucasian, Asian, African and Hispanic populations but lower (1%) in Japanese populations. Caution is warranted if Tykerb is prescribed to patients with moderate or severe hepatic impairment (see

Diarrhoea, including severe diarrhoea, has been reported with Tykerb treatment (see section 4.8). Diarrhoea can be potentially life-threatening if accompanied by dehydration, renal insufficiency, neutropenia and/or elec-trolyte imbalances and fatal cases have been reported. Diarrhoea generally occurs early during Tykerb treat-ment, with almost half of those patients with diarrhoea first experiencing it within 6 days. This usually lasts 4-5 days. Tykerb-induced diarrhoea is usually low-grade, with severe diarrhoea of NCI CTCAE grades 3 and 4 occurring in <10% and <1% of patients, respectively. At the start of therapy, the patients bowel pattern and

sections 4.2 and 5.2). Caution is advised if Tykerb is prescribed to patients with severe renal impairment (see sections 4.2 and 5.2).

any other symptoms (e.g. fever, cramping pain, nausea, vomiting, dizziness and thirst) should be determined, to allow identification of changes during treatment and to help identify patients at greater risk of diarrhoea. Patients should be instructed to promptly report any change in bowel patterns. In potentially severe cases of diarrhoea the measuring of neutrophil counts and body temperature should be considered. Proactive management of diarrhoea with anti-diarrhoeal medicinal product is important. Severe cases of diarrhoea may require administration of oral or intravenous electrolytes and fluids, use of antibiotics such as fluoroguinolones (especially if diarrhoea is persistent beyond 24 hours, there is fever, or grade 3 or 4 neutropenia) and inter ruption or discontinuation of Tykerb therapy (see section 4.2 – dose delay and dose reduction –diarrhoea).

Serious cutaneous reactions have been reported with Tykerb. If erythema multiforme or life-threatening reactions such as Stevens-Johnson syndrome, or toxic epidermal necrolysis (e.g. progressive skin rash often with blisters or mucosal lesions) are suspected, discontinue treatment with Tykerb.

Concomitant treatment with inhibitors or inducers of CYP3A4
Concomitant treatment with inducers of CYP3A4 should be avoided due to risk of decreased exposure to lapatinib (see section 4.5).

Concomitant treatment with strong inhibitors of CYP3A4 should be avoided due to risk of increased exposure to lapatinib (see section 4.5).

Coadministration of Tykerb with orally administered medicinal products with narrow therapeutic windows that are substrates of CYP3A4 and /or CYP2C8 should be avoided (see section 4.5). Concomitant treatment with substances that increase gastric pH should be avoided, as lapatinib solubility and absorption may decrease (see section 4.5).

4.5 Interaction with other medicinal products and other forms of interaction Effects of other medicinal products on lapatinib

Lapatinib is predominantly metabolised by CYP3A (see section 5.2). In healthy volunteers receiving ketoconazole, a strong CYP3A4 inhibitor, at 200 mg twice daily for 7 days, systemic exposure to lapatinib (100 mg daily) was increased approximately 3.6-fold, and half-life increased 1.7-fold. Co-administration of Tykerb with strong inhibitors of CYP3A4 (e.g. ritonavir, saquinavir, telithromycin, ketoconazole, itraconazole, voriconazole, posaconazole, nefazodone) should be avoided. Co-administration of Tykerb with moderate inhibitors of CYP3A4 should proceed with caution and clinical adverse reactions ould be carefully monitored.

In healthy volunteers receiving carbamazepine, a CYP3A4 inducer, at 100 mg twice daily for 3 days and 200 mg twice daily for 17 days, systemic exposure to lapatinib was decreased approximately 72%. Coadministration of Tykerb with known inducers of CYP3A4 (e.g. rifampicin, rifabutin, carbamazepine, phenytoin or Hypericum perforatum [St John's Wort]) should be avoided or Hypericum perforatum [St John's Wort]) should be avoided.
Lapatinib is a substrate for the transport proteins Pgp and BCRP. Inhibitors (ketoconazole, itraconazole, quinidine, verapamil, cyclosporine, erythromycin) and inducers (rifampicin, St John's Wort) of these proteins may alter the exposure and/or distribution of lapatinib (see section 5.2).

The solubility of lapatinib is pH-dependent. Concomitant treatment with substances that increase gastric pH

should be avoided, as lapatinib solubility and absorption may decrease. Pre-treatment with a proton pump inhibitor (esomeprazole) decreased lapatinib exposure by an average of 27% (range: 6% to 49%). This effect decreases with increasing age from approximately 40 to 60 years.

Effects of lapatinib on other medicinal products
Lapatinib inhibits CYP3A4 *in vitro* at clinically relevant concentrations. Co-administration of Tykerb with orally administered midazolam resulted in an approximate 45% increase in the AUC of midazolam. There was no clinically meaningful increase in AUC when midazolam was dosed intravenously. Co-administration of Tykerb with orally administered medicinal products with narrow therapeutic windows that are substrates of CYP3A4 (e.g. cisapride, pimozide and quinidine) should be avoided (see sections 4.4 and 5.2). Lapatinib inhibits CYP2C8 in vitro at clinically relevant concentrations. Co-administration of Tykerb with medicinal products with narrow therapeutic windows that are substrates of CYP2C8 (e.g. repaglinide) should be avoided (see sections 4.4 and 5.2).

Co-administration of lapatinib with intravenous paclitaxel increased the exposure of paclitaxel by 23%, due to lapatinib inhibition of CYP2C8 and/or Pgp. An increase in the incidence and severity of diarrhoea and neutropenia has been observed with this combination in clinical studies. Caution is advised if lapatinib is co-administered with paclitaxel. Co-administration of lapatinib with intravenously administered docetaxel did not significantly affect the AUC or Cmax of either active substance. However, the occurrence of docetaxel-induced neutropenia was increased

Co-administration of Tykerb with irinotecan (when administered as part of the FOLFIRI regimen) resulted in an approximate 40% increase in the AUC of SN-38, the active metabolite of irinotecan. The precise mechanism approximate 40% increase in the AUC of SN-38, the active metabolite of innotecan. The precise mechanism of this interaction is unknown, but it is assumed to be due to inhibition of one or more transport proteins by lapatinib. Adverse reactions should be carefully monitored if Tykerb is co-administered with irinotecan, and a reduction in the dose of irinotecan should be considered. Lapatinib inhibits the transport protein Pgp in vitro at clinically relevant concentrations. Co-administration of lapatinib with orally administered digoxin resulted in an approximate 80% increase in the AUC of digoxin. Caution should be exercised when dosing lapatinib concurrently with medicinal products with narrow therapeutic windows that are substrates of Pgp, and a reduction in the dose of the Pgp substrate should be considered.

Lapatinib inhibits the transport proteins BCRP and OATP1B1 in vitro. The clinical relevance of this effect has not been evaluated. It cannot be excluded that lapatinib will affect the pharmacokinetics of substrates of BCRP (e.g. topotecan) and OATP1B1 (e.g. rosuvastatin) (see section 5.2). Concomitant administration of Tykerb with capecitabine, letrozole or trastuzumab did not meaningfully alter the pharmacokinetics of these medicinal products (or the metabolites of capecitabine) or lapatinib.

Interactions with food and drink The bioavailability of lapatinib is increased up to about 4 times by food, depending on e.g. the fat content in

all. Furthermore, depending on type of food the bioavailability is approximately 2-3 times higher when the staken 1 hour after food compared with 1 hour before the first meal of the day (see sections 4.2and 3.2). Grapefruit juice may inhibit CYP3A4 in the gut wall and increase the bioavailability of lapatinib and should therefore be avoided during treatment with Tykerb.

4.6 Fertility, pregnancy and lactation

Women of childbearing potentialWomen of childbearing potential should be advised to use adequate contraception and avoid becoming pregnant while receiving treatment with Tykerb and for at least 5 days after the last dose.

There are no adequate data from the use of Tykerb in pregnant women. Studies in animals have shown reproductive toxicity (see section 5.3). The potential risk for humans is not known.

Tykerb should not be used during pregnancy unless clearly necessary. Breast-feeding
The safe use of Tykerb during breast-feeding has not been established. It is not known whether lapatinib is excreted in human milk. In rats, growth retardation was observed in pups which were exposed to lapatinib via breast fileding must be discontinued in women who are receiving therapy with Tykerb and for at least 5 days after the last dose.

There are no adequate data from the use of Tykerb in women of childbearing potential

4.7 Effects on ability to drive and use machines A detrimental effect on such activities cannot be predicted from the pharmacology of lapatinib. The clinical status of the patient and the adverse event profile of lapatinib should be borne in mind when considering the

patient's ability to perform tasks that require judgement, motor or cognitive skills.

Summary of the safety profile

The safety of lapatinib has been evaluated as monotherapy or in combination with other chemotherapies for various cancers in more than 20,000 patients, including 198 patients who received lapatinib in combination with capecitabine, 149 patients who received lapatinib in combination with leaves the received lapatinib in combination with letrozole (see section 5.1).

The most common adverse reactions (>25%) during therapy with lapatinib were gastrointestinal events (such as diarrhoea, nausea, and vomiting) and rash. Palmar-plantar erythrodysesthesia [PPE] was also common (>25%) when lapatinib was administered in combination with capecitabine. The incidence of PPE was similar in the lapatinib plus capecitabine and capecitabine alone treatment arms. Diarrhoea was the most common adverse reaction resulting in discontinuation of treatment when lapatinib was administered in combination with capecitabine, or with letrozole.

No additional adverse reactions were reported to be associated with lapatinib in combination with trastuzumand additional adverse reactions were reported to be associated with lapatinio in combination with trastituzinab. There was an increased incidence of cardiact otxicity, but these events were comparable in nature and severity to those reported from the lapatinib clinical programme (see section 4.4 − cardiac toxicity). These data are based on exposure to this combination in 149 patients in the pivotal trial. The following convention has been utilised for the classification of frequency: very common (≥1/10), common (≥1/100 to <1/10), uncommon (≥1/1,000 to <1/100), rare (≥1/10,000 to <1/1,000 and very rare (<1/10,000), not known (cannot be estimated from the available data). Within each frequency grupping, understable effects are presented in order of decreasing seriousness.

| Immune system | disorders |
|------------------|---|
| Rare | Hypersensitivity reactions including anaphylaxis (see section 4.3) |
| | I nutrition disorders |
| Very common | Anorexia |
| Psychiatric disc | orders |
| Very common | Insomnia* |
| Nervous system | n disorders |
| Very common | Headache [†] |
| Common | Headache* |
| Cardiac disorde | ers |
| Common | Decreased left ventricular ejection fraction (see section 4.2 - dose reduction – cardiac events and section 4.4). |
| Not known | Ventricular arrhythmias/Torsades de Pointes, electrocardiogram QT prolonged** |
| Vascular disord | ers |
| Very common | Hot flush† |
| Respiratory, the | oracic and mediastinal disorders |
| Very common | Epistaxis†, cough†, dyspnoea†. |
| Uncommon | Interstitial lung disease/pneumonitis. |
| Not known | Pulmonary arterial hypertension**. |
| Gastrointestinal | disorders |
| Very common | Diarrhoea, which may lead to dehydration (see section 4.2 - dose delay and dose reduction – other toxicities and section 4.4), nausea, vomiting, dyspepsia*, stomatitis*, constipation*, abdominal pain*. |
| Common | Constipation [†] |
| Hepatobiliary d | isorders |
| Common | Hyperbilirubinaemia, hepatotoxicity (see section 4.4). |
| Skin and subcur | taneous tissue disorders |
| Very common | Rash (including dermatitis acneiform) (see section 4.2 - dose delay and dose reduction – other toxicities), dry skin*†, palmar-plantar erythrodysaesthesia*, alopecia†, pruritus†. |
| Common | Nail disorders including paronychia. |
| Not known | Serious cutaneous reactions, including Stevens Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN)* * |
| Musculoskeleta | I and connective tissue disorders |
| Very common | Pain in extremity*†, back pain*†, arthralgia†. |
| General disorde | ers and administration site conditions |
| Very common | Fatigue, mucosal inflammation*, asthenia†. |

These adverse reactions were observed when lapatinib was administered in combination with capecit-

These adverse reactions were observed when lapatinib was administered in combination with letrozole.
 ** Adverse reactions from spontaneous reports and literature

Description of selected adverse reactions Description of selected adverse reactions
Decreased left ventricular ejection fraction and QT interval prolongation
Left ventricular ejection fraction (LVEF) decreases have been reported in approximately 1% of patients receiving lapatinib and were asymptomatic in more than 70% of cases. LVEF decreases resolved or improved in more than 70% of cases, in approximately 60% of these on discontinuation of treatment with lapatinib, and in approximately 40% of cases lapatinib was continued. Symptomatic LVEF decreases were observed in approximately 0.3% of patients who received lapatinib monotherapy or in combination with other anti-cancer medicates. inal products. Observed adverse reactions included dyspnoea, cardiac failure and palpitations. Overall 58 % of these symptomatic patients recovered. LVEF decreases were reported in 2.5 % of patients who received lapatinib in combination with capecitabine, as compared to 1.0 % with capecitabine alone. LVEF decreases were reported in $3.1\,\%$ of patients who received lapatinib in combination with letrozole as compared to $1.3\,\%$ of patients receiving letrozole plus placebo. LVEF decreases were reported in $6.7\,\%$ of patients who received Lapatinis in combination with trastuzumab, as compared to $2.1\,\%$ of patients who received lapatinis alone. A concentration dependent increase in QTcF (maximum mean $\Delta\Delta$ QTcF 8.75 ms; 90% Cl 4.08, 13.42) was observed in a dedicated QT study in patients with advanced solid tumours (see section 4.4).

Diarrhoea occurred in approximately 65 % of patients who received lapatinib in combination with capecitabine, in 64 % of patients who received lapatinib in combination with letrozole and in 62 % of patients who received lapatinib in combination with trastuzumab. Most cases of diarrhoea were grade 1 or 2 and did not result in discontinuation of treatment with lapatinib. Diarrhoea responds well to proactive management (see section 4.4). However, a few cases of acute renal failure have been reported secondary to severe dehydration

Rash occurred in approximately 28 % of patients who received lapatinib in combination with capecitabine, in 45 % of patients who received lapatinib in combination with letrozole and in 23 % of patients who received lapatinib in combination with trastuzumab. Rash was generally low grade and did not result in discontinuation of treatment with lapatinib. Prescribing physicians are advised to perform a skin examination prior to treat-

ment and regularly during treatment. Patients experiencing skin reactions should be encouraged to avoid exposure to sunlight and apply broad spectrum sunscreens with a Sun Protection Factor (SPF) ≥ 30. If a skin reaction occurs a full body examination should be performed at every visit until one month after resolution. Patients with extensive or persistent skin reactions should be referred to a dermatologist.

HepatotoxicityThe risk of lapatinib-induced hepatotoxicity was associated with carriage of the HLA alleles DQA1*02:01 and DRB1*07:01 (see section 4.4).

Reporting of suspected adverse reactions
Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions

There is no specific antidote for the inhibition of EGFR (ErbB1) and/or HER2 (ErbB2) tyrosine phosphorylation. The maximum oral dose of lapatinib that has been administered in clinical studies is 1800 mg once daily. Asymptomatic and symptomatic cases of overdose have been reported in patients being treated with Tykerb. In patients who took up to 5000 mg of lapatinib, symptoms observed include known lapatinib associated events (see section 4.8) and in some cases sore scalp and/or mucosal inflammation. In a single case of a patient who took 9000 mg of Tykerb, sinus tachycardia (with otherwise normal ECG) was also observed. Lapatinib is not significantly renally excreted and is highly bound to plasma proteins, therefore haemodialysis would not be expected to be an effective method to enhance the elimination of lapatinib. Further management should be as clinically indicated or as recommended by the national poisons centre, where available.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties
Pharmacotherapeutic group: Antineoplastic agent, other antineoplastic agents, protein kinase inhibitor, ATC code: L01XE07

Mechanism of action

Lapatinib, a 4-anilinoquinazoline, is an inhibitor of the intracellular tyrosine kinase domains of both EGFR (ErbB1) and of HER2 (ErbB2) receptors (estimated Ki¹⁰⁰ values of 3nM and 13nM, respectively) with a slow offrate from these receptors (half-life greater than or equal to 300 minutes). Lapatinib inhibits ErbB-driven

tumour cell growth in vitro and in various animal models.

The combination of lapatinib and trastuzumab may offer complementary mechanisms of action as well as possible non-overlapping mechanisms of resistance. The growth inhibitory effects of lapatinib were evaluated in trastuzumab-conditioned cell lines. Lapatinib retained significant activity against HER2-amplified breast cancer cell lines selected for long-term growth in trastuzumab-containing medium in vitro and was synergistic in combination with trastuzumab in these cell lines.

Clinical efficacy and safety Combination treatment with Tykerb and capecitabine

The efficacy and safety of Tykerb in combination with capecitabine in breast cancer patients with good performance status was evaluated in a randomised, phase III study. Patients eligible for enrolment had HER2-overexpressing, locally advanced or metastatic breast cancer, progressing after prior tartment that included taxanes, anthracyclines and trastuzumab. LVEF was evaluated in all patients (using echocardiogram [Echo] or multi gated acquisition scan [MUGAI) prior to initiation of treatment with Tykerb to ensure baseline LVEF was within the institutions normal limits. In the clinical study LVEF was monitored at approximately eight week intervals during treatment with Tykerb to ensure it did not decline to below the institutions lower limit of normal. The majority of LVEF decreases (greater than 60 % of events) were observed during the first nine weeks of treatment, however limited data was available for long term exposure.

treatment, however limited data was available for long term exposure. Patients were randomised to receive either Tykerb 1250 mg once daily (continuously) plus capecitabine (2000 mg/m²/day on days 1-14 every 21 days), or to receive capecitabine alone (2500 mg/m²/day on days 1-14 every 21 days). The primary endpoint was time to progression (TTP). Assessments were undertaken by the study investigators and by an independent review panel, blinded to treatment. The study was halted based on the results of a pre-specified interim analysis that showed an improvement in TTP for patients terim analysis and the end of the enrolment. Investigator analysis on data at the end of enrolment is need in Table 1. receiving Tykerb plus capecitabine. An additional 75 patients were enrolled in the study between the time of

Table 1 Time to Progression data from Study EGF100151 (Tykerb / capecitabine)

| | Investigator assessment | | |
|----------------------|---|--|--|
| | Tykerb (1,250 mg/day)+ capecitabine (2,000 mg/m²/day, days 1-14 q21 days) | Capecitabine (2,500 mg/m²/day, days 1-14 q21 days) | |
| | (N = 198) | (N = 201) | |
| Number of TTP events | 121 | 126 | |
| Median TTP, weeks | 23.9 | 18.3 | |
| Hazard Ratio | 0.72 | | |
| (95% CI) | (0.56, 0.92) | | |
| p value | 0.008 | | |

The independent assessment of the data also demonstrated that Tykerb when given in combination with capecitabine significantly increased time to progression (Hazard Ratio 0.57~[95~%~Cl~0.43,~0.77]~p=0.0001)

Results of an updated analysis of the overall survival data to 28 September 2007 are presented in Table 2.

Table 2 Overall survival data from Study EGF100151 (Tykerb / canecitabine

| | Tykerb (1,250 mg/day)+ capecitabine (2,000 mg/m²/ day, days 1-14 q21 days) | Capecitabine (2,500 mg/m² day, days 1-14 q21 days) | |
|--------------------------------|--|--|--|
| | (N = 207) | (N = 201) | |
| Number of subjects who died | 148 | 154 | |
| Median overall survival, weeks | 74.0 | 65.9 | |
| Hazard Ratio | 0.9 | | |
| (95% CI) | (0.71, 1.12) | | |
| p value | 0.3 | | |

On the combination arm, there were 4 (2%) progressions in the central nervous system as compared with the 13 (6%) progressions on the capecitabline alone arm.

Data are available on the efficacy and safety of Tykerb in combination with capecitabine relative to trastu zumab in combination with capecitabine. A randomised Phase III study (EGF111438) (N=540) compared the effect of the two regimens on the incidence of CNS as site of first relapse in women with HER2 overexpressing metastatic breast cancer. Patients were randomised to either Tykerb 1250 mg once daily (continuously) plus capecitabine (2000 mg/m²/day on days 1-14 every 21 days), or trastuzumab (loading dose of 8mg, kg followed by 6mg/kg q3 weekly infusions) plus capecitabine (2500mg/m²/day, days 1-14, every 21 days). Randomisation was stratified by prior trastuzumab treatment and number of prior treatments for metastatic disease. The study was halted as the interim analysis (N=475) showed a low incidence of CNS events and, superior efficacy of the trastuzumab plus capecitabine arm in terms of progression-free survival and overall survival (see results of final analysis in Table 3).

In the Tykerb plus capecitabine arm 8 patients (3.2%) experienced CNS as site of first progression, compared with 12 patients (4.8%) in the trastuzumab plus capecitabine arm. Lapatinib effect on CNS metastasis

Lapatinib has in terms of objective responses demonstrated modest activity in the treatment of established CNS metastases. In the prevention of CNS metastases in the metastatic and early breast cancer settings the observed activity was limited.

Analyses of Investigator-Assessed Progression-Free Survival and Overall Surviva

| | Investigator- | Assessed PFS | Overall Survival | | |
|--|--|--|--|---|--|
| | Tykerb (1,250 mg/day) + capecitabine (2,000 mg/m²/day, days 1-14 q21 days) | Trastuzumab (loading dose of 8mg/kg followed by 6mg/kg q3 weekly infusions) + capecitabine (2,500 mg/m²/ day, days 1-14 q21 days) | Tykerb (1,250 mg/day) + capecitabine (2,000 mg/m²/day, days 1-14 q21 days) | Trastuzumab (loading dose of 8mg/kg followed by 6mg/kg q3 weekly infusions capecitabine (2,500 mg/m²/day, days 1-14 q21 days) | |
| ITT population | - | | | | |
| N | 271 | 269 | 271 | 269 | |
| Number (%) with Event ¹ | 160 (59) | 134 (50) | 70 (26) | 58 (22) | |
| Kaplan-Meier estimate, months ^a | | | | | |
| Median (95% CI) | 6.6 (5.7, 8.1) | 8.0 (6.1, 8.9) | 22.7 (19.5, -) | 27.3 (23.7, -) | |
| Stratified Hazard ratio ^b | | | | | |
| HR (95% CI) | 1.30 (1.04, 1.64) | | 1.34 (0.95, 1.90) | | |
| p-value | 0.0 |)21 | 0.095 | | |
| Subjects who h | ad received prior tra | stuzumab* | | | |
| N | 167 | 159 | 167 | 159 | |
| Number (%) with Event ¹ | 103 (62) | 86 (54) | 43 (26) | 38 (24) | |
| Median (95% CI) | 6.6 (5.7, 8.3) | 6.1 (5.7, 8.0) | 22.7 (20.1,-) | 27.3 (22.5, 33.6) | |
| HR (95% CI) | 1.13 (0.8 | 35, 1.50) | 1.18 (0. | .76, 1.83) | |
| Subjects who h | ad not received prior | r trastuzumab* | | | |
| N | 104 | 110 | 104 | 110 | |
| Number (%) with Event ¹ | 57 (55) | 48 (44) | 27 (26) | 20 (18) | |
| Median (95% CI) | 6.3 (5.6, 8.1) | 10.9 (8.3, 15.0) | NE ² (14.6, -) | NE ² (21.6, -) | |
| HR (95% CI) | 1.70 (1.15, 2.50) | | 1.67 | 7 (0.94, 2.96) | |

. PFS was defined as the time from randomisation to the earliest date of disease progression or death

. PrS was defined as the time from randomisation to the earliest date of disease progression or dea from any cause, or to the date of censor.

Pike estimate of the treatment hazard ratio, <1 indicates a lower risk for Tykerb plus capecitabine compared with Trastuzumab plus capecitabine.

PS event is Progressed or Died and OS event is Died due to any cause.

NE=Median was not reached.

Post hoc analysis

Combination treatment with Tykerb and trastuzumab

The efficacy and safety of lapatinib in combination with trastuzumab in metastatic breast cancer were evaluated in a randomised trial. Eligible patients were women with Stage IV ErbB2 gene amplified (or protein overexpressing) metastatic breast cancer who had been exposed to treatment with anthracyclines and taxanes. In addition, per the protocol, patients were to be reported by the investigators as having progressed on their most recent trastuzumab containing regimen in the metastatic setting. The median number of prior trastucumab-containing regimens was three. Patients were randomised to receive either oral lapatinib 1,000 mg once daily plus trastuzumab 4 mg/kg administered as an intravenous (IV) loading dose, followed by 2 mg/kg IV weekly (N = 148), or oral lapatinib 1,500 mg once daily (N = 148). Patients who had objective disease progression after receiving at least 4 weeks of treatment with lapatinib monotherapy were eligible to crossover to combination therapy. Of the 148 patients who received monotherapy treatment, 77 (52%) patients elected at the time of disease progression to receive combination treatment.

rogression-free survival (PFS) was the primary endpoint of the study with response rate and overall survival (OS) as secondary endpoints. The median age was 51 years and 13% were 65 years or older. Ninety-four percent (94%) were Caucasian. Most patients in both treatment arms had visceral disease (215 [73%] patients overall). In addition, 150 [50%] of patients were hormone receptor negative. A summary of efficacy endpoints and overall survival data is provided in Table 4. Subgroup analysis results based on predefined stratification factor (hormone receptor status) is also shown in Table 5.

Table 4 Progression-free survival and overall survival data (Tykerb / trastuzumab)

| | Lapatinib plus trastuzumab (N=148) | Lapatinib alone (N=148) |
|--|--|----------------------------|
| Median PFS ¹ , weeks (95% CI) | 12.0 (8.1, 16.0) | 8.1 (7.6, 9.0) |
| Hazard Ratio (95% CI) | 0.73 (0. | 57, 0.93) |
| P value | 0.008 | |
| Response Rate, % (95% CI) | 10.3 (5.9, 16.4) | 6.9 (3.4, 12.3) |
| Died | 105 | 113 |
| Median overall survival ¹ , months (95% CI) | 14.0 (11.9, 17.2) | 9.5 (7.6, 12.0) |
| Hazard Ratio (95% CI) | 0.74 (0.57, 0.97) | |
| P value | 0.026 | |

PFS = progression-free survival; CI = confidence interval.

Summary of PFS and OS in Studies with Hormone Receptor negative

| Median PFS | | Median OS | |
|--|--|-----------------------|--|
| Lap+Tras 15.4 wks (8.4, 16.9) | | 17.2 mos (13.9, 19.2) | |
| Lap 8.2 wks (7.4, 9.3) 8.9 mos (6.7, 11.8 | | 8.9 mos (6.7, 11.8) | |
| HR (95% CI) 0.73 (0.52, 1.03) | | 0.62 (0.42, 0.90) | |

Combination treatment with Tykerb and letrozole Tykerb has been studied in combination with letrozole for the treatment of postmenopausal women with

hormone receptor-positive (oestrogen receptor [ER] positive and / or progesterone receptor [PgR] positive) advanced or metastatic breast cancer. The Phase III study (EGF30008) was randomised, double-blind, and placebo controlled. The study enrolled patients who had not received prior therapy for their metastatic disease.

In the HER2-overexpressing population, only 2 patients were enrolled who had received prior trastuzumab, 2 patients had received prior aromatase inhibitor therapy, and approximately half had received tamoxifen. Patients were randomised to letrozole 2.5 mg once daily plus Tykerb 1500 mg once daily or letrozole with placebo. Randomisation was stratified by sites of disease and by time from discontinuation of prior adjuvant anti-oestrogen therapy. HER2 receptor status was retrospectively determined by central laboratory testing. Of all patients randomised to treatment, 219 patients had tumours overexpressing the HER2 receptor, and

HER2-negative tumours, and a total of 115 patients whose tumour HER2 status was unconfirmed (no tumour sample, no assay result, or other reason). In patients with HER2-overexpressing MBC, investigator-determined progression-free survival (PFS) was sig-nificantly greater with letrozole plus Tykerb compared with letrozole plus placebo. In the HER2-negative pop-ulation, there was no benefit in PFS when letrozole plus Tykerb was compared with letrozole plus placebo

this was the pre-specified primary population for the analysis of efficacy. There were 952 patients with

Progression Free Survival data from Study EGF30008 (Tykerb / letr

| | HER2-Overexpressing Population | | HER2-Negative Population | |
|----------------------------------|---|---------------------------------------|---|---------------------------------------|
| | N = 111 | N = 108 | N = 478 N = 474 | N = 474 |
| | Tykerb 1500 mg / day + Letrozole 2.5 mg /day | Letrozole 2.5 mg /day + placebo | Tykerb 1500 mg / day + Letrozole 2.5 mg /day | Letrozole 2.5 mg /day + placebo |
| Median PFS, weeks (95% CI) | 35.4 (24.1, 39.4) | 13.0 (12.0, 23.7) | 59.7 (48.6, 69.7) | 58.3 (47.9, 62.0) |
| Hazard Ratio | 0.71 (0.53, 0.96) | | 0.90 (0.77, 1.05) | |
| P-value | 0.019 | | 0.188 | |
| Objective Response Rate (ORR) | 27.9% | 14.8% | 32.6% | 31.6% |
| Odds Ratio | 0.4 (0.2, 0.9) | 0.4 (0.2, 0.9) | | |
| P-value | 0.021 | 0.021 | | |
| Clinical Benefit Rate (CBR) | 47.7% | 28.7% | 58.2% | 31.6% |
| Odds Ratio | 0.4 (0.2, 0.8) | | 1.0 (0.7, 1.2) | |
| P-value | 0.003 | | 0.199 | |

At the time of the final PFS analysis (with median follow-up of 2.64 years), the overall survival data were not mature and there was no significant difference between treatment groups in the HER2-positive population; this had not changed with additional follow-up (>7.5 years median follow-up time: Table 7).

HER2 overexpression = IHC 3+ and/or FISH positive; HER2 negative = IHC 0, 1+ or 2+ and/or FISH negative Clinical Benefit Rate was defined as complete plus partial response plus stable disease for ≥6 months.

Table 7 Overall Survival (OS) results from study EGF30008 (in the HER2-positive population

| | Tykerb 1500 mg / day + Letrozole 2.5 mg /day N=111 | Letrozole 2.5 mg /day + placebo N=108 | |
|---|---|---|--|
| Pre-planned OS analysis (conducted | at the time of the final PFS | S analysis, 03 June 2008) | |
| Median Follow-up (yrs) | 2,64 | 2,64 | |
| Deaths (%) | 50 (45) | 54 (50) | |
| Hazard Ratio ^a (95% CI), p-value ^b | 0,77 (0,52; 1,14); 0,185 | | |
| Final OS analysis (post-hoc analysis, | 07 August 2013) | | |
| Median Follow-up (yrs) | 7,78 | 7,55 | |
| Deaths (%) | 86 (77) | 78 (72) | |
| Hazard Ratio (95% CI), p-value | 0,97 (0,07; 1,33); 0,848 | | |

Median values from Kaplan-Meier analysis; HR and p-values from Cox regression models adjusting for

important prognostic factors.

a. Estimate of the treatment hazard ratio, where <1 indicates a lower risk with letrozole 2.5 mg + lapatinib 1500 mg compared with letrozole 2.5 mg + placebo.

b. P-value from Cox regression model, stratifying for site of disease and prior anti-adjuvant therapy

The efficacy and safety of Tykerb in combination with an aromatase inhibitor were further confirmed in another Phase III study (Study EGF114299). Patients enrolled were postmenopausal women who had hormone receptor-positive/HER2-positive metastatic breast cancer which had progressed after prior trastuzumab-containing chemotherapy regimen and endocrine therapies. The study was primarily designed to evaluate a potential chemotherapy regimen and endocrine interaples. The Study was primarily designed to evaluate a potential benefit in PFS of dual HER2 blockade (Tykerb + trastuzumab) versus single HER2 blockade (trastuzumab), both in combination with an aromatase inhibitor (Al). The study included a third arm of Tykerb + Al. A total of 355 patients were randomised in a 1:1:1 ratio to Tykerb 1000 mg + trastuzumab (loading dose of 8 mg/kg followed by a maintenance dose of 6 mg/kg IV q3weeks) + Al (N=120), or trastuzumab (loading dose of 8 mg/kg followed by a maintenance dose of 6 mg/kg IV q3weeks) + Al arm (N=117), or Tykerb 1500 mg + Al (N=118).

Table 8 Progression Free Survival data from Study EGF114299

| | Tykerb (1500 mg) + Al | Trastuzumab (6 mg/kg) + Al | |
|------------------------------------|-----------------------|----------------------------|--|
| | N=117 | N=118 | |
| Events, n (%) | 74 (63) | 75 (64) | |
| Median PFS, months (95% CI) | 8.3 (5.8, 11.2) | 5.7 (5.5, 8.4) | |
| HR; 95% CI versus trastuzumab + AI | 0.71 (0.51, 0.98) | | |
| P-value | 0.0361 | - | |

Cardiac electrophysiology
The effect of lapatinib on the QT-interval was evaluated in a single-blind, placebo-controlled, single sequence (placebo and active treatment) crossover study in patients with advanced solid tumours (EGF114271) (n=58). During the 4-day treatment period, three doses of matching placebo were administered 12 hours apart in the morning and evening on Day 1 and in the morning on Day 2. This was followed by three doses of lapatinib 2000 mg administered in the same way. Measurements, including electrocardiograms (ECGs) and pharmacokinetic samples, were taken at baseline and at the same time points on Day 2 and Day 4. In the evaluable population (n=37), the maximum mean $\Delta\Delta$ QTcF (90% Cl) of 8.75 ms (4.08, 13.42) was observed 10 hours after ingestion of the third dose of lapatinib 2000 mg. The $\Delta\Delta$ QTcF exceeded the 5 ms threshold and the upper bound 90% Cls exceeded the 10 ms threshold at multiple time points. The results for the pharmacodynamics population (n=52) were consistent with those from the evaluable population (maximum $\Delta\Delta$) TcF (90% Cl) of 7.91 ms (4.13, 11.68) observed 10 hours after ingestion of the third dose of

imum $\Delta\Delta$ QTcF (90% CI) of 7.91 ms (4.13, 11.68) observed 10 hours after ingestion of the third dose of lapatinib 2000 mg). There is a positive relationship between lapatinib plasma concentrations and $\Delta\Delta QTcF$. Lapatinib produced a maximum mean concentration of 3920 (3450.4460) ng/ml (geometric mean/95% CI), exceeding the geometric mean C_{max.ss} and 95% CI values observed following the approved dosing regimens. An additional increase in peak exposure of lapatinib can be expected when lapatinib is taken repeatedly with food (see sections 4.2 and 5.2) or concomitantly with strong CYP3A4 inhibitors. When lapatinib is taken in combination with strong CYP3A4 inhibitors the QTc interval can be expected to be prolonged by 16.1 ms (12.6-20.3 ms) as demonstrating the control of the control

Food effects on lapatinib exposure

strated in a model-based prediction (see section 4.4).

The bioavailability and thereby the plasma concentrations of lapatinib are increased by food, in relation to the content and timing of the meal. Dosing of lapatinib one hour after a meal results in approximately 2-3 times higher systemic exposure, compared to dosing one hour before a meal (see sections 4.5 and 5.2). The European Medicines Agency has waived the obligation to submit the results of studies with Tykerb in all subsets of the paediatric population in the treatment of breast carcinoma (see section 4.2 for inform

5.2 Pharmacokinetic properties

5.2 Pharmacokinetic properties Absorption

The absolute bioavailability following oral administration of lapatinib is unknown, but it is incomplete and variable (approximately 70% coefficient of variation in AUC). Serum concentrations appear after a median lag time of 0.25 hours (range 0 to 1.5 hours). Peak plasma concentrations (c_{mus}) of lapatinib are achieved approximately 4 hours after administration. Daily dosing of 1250 mg produces steady state geometric mean (coefficient of variation) C_{mus} values of 2.43 (76%) µg/ml and AUC values of 36.2 (79%) µg/ml/ml. Systemic exposure to lapatinib is increased when administered with food. Lapatinib AUC values were approximately 3- and 4-fold higher (C_{mus} approximately 2.5 and 3-fold higher) when administered with administration in the fasted state. Systemic exposure to lapatinib is also affected by the timing of administration in rela-

tion in the fasted state. Systemic exposure to lapatinib is also affected by the timing of administration in relation to food intake. Relative to dosing 1 hour before a low fat breakfast, mean AUC values were approximately 2- and 3-fold higher when lapatinib was administered 1 hour after a low fat or high fat meal, respectively Distribution

Lapatinib is highly bound (greater than 99%) to albumin and alpha-1 acid glycoprotein. *In vitro* studies indicate that lapatinib is a substrate for the transporters BCRP (ABCG1) and p-glycoprotein (ABCB1). Lapatinib has also been shown in vitro to inhibit these efflux transporters, as well as the hepatic uptake transporter OATP 1B1, at clinically relevant concentrations (O_{50} values were equal to 2.3 µg/ml). The clinical significance of these effects on the pharmacokinetics of other medicinal products or the pharmacological activity of other anti-cancer medicinal products is not known.

Biotransformation Biotransformation
Lapatinib undergoes extensive metabolism, primarily by CYP3A4 and CYP3A5, with minor contributions from CYP2C19 and CYP2C8 to a variety of oxidated metabolites, none of which account for more than 14% of the dose recovered in the faeces or 10% of lapatinib concentration in plasma.
Lapatinib inhibits CYP3A (Ki 0.6 to 2.3 μg/ml) and CYP2C8 (0.3 μg/ml) in vitro at clinically relevant concentrations. Lapatinib did not significantly inhibit the following enzymes in human liver microsomes: CYP1A2, CYP2C9, CYP2C19, and CYP2D6 or UGT enzymes (in vitro IC₅₀ values were greater than or equal to 6.9 μg/

The half-life of lapatinib measured after single doses increases with increasing dose. However, daily dosing of lapatinib results in achievement of steady state within 6 to 7 days, indicating an effective half-life of 24 hours. Lapatinib is predominantly eliminated through metabolism by CYP3A4/5. Biliary excretion may also contribute to the elimination. The primary route of excretion for lapatinib and its metabolites is in faeces. Recovery of unchanged lapatinib in faeces accounts for a median 27% (range 3 to 67%) of an oral dose. Less than 2% of the administered oral dose (as lapatinib and metabolites) excreted in urine.

Lapatinib pharmacokinetics have not been specifically studied in patients with renal impairment or in patients

undergoing haemodialysis. Available data suggest that no dose adjustment is necessary in patients with mild to moderate renal impairment. Hepatic impairment The pharmacokinetics of lapatinib were examined in patients with moderate (n = 8) or severe (n = 4) hepatic impairment (Child-Pugh scores of 7-9, or greater than 9, respectively) and in 8 healthy control patients. Systemic exposure (AUC) to lapatinib after a single oral 100 mg dose increased approximately 56% and 85% in patients with moderate and severe hepatic impairment, respectively. Administration of lapatinib in patients with hepatic impairment should be undertaken with caution (see sections 4.2 and 4.4).

5.3 Preclinical safety data

5.3 Preclinical safety data
Lapatinib was studied in pregnant rats and rabbits given oral doses of 30, 60, and 120 mg/kg/day. There were no teratogenic effects; however, minor anomalies (left-sided umbilical artery, cervical rib and precocious ossification) occurred in rats at ≥60 mg/kg/day (4 times the expected human clinical exposure). In rabbits, lapatinib was associated with maternal toxicity at 60 and 120 mg/kg/day (8% and 23% of the expected human clinical exposure, respectively) and abortions at 120 mg/kg/day. At ≥60 mg/kg/day there were decreased foetal body weights, and minor skeletal variations. In the rat pre- and postnatal development study, a decrease in pup survival occurred between birth and postnatal day 21 at doses of 60 mg/kg/day or higher (5 times the expected human clinical exposure). The highest no-effect dose for this study was 20 mg/kg/day. In oral carcinogenicity studies with lapatinib, severe skin lesions were seen at the highest doses tested which produced exposures based on AUC up to 2-fold in mice and male rats, and up to 15-fold in female rats, compared to humans given 1250 mg of lapatinib noce daily. There was no evidence of carcinogenicity in mice. In pared to humans given 1250 mg of lapatinib once daily. There was no evidence of carcinogenicity in mice. In rats, the incidence of benign haemangioma of the mesenteric lymph nodes was higher in some groups than in concurrent controls. There was also an increase in renal infarcts and papillary necrosis in female rats at exposures 7 and 10-fold compared to humans given 1250 mg of lapatinib once daily. The relevance of these

findings for humans is uncertain. There were no effects on male or female rat gonadal function, mating, or fertility at doses up to 120 mg/kg/day (females) and up to 180 mg/kg/day (males) (8 and 3 times the expected human clinical exposure, respectively). The effect on human fertility is unknown. Lapatini was not clastogenic or mutagenic in a battery of assays including the Chinese hamster chromosome aberration assay, the Ames assay, human lymphocyte chromosome aberration assay and an *in vivo* rat bone marrow chromosome aberration assav.

6. PHARMACEUTICAL PARTICULARS 6.1 List of excipients

findings for humans is uncertain.

Tablet core Microcrystalline cellulose Povidone (K30) Sodium starch glycolate (Type A) Magnesium stearate **Tablet coating** Titanium dioxide (E171)

Macrogol 400 Polysorbate 80 Iron oxide yellow (E172) Iron oxide red (E172)

6.2 Incompatibilities

6.3 Shelf life The medicinal product must not be used after the expiry date (= EXP) printed on the pack. 6.4 Special precautions for storage

Do not store above 30°C. Keep out of the reach and sight of children

6.5 Nature and contents of container Tykerb is supplied in either blister packs or bottles

Blister packs Tykerb / capecitabine combination posology
Each pack of Tykerb contains 70 film-coated tablets in foil blisters (polyamide / aluminium / polyvinyl chloride / aluminium) of 10 tablets each. Each foil has a perforation down the middle to allow the blisters to be separated into a daily dose of 5 tablets.

Multipacks contain 140 (2 packs of 70) film-coated tablets.

Tykerb / aromatase inhibitor combination posology
Each pack of Tykerb contains 84 film-coated tablets in foil blisters (polyamide / aluminium / polyvinyl chloride

/ aluminium) of 12 tablets each. Each foil has a perforation down the middle to allow the blisters to be sepaated into a daily dose of 6 tablets Tykerb is also supplied in high density polyethylene bottles (HDPE) with a child resistant polypropylene closure containing 70, 84, 105 or 140 film-coated tablets.

6.6 Special precautions for disposalAny unused medicinal product or waste material should be disposed of in accordance with local requirements.

Glaxo Operations UK Limited, Ware, UK for Novartis Pharma AG, Basle, Switzerland 7.Manufactured by:

8. DATE OF REVISION OF THE TEXT

Not all pack sizes may be marketed.

® = registered trademark Novartis Pharma AG, Basle, Switzerland

- A medicament is a product which affects your health, and its consumption contrary to instructions is dan-

gerous for you. Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold

The doctor and the pharmacist are experts in medicine, its benefits and risks.
 Do not by yourself interrupt the period of treatment prescribed for you.
 Do not repeat the same prescription without consulting your doctor.

Keep medicaments out of reach of children