

# MUCOFALK®-POWDER

---

## Composition:

1 bag with 5 g powder contains:

Plantago afra (Plantago psyllium) seed shells 3,25 g

Note for diabetics:

MUCOFALK contains sorbitol (0.07 carbohydrate exchanged units per bag)

## Indications:

For regulation in irritable colon and diverticulosis; for regulation of fecal passage in constipation and deficient peristalsis, e.g. owing to confinement to bed, after operations, in hemorrhoids and in pregnancy; in diarrhea and tendency to diarrhea without organic findings and for support of therapy in Crohn's disease; in chronic functional constipation.

## Contraindications:

In ileus and stenoses in the gastrointestinal tract, especially in the region of the esophagus, MUCOFALK should not be employed.

## Side effects:

Symptoms possibly present such as flatulence and feeling of fullness may occur to an increased extent in the first days of treatment, but these subside in the course of consistent further treatment.

## Interactions with other agents:

MUCOFALK should not be taken with motility - inhibiting antidiarrheics e.g. diphenoxylate, diphenoxamine, loperamide hydrochloride, opiates (danger of ileus).

## Dosage and instructions for use:

If not otherwise prescribed, adults take the content of one bag 1-3 times daily stirring into abundant fluid. It should best be taken before or during a meal (at the latest, one hour before going to bed).

1. Put the content of one bag of MUCOFALK into a glass.
2. Fill the glass slowly with cold water.
3. Stir it up with a spoon and drink it immediately.
4. It is advisable to drink one more glass of fluid. Per day 1.5 l fluid should be taken.

**Package size:** Package with 20 bags

**Keep all medicines out of the reach of children.**

Dry storage not above 25°C.

Manufactured by: Co. W. Weimer, D-7550 Rastatt

under authorisation of:



Dr. Falk Pharma GmbH  
D-7800 Freiburg  
Germany