# capozide

# Scored tablets

Composition

captopril . . . . . . . . . . . Excipients: lactose, microcrystalline cellulose, starch, sunset orange-yellow lacquer, magnesium

stearate, stearic acid.

Pharmaceutical form

Scored tablets: Box of 28 tablets

Pharmaco-therapeutic class

INHIBITOR OF THE ANGIOTENSINE CONVERSION ENZYME ASSOCIATED WITH A DIURETIC OF THE DILUTION CORTICAL SEGMENT (C: cardiovascular system).

# Company name and address

Bristol-Myers Squibb La Grande Arche Nord 92044 Paris la Défense cedex, France

# **INDICATIONS FOR USE**

This drug contains two antihypertensive agents and has both diuretic and vascular effects. It is indicated for the treatment of arterial hypertension.

# **WARNING!**

# This drug should NOT BE USED in the following conditions:

- allergy to this drug, including marked facial oedema.

pregnancy and breast-feeding.

IF IN DOUBT, DO NOT FAIL TO CONSULT YOUR DOCTOR OR PHARMACIST.

#### Special warnings

If a dry cough develops, it is essential to again consult your doctor who will decide whether or not to continue treatment.

There is a risk of facial oedema and allergic reactions.

Inform your doctor if you expect to undergo haemodialysis.

# Precautions for use

Inform your doctor in case of:

- renal or cardiac failure, diabetes, atherosclerosis, renal artery stenosis,
- low blood pressure or a strict "salt-free" diet.

In case of surgery, the anaesthetist should be informed that the patient is under treatment with Capozide®.

IF IN DOUBT, DO NOT FAIL TO CONSULT YOUR DOCTOR OR PHARMACIST.

### Drug interactions and other interactions

TO AVOID POSSIBLE INTERACTIONS WITH OTHER DRUGS YOUR DOCTOR OR PHARMA-CIST SHOULD SYSTEMATICALLY BE INFORMED OF ANY OTHER ONGOING TREATMENT, especially with lithium, potassium salts or a potassium-sparing diuretic, astemizole, bepridil, I.V. erythromycine, halofantrine, pentamidine, sultropide, terfenadine, and vincamine.

DO NOT LEAVE WITHIN THE REACH OF CHILDREN.

#### Pregnancy

Treatment must be suspended if pregnancy occurs while the patient is under treatment with a conversion enzyme inhibitor. The treating physician must be informed.

Warning for athletes: this drug contains an active ingredient which may yield positive results in antidoping control tests.

# **HOW TO USE THIS DRUG**

The dosage must be determined by a physician.

The usual dose is one tablet per day.

CAREFULLY FOLLOW YOUR DOCTOR'S PRESCRIPTION.

# Modality and route of administration

Oral route.

# Administration frequency and timing

This drug should be taken in a single dose in the morning.

## **Duration of treatment**

This drug must be taken regularly for as long as prescribed by your doctor, even if for a very long period of time.

#### What to do in case of overdose

The most likely effect of an overdose is a drop in blood pressure. In case of pronounced hypotension, have the patient lie down with legs raised.

If the condition continues, immediately inform your doctor.

# **UNDESIRED AND UNPLEASANT SIDE EFFECTS**

Skin rash, cough, dizziness, fatigue, and gastrointestinal disorders may develop. Inform your doctor. NOTIFY YOUR DOCTOR OR PHARMACIST OF ANY ADVERSE SIDE EFFECTS NOT MENTIONED IN THIS LEAFLET.

### **STORAGE**

Use before the date of expiry indicated on the box.

# Special storage precautions

This drug must be stored in a dry place away from direct heat.

#### **RELEASE CONDITIONS**

THIS DRUG IS REGISTERED IN LIST I.

DATE OF REVISION OF THIS LEAFLET: January 1996

# THIS IS A MEDICAMENT

- . Medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
- . Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.
- . The doctor and the pharmacist are experts in medicine, its benefits and risks.
- . Do not by yourself interrupt the period of treatment prescribed for you.
- . Do not repeat the same prescription without consulting your doctor.

# Keep Medicament out of reach of children

Council of Arab Health Ministers Union of Arab Pharmacists

Wereline to athless "to food contains an active of referrit