

1. NAME OF THE MEDICINAL PRODUCT

IVEMEND® 150 mg powder for solution for infusion.

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each vial contains fosaprepitant dimethylumine equivalent to 150 mg fosaprepitant, which corresponds to 130.5 mg of aprepitant. After reconstitution and dilution 1 ml of solution contains 1 mg fosaprepitant (1 mg/ml) (see section 6.6).

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Powder for solution for infusion.
White to off-white amorphous powder.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Prevention of acute and delayed nausea and vomiting associated with highly emetogenic cisplatin-based cancer chemotherapy in adults.

Prevention of nausea and vomiting associated with moderately emetogenic cancer chemotherapy in adults.

IVEMEND 150 mg is given as part of a combination therapy (see section 4.2).

4.2 Posology and method of administration

Posology

The recommended dose is 150 mg administered as an infusion **over 20-30 minutes** on Day 1, initiated approximately 30 minutes prior to chemotherapy (see section 6.6). IVEMEND should be administered in conjunction with a corticosteroid and a 5-HT3 antagonist as specified in the tables below.

The following regimens are recommended for the prevention of nausea and vomiting associated with emetogenic cancer chemotherapy.

Highly Emetogenic Chemotherapy Regimen

| | Day 1 | Day 2 | Day 3 | Day 4 |
|-------------------------------|--|-------------|--------------------------------|--------------------------------|
| IVEMEND | 150 mg intravenously | none | none | none |
| Dexamethasone | 12 mg orally | 8 mg orally | 8 mg orally twice daily | 8 mg orally twice daily |
| 5-HT ₃ antagonists | Standard dose of 5-HT ₃ antagonists. See the product information for the selected 5-HT ₃ antagonist for appropriate dosing information | none | none | none |

Dexamethasone should be administered 30 minutes prior to chemotherapy treatment on Day 1 and in the morning on Days 2 to 4. Dexamethasone should also be administered in the evenings on Days 3 and 4. The dose of dexamethasone accounts for active substance interactions.

Moderately Emetogenic Chemotherapy Regimen

| | Day 1 |
|-------------------------------|--|
| IVEMEND | 150 mg intravenously |
| Dexamethasone | 12 mg orally |
| 5-HT ₃ antagonists | Standard dose of 5-HT ₃ antagonists. See the product information for the selected 5-HT ₃ antagonist for appropriate dosing information |

Dexamethasone should be administered 30 minutes prior to chemotherapy treatment on Day 1. The dose of dexamethasone accounts for active substance interactions.

Efficacy data in combination with other corticosteroids and 5-HT₃ antagonists are limited. For additional information on the co-administration with corticosteroids, see section 4.5.

Refer to the Summary of Product Characteristics of co-administered 5-HT₃ antagonist medicinal products.

Special populations

Elderly (≥65 years)

No dose adjustment is necessary for the elderly (see section 5.2).

Gender

No dose adjustment is necessary based on gender (see section 5.2).

Renal impairment

No dose adjustment is necessary for patients with renal impairment or for patients with end stage renal disease undergoing haemodialysis (see section 5.2).

Hepatic impairment

No dose adjustment is necessary for patients with mild hepatic impairment. There are limited data in patients with moderate hepatic impairment and no data in patients with severe hepatic impairment. IVEMEND should be used with caution in these patients (see sections 4.4 and 5.2).

Paediatric population

The safety and efficacy of IVEMEND in children and adolescents below 18 years of age has not yet been established. Currently available data are described in sections 5.1 and 5.2, but no recommendation on a posology can be made.

Method of administration

IVEMEND 150 mg should be administered intravenously and should not be given by the intramuscular or subcutaneous route. Intravenous administration occurs preferably through a running intravenous infusion over 20-30 minutes (See section 6.6). Do not administer IVEMEND as a bolus injection or undiluted solution.

For instructions on reconstitution and dilution of the medicinal product before administration, see section 6.6.

4.3 Contraindications

Hypersensitivity to the active substance or to polysorbate 80 or any of the other excipients listed in section 6.1.

Co-administration with pimozide, terfenadine, astemizole or cisapride (see section 4.5).

4.4 Special warnings and precautions for use

Patients with moderate to severe hepatic impairment

There are limited data in patients with moderate hepatic impairment and no data in patients with severe hepatic impairment. IVEMEND should be used with caution in these patients (see section 5.2).

CYP3A4 interactions

IVEMEND should be used with caution in patients receiving concomitant active substances that are metabolised primarily through CYP3A4 and with a narrow therapeutic range, such as ciclosporin, tacrolimus, sirolimus, everolimus, alfentanil, ergot alkaloid derivatives, fentanyl, and quinidine (see section 4.5). Additionally, concomitant administration with irinotecan should be approached with particular caution as the combination might result in increased toxicity.

Co-administration with warfarin (a CYP2C9 substrate)

In patients on chronic warfarin therapy, the International Normalised Ratio (INR) should be monitored closely for 14 days following the use of fosaprepitant (see section 4.5).

Co-administration with hormonal contraceptives

The efficacy of hormonal contraceptives may be reduced during and for 28 days after administration of fosaprepitant. Alternative non-hormonal back-up methods of contraception should be used during treatment with fosaprepitant and for 2 months following the use of fosaprepitant (see section 4.5).

Hypersensitivity reactions

Immediate hypersensitivity reactions including flushing, erythema, dyspnoea, and anaphylaxis/anaphylactic shock have occurred during or soon after infusion of fosaprepitant. These hypersensitivity reactions have generally responded to discontinuation of the infusion and administration of appropriate therapy. It is not recommended to reinstitute the infusion in patients who experience hypersensitivity reactions.

Administration and infusion site reactions

Infusion site reactions (ISRs) have been reported with the use of IVEMEND (see section 4.8). The majority of severe ISRs, including thrombophlebitis and vasculitis, were reported with concomitant vesicant (e.g., anthracycline-based) chemotherapy administration, particularly when associated with extravasation. Necrosis was also reported in some patients with concomitant vesicant chemotherapy. Mild injection site thrombosis has been observed at higher doses without concomitant vesicant chemotherapy.

IVEMEND should not be given as a bolus injection, but should always be diluted and given as a slow intravenous infusion (see section 4.2). IVEMEND should not be administered intramuscularly or subcutaneously (see section 5.3). If signs or symptoms of local irritation occur, the injection or infusion should be terminated and restarted in another vein.

Sodium

This medicinal product contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially 'sodium-free'.

4.5 Interaction with other medicinal products and other forms of interaction

When administered intravenously fosaprepitant is rapidly converted to aprepitant. Interactions with other medicinal products following administration of intravenous fosaprepitant are likely to occur with active substances that interact with oral aprepitant. The following information was derived from studies conducted with oral aprepitant and studies conducted with intravenous fosaprepitant co-administered with dexamethasone, midazolam, or diltiazem.

Fosaprepitant 150 mg, given as a single dose, is a weak inhibitor of CYP3A4. Fosaprepitant does not seem to interact with the P-glycoprotein transporter, as demonstrated by the lack of interaction of oral aprepitant with digoxin. It is anticipated that fosaprepitant would cause less or no greater induction of CYP2C9, CYP3A4 and glucuronidation than that caused by the administration of oral aprepitant. Data are lacking regarding effects on CYP2C8 and CYP2C19.

Effect of fosaprepitant on the pharmacokinetics of other active substances

CYP3A4 inhibition

As a weak inhibitor of CYP3A4, the fosaprepitant 150 mg single dose can cause a transient increase in plasma concentrations of co-administered active substances that are metabolised through CYP3A4. The total exposure of CYP3A4 substrates may increase up to 2-fold on Days 1 and 2 after co- administration with a single 150 mg fosaprepitant dose. Fosaprepitant must not be used concurrently with pimozide, terfenadine, astemizole, or cisapride. Inhibition of CYP3A4 by fosaprepitant could result in elevated plasma concentrations of these active substances, potentially causing serious or life- threatening reactions (see section 4.3). Caution is advised during concomitant administration of fosaprepitant and active substances that are metabolised primarily through CYP3A4 and with a narrow therapeutic range, such as ciclosporin, tacrolimus, sirolimus, everolimus, alfentanil, diergotamine, ergotamine, fentanyl, and quinidine (see section 4.4).

Corticosteroids

Dexamethasone: The oral dexamethasone dose on Days 1 and 2 should be reduced by approximately 50 % when co-administered with fosaprepitant 150 mg on Day 1 to achieve exposures of dexamethasone similar to those obtained when given without fosaprepitant 150 mg. Fosaprepitant 150 mg administered as a single intravenous dose on Day 1 increased the AUC_{0-24hr} of dexamethasone, a CYP3A4 substrate, by 100 % on Day 1 86 % on Day 2 and 18 % on Day 3 when dexamethasone was co-administered as a single 8 mg oral dose on Days 1, 2, and 3.

Chemotherapeutic medicinal products

Interaction studies with fosaprepitant 150 mg and chemotherapeutic medicinal products have not been conducted; however, based on studies with oral aprepitant and docetaxel and vinorelbine, IVEMEND 150 mg is not expected to have a clinically relevant interaction with intravenously administered docetaxel and vinorelbine. An interaction with orally administered chemotherapeutic medicinal products metabolised primarily or partly by CYP3A4 (e.g. etoposide, vinorelbine) cannot be excluded. Caution is advised and additional monitoring may be appropriate in patients receiving medicinal products metabolized primarily or partly by CYP3A4 (see section 4.4). Post-marketing events of neurotoxicity, a potential adverse reaction of ifosfamide, have been reported after aprepitant and ifosfamide coadministration.

Immunosuppressants

Following a single 150 mg fosaprepitant dose, a transient moderate increase for two days possibly followed by a mild decrease in exposure of immunosuppressants metabolised by CYP3A4 (e.g., ciclosporin, tacrolimus, everolimus and sirolimus) is expected. Given the short duration of increased exposure, dose reduction of the immunosuppressant based on Therapeutic Dose Monitoring is not recommended on the day of and the day after administration of IVEMEND.

Midazolam

Fosaprepitant 150 mg administered as a single intravenous dose on Day 1 increased the AUC₀₋₆ of midazolam by 77 % on Day 1 and had no effect on Day 4 when midazolam was co-administered as a single oral dose of 2 mg on Days 1 and 4. Fosaprepitant 150 mg is a weak CYP3A4 inhibitor as a single dose on Day 1 with no evidence of inhibition or induction of CYP3A4 observed on Day 4.

The potential effects of increased plasma concentrations of midazolam or other benzodiazepines metabolised via CYP3A4 (alprazolam, triazolam) should be considered when co-administering these medicinal products with IVEMEND.

Diltiazem

Interaction studies with fosaprepitant 150 mg and diltiazem have not been conducted; however, the following study with 100 mg of fosaprepitant should be considered when using IVEMEND 150 mg with diltiazem. In patients with mild to moderate hypertension, infusion of 100 mg of fosaprepitant over 15 minutes with diltiazem 120 mg 3 times daily, resulted in a 1.4-fold increase in diltiazem AUC and a small but clinically meaningful decrease in blood pressure, but did not result in a clinically meaningful change in heart rate, or PR interval.

Induction

The fosaprepitant 150 mg single dose did not induce CYP3A4 on Days 1 and 4 in the midazolam interaction study. It is anticipated that IVEMEND would cause less or no greater induction of CYP2C9, CYP3A4, and glucuronidation than that caused by the administration of the 3-day oral aprepitant regimen, for which a transient induction with its maximum effect 6-8 days after first aprepitant dose has been observed. The 3-day oral aprepitant regimen resulted in an about 30-35 % reduction in AUC of CYP2C9 substrates and up to a 64 % decrease in ethinyl estradiol trough concentrations. Data are lacking regarding effects on CYP2C8 and CYP2C19. Caution is advised when warfarin, acenocoumarol, tolbutamide, phenytoin or other active substances that are known to be metabolised by CYP2C9 are administered with IVEMEND.

Warfarin

In patients on chronic warfarin therapy, the prothrombin time (INR) should be monitored closely during treatment with and for 14 days following the use of IVEMEND for the prevention of chemotherapy induced nausea and vomiting (see section 4.4).

Hormonal contraceptives

The efficacy of hormonal contraceptives may be reduced during and for 28 days after administration of fosaprepitant. Alternative non-hormonal back-up methods of contraception should be used during treatment with fosaprepitant and for 2 months following the use of fosaprepitant.

5-HT₃ antagonists

Interaction studies with fosaprepitant 150 mg and 5-HT₃ antagonists have not been conducted; however, in clinical interaction studies, the oral aprepitant regimen did not have clinically important effects on the pharmacokinetics of ondansetron, granisetron, or hydrodolasetron (the active metabolite of dolasetron). Therefore, there is no evidence of interaction with the use of IVEMEND 150 mg and 5-HT₃ antagonists.

Effect of other medicinal products on the pharmacokinetics of aprepitant resulting from administration of fosaprepitant 150 mg

Concomitant administration of fosaprepitant with active substances that inhibit CYP3A4 activity (e.g., ketoconazole, itraconazole, voriconazole, posaconazole, clarithromycin, telithromycin, nefazodone, and protease inhibitors) should be approached cautiously, as the combination is expected to result in several-fold increased plasma concentrations of aprepitant (see section 4.4). Ketoconazole increased the terminal half-life of oral aprepitant about 3-fold.

Concomitant administration of fosaprepitant with active substances that strongly induce CYP3A4 activity (e.g., rifampicin, phenytoin, carbamazepine, phenobarbital) should be avoided as the combination could result in reductions of the plasma concentrations of aprepitant that may result in decreased efficacy. Concomitant administration of fosaprepitant with herbal preparations containing St. John's Wort (*Hypericum perforatum*) is not recommended. Rifampicin decreased the mean terminal half-life of oral aprepitant by 68 %.

Diltiazem

Interaction studies with fosaprepitant 150 mg and diltiazem have not been conducted; however, the following study with 100 mg of fosaprepitant should be considered when using IVEMEND 150 mg with diltiazem. Infusion of 100 mg fosaprepitant over 15 minutes with diltiazem 120 mg 3 times daily, resulted in a 1.5-fold increase of aprepitant AUC. This effect was not considered clinically important.

4.6 Fertility, pregnancy and lactation

Contraception in males and females

The efficacy of hormonal contraceptives may be reduced during and for 28 days after administration of fosaprepitant. Alternative non-hormonal back-up methods of contraception should be used during treatment with fosaprepitant and for 2 months following the last dose of fosaprepitant (see sections 4.4 and 4.5).

Pregnancy

For fosaprepitant and aprepitant no clinical data on exposed pregnancies are available. The potential for reproductive toxicities of fosaprepitant and aprepitant have not been fully characterised, since exposure levels above the therapeutic exposure in humans could not be attained in animal studies. These studies did not indicate direct or indirect harmful effects with respect to pregnancy, embryonal/foetal development, parturition or postnatal development (see section 5.3). The potential effects on reproduction of alterations in neurokinin regulation are unknown. IVEMEND should not be used during pregnancy unless clearly necessary.

Breast-feeding

Aprepitant is excreted in the milk of lactating rats after intravenous administration of fosaprepitant as well as after oral administration of aprepitant. It is not known whether aprepitant is excreted in human milk. Therefore, breast-feeding is not recommended during treatment with IVEMEND.

Fertility

The potential for effects of fosaprepitant and aprepitant on fertility has not been fully characterised because exposure levels above the therapeutic exposure in humans could not be attained in animal studies. These fertility studies did not indicate direct or indirect harmful effects with respect to mating performance, fertility, embryonic/foetal development, or sperm count and motility (see section 5.3).

4.7 Effects on ability to drive and use machines

IVEMEND may have minor influence on the ability to drive and use machines. Dizziness and fatigue may occur following administration of IVEMEND (see section 4.8).

4.8 Undesirable effects

Summary of the safety profile

In clinical studies, various formulations of fosaprepitant have been administered to a total of 2,687 adults including 371 healthy subjects and 2,084 patients with chemotherapy induced nausea and vomiting (CINV). Since fosaprepitant is converted to aprepitant, those adverse reactions associated with aprepitant are expected to occur with fosaprepitant. The safety profile of aprepitant was evaluated in approximately 6,500 individuals.

Oral aprepitant

The most common adverse reactions reported at a greater incidence in adults treated with the aprepitant regimen than with standard therapy in patients receiving Highly Emetogenic Chemotherapy (HEC) were: hiccups (4.6 % versus 2.9 %), alanine aminotransferase (ALT) increased (2.8 % versus 1.1 %), dyspepsia (2.6 % versus 2.0 %), constipation (2.4 % versus 2.0 %), headache (2.0 % versus 1.8 %), and decreased appetite (2.0 % versus 0.5 %). The most common adverse reaction reported at a greater incidence in patients treated with the aprepitant regimen than with standard therapy in patients receiving Moderately Emetogenic Chemotherapy (MEC) was fatigue (1.4 % versus 0.9 %).

Tabulated list of adverse reactions - aprepitant

The following adverse reactions were observed in a pooled analysis of the HEC and MEC studies in adults at a greater incidence with oral aprepitant than with standard therapy or in postmarketing use:

Frequencies are defined as: very common (≥1/10); common (≥1/100 to <1/10); uncommon (≥1/1,000 to <1/100); rare (≥1/10,000 to <1/1,000) and very rare (<1/10,000), not known (cannot be estimated from the available data).

| System organ class | Adverse reaction | Frequency |
|--|---|-------------------------------|
| Infection and infestations | candidiasis, staphylococcal infection | rare |
| Blood and lymphatic system disorders | febrile neutropenia, anaemia | uncommon |
| Immune system disorders | hypersensitivity reactions including anaphylactic reactions | not known |
| Metabolism and nutrition disorders | decreased appetite polydipsia | common rare |
| Psychiatric disorders | anxiety disorientation, euphoric mood | uncommon rare |
| Nervous system disorders | headache dizziness, somnolence cognitive disorder, lethargy, dysgeusia | common uncommon rare |
| Eye disorders | conjunctivitis | rare |
| Ear and labyrinth disorders | tinnitus | rare |
| Cardiac disorders | palpitations bradycardia, cardiovascular disorder | uncommon rare |
| Vascular disorders | hot flush | uncommon |
| Respiratory, thoracic and mediastinal disorders | hiccups oropharyngeal pain, sneezing, cough, postnasal drip, throat irritation | common rare |
| Gastrointestinal disorders | constipation, dyspepsia eructation, nausea*, vomiting*, gastroesophageal reflux disease, abdominal pain, dry mouth, flatulence duodenal ulcer perforation, stomatitis, abdominal distension, faeces hard, neutropenic colitis | common uncommon rare |
| Skin and subcutaneous tissue disorders | rash, acne photosensitivity reaction, hyperhidrosis, seborrhoea, skin lesion, rash pruritic, Stevens-Johnson syndrome/toxic epidermal necrolysis pruritus, urticaria | uncommon rare not known |
| Musculoskeletal and connective tissue disorders | muscular weakness, muscle spasms | rare |
| Renal and urinary disorders | dysuria pollakisuria | uncommon rare |
| General disorders and administration site conditions | fatigue asthenia, malaise oedema, chest discomfort, gait disturbance | common uncommon rare |
| Investigations | ALT increased AST increased, blood alkaline phosphatase increased red blood cells urine positive, blood sodium decreased, weight decreased, neutrophil count decreased, glucose urine present, urine output increased | common uncommon rare |

*Nausea and vomiting were efficacy parameters in the first 5-days of post-chemotherapy treatment and were reported as adverse reactions only thereafter.

Description of selected adverse reactions

The adverse reactions profiles in the Multiple-Cycle extension of HEC and MEC studies in adults for up to 6 additional cycles of chemotherapy were generally similar to those observed in Cycle 1.

In an additional active-controlled clinical study in 1,169 adult patients receiving aprepitant and HEC, the adverse reactions profile was generally similar to that seen in the other HEC studies with aprepitant.

Additional adverse reactions were observed in adult patients treated with aprepitant for postoperative nausea and vomiting (PONV) and a greater incidence than with ondansetron: abdominal pain upper, bowel sounds abnormal, constipation*, dysarthria, dyspnoea, hypoaesthesia, insomnia, miosis, nausea, sensory disturbance, stomach discomfort, sub-ileus*, visual acuity reduced, wheezing.
*Reported in patients taking a higher dose of aprepitant.

Fosaprepitant

In an active-controlled clinical study in adult patients receiving HEC, safety was evaluated for 1,143 patients receiving the 1-day regimen of IVEMEND 150 mg compared to 1,169 patients receiving the 3-day regimen of aprepitant. Additionally, in a placebo-controlled clinical trial in adult patients receiving MEC, safety was evaluated for 504 patients receiving a single dose of IVEMEND 150 mg compared to 497 patients receiving the control regimen. The safety profile was generally similar to that seen in the aprepitant table above.
Tabulated list of adverse reactions – fosaprepitant
The following are adverse reactions reported in adult patients receiving fosaprepitant in clinical studies or postmarketing that have not been reported with aprepitant as described above: Infusion site reactions (ISRs) have been reported with the use of IVEMEND (see section 4.4).

Frequencies are defined as: very common (≥1/10); common (≥1/100 to <1/10); uncommon (≥1/1,000 to <1/100); rare (≥1/10,000 to <1/1,000) and very rare (<1/10,000), not known (cannot be estimated from the available data).

| System organ class | Adverse reaction | Frequency |
|--|--|-----------|
| Vascular disorders | flushing, thrombophlebitis (predominantly, infusion site thrombophlebitis) | uncommon |
| Skin and subcutaneous tissue disorders | erythema | uncommon |
| General disorders and administration site conditions | infusion site erythema, infusion site pain, infusion site pruritus | uncommon |
| | infusion site induration | rare |
| | immediate hypersensitivity reactions including flushing, erythema, dyspnoea, anaphylactic reactions/anaphylactic shock | not known |
| Investigations | blood pressure increased | uncommon |

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product.

4.9 Overdose

In the event of overdose, fosaprepitant should be discontinued and general supportive treatment and monitoring should be provided. Because of the antiemetic activity of aprepitant, emesis induced by a medicinal product may not be effective.

Aprepitant cannot be removed by haemodialysis.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antiemetics and antinauseants, ATC code: A04AD12.

Fosaprepitant is the prodrug of aprepitant and when administered intravenously is converted rapidly to aprepitant (see section 5.2). The contribution of fosaprepitant to the overall antiemetic effect has not fully been characterised, but a transient contribution during the initial phase cannot be ruled out. Aprepitant is a selective high-affinity antagonist at human substance P neurokinin 1 (NK₁) receptors. The pharmacological effect of fosaprepitant is attributed to aprepitant.

Highly Emetogenic Chemotherapy (HEC)

In a randomised, parallel, double-blind, active-controlled study, IVEMEND 150 mg (N=1,147) was compared with a 3-day aprepitant regimen (N=1,175) in adult patients receiving a HEC regimen that included cisplatin (≥ 70 mg/m²). The fosaprepitant regimen consisted of fosaprepitant 150 mg on Day 1 in combination with ondansetron 32 mg IV on Day 1 and dexamethasone 12 mg on Day 1, 8 mg on Day 2, and 8 mg twice daily on Days 3 and 4. The aprepitant regimen consisted of aprepitant 125 mg on Day 1 and 80 mg/day on Days 2 and 3 in combination with ondansetron 32 mg IV on Day 1 and dexamethasone 12 mg on Day 1 and 8 mg daily on Days 2 through 4. Fosaprepitant placebo, aprepitant placebo, and dexamethasone placebo (in the evenings on Days 3 and 4) were used to maintain blinding (see section 4.2). Although a 32 mg intravenous dose of ondansetron was used in clinical trials, this is no longer the recommended dose. See the product information for the selected 5-HT₃ antagonist for appropriate dosing information.

Efficacy was based on evaluation of the following composite measures: complete response in both the overall and delayed phases and no vomiting in the overall phase. IVEMEND 150 mg was shown to be non-inferior to that of the 3-day regimen of aprepitant. A summary of the primary and secondary endpoints is shown in Table 1.

| Table 1 Percent of adult patients receiving Highly Emetogenic Chemotherapy responding by treatment group and phase — Cycle 1 | | | |
|---|---|--|--------------------------------------|
| ENDPOINTS* | Fosaprepitant regimen (N =1,106) ** % | Aprepitant regimen (N =1,134) ** % | Difference [†] (95 % CI) |
| Complete response[‡] | | | |
| Overall [§] | 71.9 | 72.3 | -0.4 (-4.1, 3.3) |
| Delayed phase ^{§§} | 74.3 | 74.2 | 0.1 (-3.5, 3.7) |
| No vomiting | | | |
| Overall [¶] | 72.9 | 74.6 | -1.7 (-5.3, 2.0) |

*Primary endpoint is bolded.

**N: Number of patients included in the primary analysis of complete response.

†Difference and confidence interval (CI) were calculated using the method proposed by Miettinen and Nurminen and adjusted for gender.

‡Complete response = no vomiting and no use of rescue therapy.

§Overall = 0 to 120 hours post-initiation of cisplatin chemotherapy.

§§Delayed phase = 25 to 120 hours post-initiation of cisplatin chemotherapy.

Moderately Emetogenic Chemotherapy (MEC)

In a randomised, parallel, double-blind, placebo-controlled study, IVEMEND 150 mg (N=502) in combination with ondansetron and dexamethasone was compared with ondansetron and dexamethasone alone (control regimen) (N=498) in adult patients receiving a moderately emetogenic chemotherapy regimen. The fosaprepitant regimen consisted of fosaprepitant 150 mg on Day 1 in combination with oral ondansetron 8 mg for 2 doses and oral dexamethasone 12 mg. On Days 2 and 3, patients in the fosaprepitant group received placebo for ondansetron every 12 hours. The control regimen consisted of fosaprepitant placebo 150 mg IV on Day 1 in combination with oral ondansetron 8 mg for 2 doses and oral dexamethasone 20 mg. On Days 2 and 3, patients in the control group received 8 mg oral ondansetron every 12 hours. Fosaprepitant placebo and dexamethasone placebo (on Day 1) were used to maintain blinding.

The efficacy of fosaprepitant was evaluated based on the primary and secondary endpoints listed in Table 2 and was shown to be superior to the control regimen with regard to complete response in the delayed and overall phases.

| Table 2 Percent of adult patients receiving Moderately Emetogenic Chemotherapy responding by treatment group and phase | | | |
|---|---|-------------------------------------|---------|
| ENDPOINTS* | Fosaprepitant regimen (N =502) ** % | Control regimen (N =498) ** % | P-Value |
| Complete response[†] | | | |
| Delayed phase [‡] | 78.9 | 68.5 | <0.001 |
| Complete response [†] | | | |
| Overall [§] | 77.1 | 66.9 | <0.001 |
| Acute phase ^{§§} | 93.2 | 91 | 0.184 |

*Primary endpoint is bolded.

**N: Number of adult patients included in the intention to treat population.


†Complete response = no vomiting and no use of rescue therapy.

‡Delayed phase = 25 to 120 hours post-initiation of chemotherapy.

§Overall = 0 to 120 hours post-initiation of chemotherapy.

§§Acute = 0 to 24 hours post-initiation of chemotherapy.

The estimated time to first emesis is depicted by the Kaplan-Meier plot in Figure 1.



Paediatric population

The pharmacokinetics, safety and tolerability, and exploratory efficacy of intravenous fosaprepitant, administered concomitantly with ondansetron, with or without dexamethasone, were evaluated in a Phase I clinical study (N=34) in paediatric cancer patients receiving moderately or highly emetogenic chemotherapy. However, the efficacy and safety data from this small study do not support a conclusion on the optimal dosing regimen. Further studies evaluating the use of fosaprepitant in paediatric patients are on-going.

5.2 Pharmacokinetic properties

Fosaprepitant, a prodrug of aprepitant, when administered intravenously is rapidly converted to aprepitant. Plasma concentrations of fosaprepitant are below quantifiable levels within 30 minutes of the completion of infusion.

Aprepitant after fosaprepitant administration

Following a single intravenous 150-mg dose of fosaprepitant administered as a 20-minute infusion to healthy adult volunteers, the mean AUC_{0-∞} of aprepitant was 35.0 µg-hr/ml and the mean maximal aprepitant concentration was 4.01 µg/ml.

Distribution

Aprepitant is highly protein bound, with a mean of 97 %. The geometric mean volume of distribution at steady state (Vd_{ss}) of aprepitant estimated from a single 150 mg intravenous dose of fosaprepitant is approximately 82 l in humans.

Biotransformation

Fosaprepitant was rapidly converted to aprepitant in *in vitro* incubations with liver preparations from humans. Furthermore, fosaprepitant underwent rapid and nearly complete conversion to aprepitant in S9 preparations from other human tissues including kidney, lung and ileum. Thus, it appears that the conversion of fosaprepitant to aprepitant can occur in multiple tissues. In humans, fosaprepitant administered intravenously was rapidly converted to aprepitant within 30 minutes following the end of infusion.

Aprepitant undergoes extensive metabolism. In healthy young adults, aprepitant accounts for approximately 19 % of the radioactivity in plasma over 72 hours following a single intravenous administration 100 mg dose of [¹⁴C]- fosaprepitant, a prodrug for aprepitant, indicating a substantial presence of metabolites in the plasma. Twelve metabolites of aprepitant have been identified in human plasma. The metabolism of aprepitant occurs largely via oxidation at the morpholine ring and its side chains and the resultant metabolites were only weakly active. *In vitro* studies using human liver microsomes indicate that aprepitant is metabolised primarily by CYP3A4 and potentially with minor contribution by CYP1A2 and CYP2C19.

All metabolites observed in urine, faeces and plasma following an intravenous 100 mg [¹⁴C]- fosaprepitant dose were also observed following an oral dose of [¹⁴C]-aprepitant.

Upon conversion of

245.3 mg of fosaprepitant dimeglumine (equivalent to 150 mg fosaprepitant) to aprepitant, 23.9 mg of phosphoric acid and 95.3 mg of meglumine are liberated.

Elimination

Aprepitant is not excreted unchanged in urine. Metabolites are excreted in urine and via biliary excretion in faeces. Following a single intravenously administered 100 mg dose of [¹⁴C]- fosaprepitant to healthy subjects, 57 % of the radioactivity was recovered in urine and 45 % in faeces.

The pharmacokinetics of aprepitant is non-linear across the clinical dose range. The terminal half-life of aprepitant following a 150 mg intravenous dose of fosaprepitant was approximately 11 hours. The geometric mean plasma clearance of aprepitant following a 150 mg intravenous dose of fosaprepitant was approximately 73 ml/min.

Pharmacokinetics in special populations

Hepatic impairment: Fosaprepitant is metabolised in various extrahepatic tissues; therefore hepatic impairment is not expected to alter the conversion of fosaprepitant to aprepitant. Mild hepatic impairment (Child-Pugh class A) does not affect the pharmacokinetics of aprepitant to a clinically relevant extent. No dose adjustment is necessary for patients with mild hepatic impairment. Conclusions regarding the influence of moderate hepatic impairment (Child-Pugh class B) on aprepitant pharmacokinetics cannot be drawn from available data. There are no clinical or pharmacokinetic data in patients with severe hepatic impairment (Child-Pugh class C).

Renal impairment: A single 240 mg dose of oral aprepitant was administered to patients with severe renal impairment (CrCl< 30 ml/min) and to patients with end stage renal disease (ESRD) requiring haemodialysis.

In patients with severe renal impairment, the AUC_{0-∞} of total aprepitant (unbound and protein bound) decreased by 21 % and C_{max} decreased by 32 %, relative to healthy subjects. In patients with ESRD undergoing haemodialysis, the AUC_{0-∞} of total aprepitant decreased by 42 % and C_{max} decreased by 32 %. Due to modest decreases in protein binding of aprepitant in patients with renal disease, the

AUC of pharmacologically active unbound aprepitant was not significantly affected in patients with renal impairment compared with healthy subjects. Haemodialysis conducted 4 or 48 hours after dosing had no significant effect on the pharmacokinetics of aprepitant; less than 0.2 % of the dose was recovered in the dialysate.

No dose adjustment is necessary for patients with renal impairment or for patients with ESRD undergoing haemodialysis.

Paediatric population: Following administration of a single dose of 150 mg IV fosaprepitant to adolescent patients (aged 12 to 17 years), the mean aprepitant C_{max} and AUC_{0-∞} were approximately 5.9 µg/mL and 43.6 µg-hr/mL, respectively. Following administration of a single dose of 3 mg/kg IV fosaprepitant to paediatric patients aged 6 months to <12 years, the mean aprepitant C_{max} and AUC_{0-∞} were approximately 2.4 µg/mL and 20.8 µg-hr/mL, respectively.

Relationship between concentration and effect

Positron emission tomography (PET) imaging studies, using a highly specific NK₁-receptor tracer, in healthy young men administered a single intravenous dose of 150 mg fosaprepitant (N=8) demonstrated brain NK₁ receptor occupancy of ≥100 % at T_{max} and 24 hours, ≥97 % at 48 hours, and between 41 % and 75 % at 120 hours, following dosing. Occupancy of brain NK₁ receptors, in this study, correlate well with aprepitant plasma concentrations.

5.3 Pre-clinical safety data

Pre-clinical data obtained with intravenous administration of fosaprepitant and oral administration of aprepitant reveal no special hazard for humans based on conventional studies of single and repeated dose toxicity, genotoxicity (including *in vitro* tests), and toxicity to reproduction and development.

Carcinogenic potential in rodents was only investigated with orally administered aprepitant. However, it should be noted that the value of the toxicity studies carried out with rodents, rabbits and monkeys, including the reproduction toxicity studies, are limited since systemic exposures to fosaprepitant and aprepitant were only similar or even lower than therapeutic exposure in humans. In the performed safety pharmacology and repeated dose toxicity studies with dogs, fosaprepitant C_{max} and aprepitant AUC values were up to 3 times and 40 times, respectively, higher than clinical values.

In a toxicity study in juvenile dogs treated with fosaprepitant from postnatal day 14 to day 42, a decreased testicular weight and Leydig cell size were seen in the males at 6 mg/kg/day and increased uterine weight, hypertrophy of the uterus and cervix, and oedema of vaginal tissues were seen in females from 4 mg/kg/day. In a juvenile toxicity study in rats treated with aprepitant from postnatal day 10 to day 63, earlier vaginal opening in females from 250 mg/kg b.i.d. and delayed preputial separation in males from 10 mg/kg b.i.d was seen. There were no treatment-related effects on mating, fertility or embryonic/foetal survival, and no pathological changes in the reproductive organs. There were no margins to clinically relevant exposure of aprepitant. For short term treatment, these findings are considered unlikely to be clinically relevant.

In laboratory animals, fosaprepitant in non-commercial formulations caused vascular toxicity and haemolysis at concentrations below 1 mg/ml and higher, dependent on the formulation. In human washed blood cells also evidence of haemolysis was found with non-commercial formulations at fosaprepitant concentrations of 2.3 mg/ml and higher, although tests in human whole blood were negative. No haemolysis was found with the commercial formulation up to a fosaprepitant concentration of 1 mg/ml in human whole blood and washed human erythrocytes.

In rabbits, fosaprepitant caused initial transient local acute inflammation following paravenous, subcutaneous and intramuscular administration. At the end of the follow-up period (post-dose day 8), up to slight local subacute inflammation was noted following paravenous and intramuscular administration and additional up to moderate focal muscle degeneration/necrosis with muscle regeneration following intramuscular administration.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Disodium edetate (E386)
Polysorbate 80 (E433)
Lactose anhydrous
Sodium hydroxide (E524) (for pH adjustment) and/or
Hydrochloric acid diluted (E507) (for pH adjustment)

6.2 Incompatibilities

IVEMEND is incompatible with any solutions containing divalent cations (e.g., Ca²⁺, Mg²⁺), including Hartman's and lactated Ringer's solutions. This medicinal product must not be mixed with other medicinal products except those mentioned in section 6.6.

6.3 Shelf-life

2 years.

After reconstitution and dilution, chemical and physical in-use stability has been demonstrated for 24 hours at 25°C.

From a microbiological point of view, the medicinal product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2 to 8°C.

6.4 Special precautions for storage

Store in a refrigerator (2°C - 8°C).

For storage conditions after reconstitution and dilution of the medicinal product, see section 6.3.

6.5 Nature and contents of container

10 ml Type I clear glass vial with a chlorobutyl or bromobutyl rubber stopper and an aluminum seal with a grey plastic flip off cap.

Pack sizes: 1 or 10 vials.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

IVEMEND must be reconstituted and then diluted prior to administration.

Preparation of IVEMEND 150 mg for intravenous administration:

- Inject 5 ml sodium chloride 9 mg/ml (0.9 %) solution for injection into the vial. Assume that sodium chloride 9 mg/ml (0.9 %) solution for injection is added to the vial along the vial wall in order to prevent foaming. Swirl the vial gently. Avoid shaking and jetting sodium chloride 9 mg/ml (0.9 %) solution for injection into the vial.
- Prepare an infusion bag filled with **145 ml** of sodium chloride 9 mg/ml (0.9 %) solution for injection (for example, by removing 105 ml of sodium chloride 9 mg/ml (0.9 %) solution for injection from a 250 ml sodium chloride 9 mg/ml (0.9 %) solution for injection infusion bag).
- Withdraw the entire volume from the vial and transfer it into an infusion bag containing 145 ml of sodium chloride 9 mg/ml (0.9 %) solution for injection to **yield a total volume of 150 ml**. Gently invert the bag 2-3 times.

The medicinal product must not be reconstituted or mixed with solutions for which physical and chemical compatibility has not been established (see section 6.2).

The appearance of the reconstituted solution is the same as the appearance of the diluent.

The reconstituted and diluted medicinal product should be inspected visually for particulate matter and discoloration before administration.

No special requirements for disposal.

7. MARKETING AUTHORISATION HOLDER

Merck Sharp & Dohme Ltd.
Hertford Road, Hoddesdon
Hertfordshire EN 11 9BU
United Kingdom

8. Manufactured by:

Patheon Manufacturing Services LLC ,
5900 Martin Luther King Jr. Highway, Greenville, NC 27834,
USA

9. Released by:

Merck Sharp & Dohme B.V.
Waarderweg 39
2031 BN Haarlem
The Netherlands

10. DATE OF REVISION OF THE TEXT May 2020

Detailed information on this medicinal product is available on the website of the European Medicines Agency http://www.ema.europa.eu.

(THIS IS A MEDICAMENT)
-Medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
-Follow strictly the doctor's prescription, the method of use, and the instructions of the pharmacist who sold the medicament.
-The doctor and the pharmacist are experts in medicine, its benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed for you.
-Do not repeat the same prescription without consulting your doctor.
-Keep medicament out of reach of children
Council of Arab Health Ministers and Union of Arab Pharmacists

