TRIUMEO

Dolutegravir-abacavir-lamivudine

QUALITATIVE AND QUANTITATIVE COMPOSITIO

Purple biconvex avail tablets debassed with "572 Tri" on one side

Each film-coated tablet contains 50 mg of dolutegravir as dolutegravir sodium, 600 mg of abacavir as abacavir sulfate and 300 mg of lamivudine. PHARMACEUTICAL FORM

Film-coated tablets CLINICAL PARTICULARS

TRIUMEO is indicated for the treatment of Human Immunodeficiency Virus (HIV) infection in adults and adolescents from 12 years of age who are antiretroviral treatment-naïve or are infected with HIV without documented or clinically suspected resistance to any of the three antiretroviral agents in TRIUMEO.

Dosage and Administration TRIUMEO therapy should be initiated by a physician experienced in the management of HIV infection.

TRIUMEO should not be administered to adults or adolescents who weigh less than 40 kg because it is a fixed-dose tablet that cannot be dose reduced. TRIUMEO can be taken with or without food.

TRIUMEO is a fixed-dose tablet and should not be prescribed for patients requiring dosage adjustments, such as those with creatinine clearance less than 50 ml/min. Separate preparations of TIVICAY, ZIAGEN or EPIVIR should be administered in cases where discontinuation or dose adjustment is indicated. In these cases the physician should refer to the individual product information for these medicinal products.

Since the recommended dose of dolutegravir is 50 mg twice daily for patients with resistance to integrase inhibitors, the use of TRIUMEO is not

Adults and adolescents The recommended dose of TRIUMEQ in adults and adolescents weighing at least 40 kg is one tablet once daily.

TRIUMEQ is not currently recommended for treatment of children less than 12 years of age as the necessary dose adjustment cannot be made. Clinical

data is currently not available for this combination. Physicians should refer to the individual product information for TIVICAY, ZIAGEN and EPIVIR. Elderly There are limited data available on the use of TIVICAY, ZIAGEN and EPIVIR in patients aged 65 years and over. However, there is no evidence that elderly patients require a different dose than younger adult patients (see Pharmacokinetics — Special Patient Populations). When treating elderly

patients, consideration needs to be given to the greater frequency of decreased hepatic, renal and cardiac function, concomitant medicinal Renal impairment

Whilst no dosage adjustment of dolutegravir or abacavir is necessary in patients with renal impairment, a dose reduction of EPIVIR is required due to decreased clearance. Therefore TRIUMEQ is not recommended for use in patients with a creatinine clearance less than 50 ml/min (see Pharmacokinetics — Special Patient Populations). Hepatic impairment

A dose reduction of ZIAGEN may be required for patients with mild hepatic impairment (Child-Pugh grade A). As dose reduction is not possible with RIUMEQ, the separate preparations of TIVICAY, ZIAGEN and EPIVIR should be used when this is judged necessary. TRIUMEQ is not recommended in patients with moderate and severe hepatic impairment (Child-Pugh grade B or C) (see Pharmacokinetics — Special Patient Populations). Contraindications

TRIUMEO is contraindicated in patients with known hypersensitivity to dolutegravir, abacavir or lamivudine, or to any of the excipients. TRIUMEO is contraindicated in combination with dofetilide or nilsicainide

The special warnings and precautions relevant to TIVICAY. ZIAGEN and EPIVIR are included in this section. There are no additional precautions and

Warnings and Precautions

Hypersensitivity reactions (see also Adverse Reactions): Both abacavir and dolutegravir are associated with a risk for hypersensitivity reactions (HSR) (see Clinical Description of HSR below, and Adverse Reactions), and share some common features such as fever and/or rash with other symptoms indicating multi-organ involvement

peen observed more commonly with abacavir, some of which have been life-threatening, and in rare cases fatal. The risk for abacavir HSR to occur is significantly increased for patients who test positive for the HLA-B*5701 allele. However, abacavir HSRs have been reported in patients who do not carry this allele.

The following should be adhered to:

 Testing for HI A-R*5701 status should be considered before initiating abacavir treatment and also before re-starting abacavir treatment in patients of unknown *HLA-B*5701* status who have previously tolerated abacavir.

Clinically it is not possible to determine whether a HSR with *TRIUMEQ* is caused by abacavir or dolutegravir. Hypersensitivity reactions have

- TRIUMEQ is not recommended for use in patients with the HLA-B*5701 allele, or in patients who have had a suspected abacave. HSR while taking any other medicinal product containing abacavir (e.g. ZIAGEN. KIVEXA, TRIZIVIR) regardless of HLA-B*5701 status. Each patient should be reminded to read the Patient Leaflet included in the TRIUMEO pack. They should be reminded of the
- importance of removing the Alert Card included in the pack, and keeping it with them at all times. • In any patient treated with TRIUMEO, the clinical diagnosis of suspected hypersensitivity reaction must remain the basis of clinical decision making.
- TRIUMEQ must be stopped without delay, even in the absence of the HLA-B*5701 allele, if a HSR is suspected. Delay in stopping treatment with TRIUMEQ after the onset of hypersensitivity may result in a life-threatening **reaction**. Clinical status including liver aminotransferases and bilirubin should be monitored.
- Patients who have experienced a hypersensitivity reaction should be instructed to dispose of their remaining TRIUMEQ tablets,
- After stopping treatment with TRIUMEQ for reasons of a suspected HSR, TRIUMEQ or any other medicinal product containing abacavir or dolutegravir must never be re-initiated.
- Restarting abacavir containing products following a suspected abacavir HSR can result in a prompt return of symptoms within hours and may include life-threatening hypotension and death. If a hypersensitivity reaction is ruled out, patients may restart TRIUMEQ. Rarely, patients who have stopped abacavir for
- reasons other than symptoms of HSR have also experienced life-threatening reactions within hours of re-initiating abacavir therapy (*see Adverse Reactions, Description of selected adverse reactions*). Patients must be made aware that HSR can occu with reintroduction of TRIUMEQ or any other medicinal product containing abacavir (e.g. ZIAGEN, KIVEXA, TRIZIVIR) and that reintroduction of TRIUMEQ or any other medicinal product containing abacavir (e.g. ZIAGEN, KIVEXA, TRIZIVIR) should be undertaken only if medical care can be readily accessed. Clinical Description of HSR with dolutegravir:

lypersensitivity reactions have been reported with integrase inhibitors, including dolutegravir, and were characterized by rash, constitutional findings, and sometimes, organ dysfunction, including liver injury Clinical Description of HSR with abacavir:

Abacavir HSR has been well characterised through clinical studies and during post marketing follow-up. Symptoms usually appeared within the first six weeks (median time to onset 11 days) of initiation of treatment with abacavir, although these reactions may occur at any

Other signs and symptoms that have been observed as part of abacavir HSR include respiratory and gastrointestinal symptoms, which may lead to misdiagnosis of HSR as respiratory disease (pneumonia, bronchitis, pharyngitis), or gastroenteritis (see Adverse Reactions Description of Selected Adverse Reactions). The symptoms related to HSR worsen with continued therapy and can be life

threatening. These symptoms usually resolve upon discontinuation of abacavir.

Almost all HSR to abacavir include fever and/or rash as part of the syndrome.

actic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported with the use of antiretroviral nucleoside nalogues either alone or in combination, including abacavir and lamivudine. A maiority of these cases have been in women, Clinical features which may be indicative of the development of lactic acidosis include generalised weakness, anorexia, and sudden unexplained

weight loss, gastrointestinal symptoms and respiratory symptoms (dyspnoea and tachypnoea). ution should be exercised when administering TRIUMEQ particularly to those with known risk factors for liver disease. Treatment with TRIUMEQ should be suspended in any natient who develops clinical or laboratory findings suggestive of lactic acidosis with or without benatitis (which

Serum lipid and blood glucose levels may increase during antiretroviral therapy. Disease control and life style changes may also be contributing factors. Consideration should be given to the measurements of serum lipids and blood glucose. Lipid disorders should be managed as clinically

In HIV-infected patients with severe immune deficiency at the time of initiation of anti-retroviral therapy (ART), an inflammatory reaction to

actions have been observed within the first few weeks or months of initiation of ART. Relevant examples are cytomegalovirus retinitis generalised and/or focal mycobacterial infections and *Pneumocystis jiroyeci* pneumonia (often referred to as PCP). Any inflammatory symptoms st be evaluated without delay and treatment initiated when necessary. Autoimmune disorders (such as Graves' disease, polymyositis and Guillain-Barre syndrome) have also been reported to occur in the setting of immune reconstitution, however, the time to onset is more variable. and can occur many months after initiation of treatment and sometimes can be an atypical presentatio iver chemistry elevations consistent with immune reconstitution syndrome were observed in some hepatitis B and/or C co-infected patients at

the start of dolutegravir therapy. Monitoring of liver chemistries is recommended in patients with hepatitis B and/or C co-infection (see Patients infected with hepatitis B virus (HBV) later in this section and Adverse Reactions). Patients co-infected with henatitis R virus (HRV)

Particular diligence should be applied in initiating or maintaining effective hepatitis B therapy when starting therapy with TRIUMEQ in hepatitis B co-infected patients. Clinical study and marketed use of lamivudine, have shown that some patients with chronic HBV disease may experience clinical or laboratory idence of recurrent hepatitis upon discontinuation of lamivudine, which may have more severe consequences in patients with decompensated liver disease. If TRIUMEQ is discontinued in patients co-infected with HBV, periodic monitoring of both liver function tests and markers of HBV

eplication should be considered. Opportunistic infections: Patients receiving TRIUMEQ or any other antiretroviral therapy may still develop opportunistic infections and other complications of HIV infection.

Therefore patients should remain under close clinical observation by physicians experienced in the treatment of these associated HIV diseases.

Patients should be advised that current antiretroviral therapy, including TRIUMEQ has not been proven to prevent the risk of transmission of HIV to others through sexual contact or blood contamination. Appropriate precautions should continue to be taken. Myocardial Infarction: Several observational, epidemiological studies have reported an association with abacavir use and risk of myocardial infarction. Meta-analyses of

randomised controlled trials have observed no excess risk of myocardial infarction with abacavir use. To date, there is no established biological nechanism to explain a notential increase in risk. In totality the available data from observational studies and from controlled clinical trials show inconsistency and therefore the evidence for a causal relationship between abacavir treatment and the risk of myocardial infarction is inconclusive As a precaution the underlying risk of coronary heart disease should be considered when prescribing antiretroviral therapies, including abacavir. and action taken to minimize all modifiable risk factors (e.g. hypertension, hyperlipidaemia, diabetes mellitus and smoking).

Caution should be given to co-administering medications (prescription and non-prescription) that may change the exposure of dolutegravir. abacavir, lamivudine or medications that may have their exposure changed by TRIUMEQ (see Contraindications and Interac TRIUMEQ should not be administered concurrently with other medicinal products containing any of the same active components (dolutegravir abacavir, and/or lamivudine).

Since the recommended dose of TIVICAY is 50 mg twice daily for patients taking etravirine (without boosted protease inhibitors), efavirenz, evirapine, rifampicin, tipranavir/ritonavir, carbamazepine, phenytoin, phenobarbital and St. John's wort, the use of TR/UMEQ is not recommended for patients taking these medicines (see Interactions).

Concomitant Drug Class: | Effect on Concentration | Clinical Comment

Concomitant Drug

Cmax ↓3%

Pilsicainide ↑

Tenofovir did not change dolutegravir plasma concentration to a clinically

Co-administration of dolutegravir has the notential to increase dofetilide of

pilsicainide plasma concentration via inhibition of OCT2 transporter;

co-administration has not been studied. Dofetilide or pilsicainide

life-threatening toxicity caused by high dofetilide or pilsicainide

Carbamazepine decreased dolutegravir plasma concentration. Since the

recommended dose of dolutegravir is 50 mg twice daily when co-administered with carbamazepine, TRIUMEQ is not recommended for

Co-administration with these metabolic inducers may decrease dolutegra

plasma concentration and has not been studied. Co-administration with

these metabolic inducers should be avoided. Co-administration with these

concentration due to enzyme induction and has not been studied. The effect

metabolic inducers has the potential to decrease dolutegravir plasma

of these metabolic inducers on dolutegravir exposure is likely similar to

commended for patients taking these metabolic inducers.

not expected. No dose adjustment is necessary.

containing polyvalent cations.

carbamazepine. Since the recommended dose of dolutegravir is 50 mg twice

daily when co-administered with these metabolic inducers. TRIUMEQ is not

This interaction has not been studied. Although an inducer of CYP3A4, based

on data from other inducers, a clinically significant decrease in dolutegravir is

inistration of antacids containing polyvalent cations decrease

dolutegravir plasma concentration. Dolutegravir is recommended to be

TRIUMEO is recommended to be administered 2 hours before or 6 hours after

taking products containing calcium, or alternatively, administer with food.

TRIUMEO is recommended to be administered 2 hours before or 6 hours after

taking products containing iron, or alternatively, administer with food.

Rifampicin decreased dolutegravir plasma concentration. Since the dose of

co-administration of rifampicin with TRIUMEQ is not recommended.

concentrations to a clinically relevant extent. No dose adjustment of oral

ntraceptives is necessary when co-administered with dolutegravi

Dolutegravir did not change methadone plasma concentrations to a clinically

relevant extent. No dose adjustment of methadone is necessary when

co-administered with dolutegravir.

dolutegravir is 50 mg twice daily when co-administered with rifamnicin the

Co-administration of dolutegravir increased metformin plasma

When co-administered with | concentration. A dose adjustment of metformin should be considered wh

dolutegravir 50mg 0D: starting and stopping coadministration of dolutegravir with metformin, to

administered 2 hours before or 6 hours after taking antacid products

natients taking carbamazepine.

relevant extent. No dose adjustment is necessary.

Nucleoside Reverse

Tenofovir (TDF)

Other Agents

St. John's wort

polyvalent cations (e.g., Mg, │ AUC ↓ 74%

When co-administered wit

dolutegravir 50mg BID

Effect of dolutegray

Effect of dolutegravir:

AUC ↓ 2%

Dolutegravir should not be co-administered with polyvalent cation-containing antacids. TRIUMEQ is recommended to be administered 2 hours refore or 6 hours after these agents (see Interactions) TRIUMEQ is recommended to be administered 2 hours before or 6 hours after taking calcium or iron supplements, or alternatively, administered

Dolutegravir increased metformin concentrations. A dose adjustment of metformin should be considered when starting and stopping oadministration of dolutegravir with metformin, to maintain glycaemic control (see Interactions)

As TRIUMFO contains dolutedravir. abacavir and lamivudine, any interactions that have been identified with these agents individually may occur with TRIUMEQ. Due to the different routes of metabolism and elimination, no clinically significant drug interactions are expected between dolutegravir, abacavir and lamivudine. In a cross study comparison, abacavir and lamivudine exposures were similar when given as TRIUMEQ

compared to KIVEXA alone. Effect of Dolutegravir, Abacavir and Lamivudine on the Pharmacokinetics of Other Agents 1 vitro, dolutegravir demonstrated no direct, or weak inhibition (IC50>50 μM) of the enzymes cytochrome P450 (CYP)1A2, CYP2A6, CYP2B6,

with food (see Interactions).

CYP2C8, CYP2C9, CYP2C19, CYP2D6 CYP3A, uridine diphosphate glucuronosyl transferase (UGT)1A1 or UGT2B7, or the transporters P-glycon (Pgp), breast cancer resistance protein (BCRP), bile salt export pump (BSEP), organic anion transporting polypeptide 1B1 (OATP1B1), OATP1B3, organic anion transporter 1(OCT1), multidrug resistance associated protein 2 (MRP2), or MRP4, *In vitro*, dolutegravir did not induce CYP1A2. CYP2B6 or CYP3A4. In vivo, dolutegravir did not have an effect on midazolam, a CYP3A4 probe. Based on these data, dolutegravir is not expected to affect the pharmacokinetics of drugs that are substrates of these enzymes or transporters.

In drug interaction studies, dolutegravir did not have a clinically relevant effect on the pharmacokinetics of the following: tenofovir, ritonavir, methadone, efavirenz, lopinavir, atazanavir, darunavir, etravirine, fosamprenavir, rilpivirine, boceprevir, telaprevir, daclatasvir and oral contraceptives containing norgestimate and ethinyl estradiol.

In vitro, dolutegravir inhibited the renal organic cation transporter 2 (OCT2) (ICSO = $1.93 \, \mu M$), multidrug and toxin extrusion transporter (MATE) $(ICSO = 6.34 \,\mu\text{M})$ and MATE2-K ($ICSO = 24.8 \,\mu\text{M})$. Given dolutegravir's *in vivo* exposure, it has a low potential to affect the transport of MATE2-K substrates in vivo. In vivo, dolutegravir may increase plasma concentrations of drugs in which excretion is dependent upon OCT2 or ATE1 (dofetilide, pilsicainide or metformin) (see Table 1).

In vitro, dolutegravir inhibited the basolateral renal transporters: organic anion transporter (OAT) 1 (IC50 = $2.12 \,\mu\text{M}$) and OAT3 (IC50 = $1.97 \,\mu\text{M}$). However, dolutegravir had no notable effect on the pharmacokinetics in vivo of the OAT substrates tenofovir and para aminohippurate, and efore has low propensity to cause drug interactions *via* inhibition of OAT transporter

pacavir and lamivudine do not inhibit or induce CYP enzymes (such as CYP 3A4, CYP 2C9 or CYP 2D6) and demonstrate no or weak inihibition of the OATP1B3, BCRP and Pgp or MATE2-K. In addition, lamivudine demonstrates no or weak inhibition of the drug transporters MATE1or OCT3 and imal inhibition of OCT1 and OCT2. Abacavir and lamivudine are therefore not expected to affect the plasma

oncentrations of drugs that are substrates of these enzymes or transporters. Although abacavir is an inhibitor of MATE1 and lamivudine is an inhibitor of OCT1 and OCT2 in vitro, they have low potential to affect the plasma ntrations of substrates of these transporters at therapeutic drug exposures (up to 600 mg for abacavir or 300 mg for lamivudine).

Effect of Other Agents on the Pharmacokinetics of Dolutegravir, Abacavir and Lamivudine egravir is eliminated mainly through metabolism by UGT1A1. Dolutegravir is also a substrate of UGT1A3, UGT1A9, CYP3A4, Pgp, and BCRP; therefore drugs that induce these enzymes or transporters may theoretically decrease dolutegravir plasma concentration and reduce the therapeutic effect of TRIUMEQ. Co-administration of dolutegravir and other drugs that inhibit UGT1A1, UGT1A3, UGT1A9, CYP3A4, and/or Pgp may increase dolutegravir plasma concentration (see Table 1).

In vitro, dolutegravir is not a substrate of human organic anion transporting polypeptide (OATP)1B1, OATP1B3, or OCT1, therefore drugs that lely modulate these transporter are not expected to affect dolutegravir plasma concentration Efavirenz, etravirine, nevirapine, rifampicin, carbamazepine and tipranavir in combination with ritonavir each reduced the plasma concentration

of dolutegravir significantly, and require dolutegravir dose adjustment to 50 mg twice daily. The effect of etravirine was mitigated by ninistration of the CYP3A4 inhibitors lopinavir/ritonavir, darunavir/ritonavir, and is expected to be mitigated by atazanavir/ritonavir. herefore no dolutegravir dose adjustment is necessary when co-administered with etravirine and either lopinavir/ritonavir, darunavir/ritonavir, or atazanavir/ritonavir. Another inducer, fosamprenavir in combination with ritonavir decreased plasma concentrations of dolutegravir but does not require a dosage adjustment of dolutegravir. A drug interaction study with the UGT1A1 inhibitor, atazanavir, did not result in a clinically eaningful increase in the plasma concentrations of dolutegravir. Tenofovir, lopinavir/ritonavir, darunavir/ritonavir, rilpivirine, boceprevi telaprevir, prednisone, rifabutin, daclatasvir and omeprazole had no or a minimal effect on dolutegravir pharmacokinetics, therefore no dolutegravir dose adjustment is required when co-administered with these drugs. The likelihood of metabolic interactions with abacavir and lamivudine is low. Abacavir and lamivudine are not significantly metabolised by CYP

enzymes. The primary pathways of abacavir metabolism in human are by alcohol dehydrogenase and by glucuronidation to produce the 5'-carboxylio cid and 5'-glucuronide which account for about 66% of the administered dose. These metabolites are excreted in the urine. The likelihood of metabolic interactions with lamivudine is low due to limited metabolism and plasma protein binding, and almost complete renal clearance. *In vitro*, bacavir is not a substrate of OATP1B1, OATP1B3, OCT1, OCT2, OAT1, MATE1, MATE2-K, MRP2 or MRP4 therefore drugs that modulate these transporters are not expected to affect abacavir plasma concentrations. Although abacavir and lamivudine are substrates of BCRP and Pop *in vitro*. ical studies demonstrate no clinically significant changes in abacavir pharmacokinetics when co-administered with lopinavir/ritonavir (Pgp and BCRP inhibitors) and inhibitors of these efflux transporters are unlikely to affect the disposition of lamivudine due to its high bioavailability. mivudine is an in vitro substrate of MATE1. MATE2-K and OCT2. Trimethoprim (an inhibitor of these drug transporters) has been shown to increase lamivudine plasma concentrations however; the resulting increase was of such magnitude that a dose adjustment is not recommended as it is not pected to have clinical significance. Larnivudine is a substrate of the hepatic uptake transporter OCT1. As hepatic elimination plays a minor role in the clearance of lamivudine, drug interactions due to inhibition of OCT1 are unlikely to be of clinical significance. TRIUMEO is a fixed-dose tablet and should not be prescribed for patients requiring dosage adjustments due to interacting concomitant

nedications. Separate preparations of TIVICAY, ZIAGEN or EPIVIR should be administered in cases where dose adjustment is required. In these cases the physician should refer to the individual product information for these medicinal products. Selected drug interactions are presented in Tables 1, 2 and 3. Recommendations are based on either drug interaction studies or predicted nteractions due to the expected magnitude of interaction and potential for serious adverse events or loss of efficacy

Table 1 Drug Interactions studied with dolutegravir Concomitant Drug Class: | Effect on Concentration | Clinical Comment

Non-nucleoside Reverse Dolutegravir

of Dolutegravir or

AUC ↑ 62% C_{max} ↑ 34%

AUC ↓ 59%

AUC ↓ 35%

Dolutegravir ←

AUC ↓ 4%

AUC ↓ 22%

Protease Inhibitor

Lopinavir/ritonavir

Drug Name

| Transcriptase Inhibitor: | AUC ↓ 71% | concentration. Since the recommended dose of dolutegravir is 50 mg twice daily | | | cπ ↓ 1% | | |
|-----------------------------------------------------|----------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------|---------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Etravirine (ETR) without boosted protase inhibitors | $C_{max} \downarrow 52\%$ $C\tau \downarrow 88\%$ ETR \leftrightarrow | for patients taking etravirine without boosted protease inhibitors, TRIUMEQ is not recommended for patients taking etravirine without co-administration of atazanavir/ritonavir, darunavir/ritonavir or lopinavir/ritonavir. | Da | | AUC ↑ 33% C _{max} ↑ 29% | relevar | Daclatasvir did not change dolutegravir plasma concentration to a clinically relevant extent. Dolutegravir did not change daclatasvir plasma concentration. No dose adjustment is necessary. |
| Protease Inhibitor: Lopinavir/ritonavir + | Dolutegravir ↔ AUC ↑ 11% | Lopinavir/ritonavir and etravirine did not change dolutegravir plasma concentration to a clinically relevant extent. No dose adjustment is necessary. | | | Cτ ↑ 45% Daclatasvir ↔ | | |
| Etravirine (LPV/RTV+ETR) | $C_{max} \uparrow 7\%$ $C\tau \uparrow 28\%$ $LPV \leftrightarrow$ $RTV \leftrightarrow$ | , | | observed concentration, Cτ=con | | nterval | ; AUC = area under the concentration versus time curve; C_{max} = maximum |
| Protease Inhibitor: Darunavir/ritonavir + | Dolutegravir↓ AUC↓25% | Darunavir/ritonavir and etravirine did not change dolutegravir plasma concentration to a clinically relevant extent. No dose adjustment is necessary. | | Concomitant Drug Class: Drug Name | Effect on Concentration of abacavir or Concomitant D | | Clinical Comment |
| Etravirine (DRV/RTV+ETR) | $C_{\text{max}} \downarrow 12\%$ $C\tau \downarrow 36\%$ DRV \leftrightarrow RTV \leftrightarrow | Circulation of Cinical Processing Street Survey Street Survey Sur | | Methadone (40 to 90 mg once daily for 14 days/600 mg single dose, then 600 mg twice daily for 14 days) | Abacavir AUC ←→ C _{max} ↓35% Methadone CL/F ↑22% | | The changes in abacavir pharmacokinetics are not considered clinically relevant. The changes in methadone pharmacokinetics are not considered clinically relevant for the majority of patients, however occasionally methadone dose re-titration may be required. |
| Non-nucleoside Reverse Transcriptase Inhibitor: | Dolutegravir↓ AUC↓57% | Efavirenz decreased dolutegravir plasma concentrations. Since the dose of dolutegravir is 50 mg twice daily when co-administered with efavirenz the | | Ethanol | Abacavir AUC ↑41% Ethanol AUC ↔ | | Given the safety profile of abacavir, these findings are not considered clinically significant. |
| Efavirenz (EFV) | $\begin{array}{c} C_{\text{max}} \downarrow 39\% \\ C\tau \downarrow 75\% \\ \text{EFV} \leftrightarrow \end{array}$ | co-administration of efavirenz with <i>TRIUMEQ</i> is not recommended. | | Abbreviations: \uparrow = Increase; \downarrow = decrease; \leftrightarrow = no significant change; AUC = area under the concentration versus time curve; C_{max} = maximum observed concentration, CL/F = apparent clearance Table 3 Drug Interactions studied with lamivudine | | | |
| Non-nucleoside Reverse Transcriptase Inhibitor: | Dolutegravir↓ | Co-administration with nevirapine has the potential to decrease dolutegravir plasma concentration due to enzyme induction and has not been studied. | | Concomitant Drug Class: | Effect on Concentration | | Clinical Comment |

Effect of nevirapine on dolutegravir exposure is likely similar to or less than

co-administered with nevirapine, the co-administration of nevirapine with

Atazanavir increased dolutegravir plasma concentration. No dose adjustmen

Atazanavir/ritonavir increased dolutegravir plasma concentration. No dose

Tipranavir/ritonavir decreases dolutegravir concentrations. Since the dose of

Fosamprenavir/ritonavir decreases dolutegravir concentrations, but based or

limited data, did not result in decreased efficacy in Phase III studies. No dose

This interaction has not been studied. Although an inhibitor of CYP3A4, based

on data from other inhibitors, an increase is not expected. No dose

Darunavir/ritonavir did not change dolutegravir plasma concentration to

clinically relevant extent. No dose adjustment is necessary.

clinically relevant extent. No dose adjustment is necessary.

tipranavir/ritonavir, the co-administration of tipranavir/ritonavir with

dolutegravir is 50 mg twice daily when co-administered with

RIUMEQ is not recommended.

adjustment is necessary.

adjustment is necessary in INI-naïve patients.

that of efavirenz. Since the dose of dolutegravir is 50 mg twice daily when

| imethopmm/sulfamethoxazole io-trimoxazole) (160 mg/800 mg nce daily for 5 days/300 mg ngle dose) | Lamrudine: AUC 40% Trimethoprim: AUC ↔ Sulfamethoxazole: AUC ↔ | Unless the patient has renal impairment, no dosage adjustment or lamivudine is necessary (see Dosage and Administration). Lamivudine has no effect on the pharmacokinetics of trimethoprim or sulfamethoxazole. The effect of co-administration of lamivudine with higher doses of co-trimoxazole used for the treatment of Pneumocystis jiroveci pneumoia and toxoplasmosis has not been studied. TRIUMEQ is not recommended for subjects with CrCl of <50 ml/min. | |
|-----------------------------------------------------------------------------------------------------------|------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|
| mtricitabine | | Lamivudine may inhibit the intracellular phosphorylation of emtricitabine when the two medicinal products are used concurrently. Additionally, the mechanism of viral resistance for both lamivudine and emtricitabine is mediated via mutation of the same viral reverse transcriptase gene (M184V) and therefore the therapeutic efficacy of these drugs in combination therapy may be limited. Lamivudine is not recommended for use in combination with emtricitabine or emtricitabine-containing fixed-dose combinations. | |

| | | emtricitabine is mediated via mutation of the same viral reverse transcriptase gene (M184V) and therefore the therapeutic efficacy of these drugs in combination therapy may be limited. Lamivudine is not recommended for use in combination with emtricitabine or emtricitabine-containing fixed-dose combinations. | | | | |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|--|--|--|
| Other Agents | | | | | | |
| Sorbitol solution (3.2 , 10.2 g, 13.4 g) | Single dose lamivudine oral solution 300 mg Lamivudine: AUC \downarrow 14%; 32%; 36% $c_{max} \downarrow$ 28%; 52%, 55%. | When possible, avoid chronic coadministration of sorbitol-containing medicines with lamivudine. Consider more frequent monitoring of HIV-1 viral load when chronic coadministration cannot be avoided. | | | | |
| lbbreviations: \uparrow = Increase: \downarrow = decrease: \leftrightarrow = no significant change: AUC = area under the concentration versus time curve: C_{++} = maximum | | | | | | |

Pregnancy and Lactation

There are no data on the effects of dolutegravir, abacavir or lamivudine on human male or female fertility. Animal studies indicate no effects of dolutegravir, abacavir or lamivudine on male or female fertility (see Pre-clinical Safety Data).

The safe use of TRIUMEQ in human pregnancy has not been established. Dolutegravir was shown to cross the placenta in reproductive toxicity studies in animals. Lamivudine and abacavir were associated with findings in animal reproductive toxicity studies (see Pre-clinical Safety Data). herefore administration of TRIUMEQ in pregnancy should be considered only if the benefit to the mother outweighs the possible risk to the foetus. here have been reports of mild, transient elevations in serum lactate levels. which may be due to mitochondrial dysfunction, in neonates and infants exposed in utero or peri-partum to nucleoside reverse transcriptase inhibitors (NRTIs). The clinical relevance of transient elevations in serum lactate is unknown. There have also been very rare reports of developmental delay, seizures and other neurological disease. However, a causal relationship between these events and NRTI exposure in utero or peri-partum has not been established. These findings do not affect current recommendations to use antiretroviral therapy in pregnant women to prevent vertical transmission of HIV.

Health experts recommend that where possible HIV infected women do not breast-feed their infants in order to avoid transmission of HIV. In settings where formula feeding is not feasible, local official lactation and treatment quidelines should be followed when considering breast

t is expected that dolutegravir will be secreted into human milk based on animal data, although this has not been confirmed in humans. In a study following repeat oral dose of either 150 mg lamivudine twice daily (given in combination with 300 mg zidovudine twice daily) or 300 mg mivudine twice daily, lamivudine was excreted in human breast milk (0.5 to 8.2 micrograms/ml) at similar concentrations to those found in serur In other studies following repeat oral dose of 150 mg lamivudine twice daily (given either in combination with 300 mg zidovudine or as COMBIVIR or TRIZIVIR) the breast milk:maternal plasma ratio ranged between 0.6 and 3.3. In a study after repeat oral administration of 300 mg abaravir twice daily (given as Trizivir), the breast milk:maternal plasma ratio was 0.9. No pharmacokinetic studies were conducted with abacavir once daily oral inistration. Lamivudine median infant serum concentrations ranged between 18 and 28 ng/mL and were not detectable in one of the studies (assay sensitivity 7 ng/mL). Most infants (8 out of 9) had non-detectable levels of abacavir (assay sensitivity 16 ng/mL). Intracellular carbovir and mivudine triphosphate (active metabolites of abacavir and lamivudine) levels in breastfed infants were not measured therefore the clinical relevance of the serum concentrations of the parent compounds measured is unknown.

Effects on Ability to Drive and Use Machines here have been no studies to investigate the effect of dolutegravir, abacavir or lamivudine, on driving performance or the ability to operate machinery. A detrimental effect on such activities would not be anticipated given the pharmacology of these medicinal products. The clinical status of the patient and the adverse event profile of TRIUMEQ should be borne in mind when considering the patient's ability to drive or operate machinery. PHARMACOLOGICAL PROPERTIES

TRIUMEQ contains dolutegravir, abacavir and lamivudine, therefore the adverse events associated with these may be expected. For many of the adverse events listed it is unclear whether they are related to the active substance, the wide range of other medicinal products used in the

management of HIV infection, or whether they are a result of the underlying disease process Many of the adverse events listed occur commonly (nausea, vomiting, diarrhoea, fever, lethargy, rash) in patients with abacavir hypersensitivity Therefore, patients with any of these symptoms should be carefully evaluated for the presence of this hypersensitivity reaction. If TRIUMEO has been discontinued in patients due to experiencing any one of these symptoms and a decision is made to restart abacavir, this must be done only under direct medical supervision (see Special considerations following an interruption of TRIUMEQ therapy in Warnings and Precautions). Adverse drug reactions for dolutegravic abacavic or lamivudine are listed in the tables below by MedDRA system organ class and by frequency

Frequencies are defined as very common (≥1/10), common (≥1/100, <1/10), uncommon (≥1/1,000, <1/100), rare (≥1/10,000, <1/1000) and ery rare (<1/10,000), including isolated reports. Clinical Trial Data Clinical safety data with TRIUMFO are limited. The adverse reactions observed for the combination of dolutegravic plus abacavic/lamivudine in

analysis of pooled data from Phase IIb to Phase IIIb clinical trials were generally consistent with the adverse reaction profiles for the individual omponents dolutegravir, abacavir and lamivudine. However, the following common treatment-emergent adverse reactions were observed with he combination but were not listed in the prescriber information for any of the individual components: iastrointestinal disorders: abdominal distension, gastro-oesophageal reflux disease, dyspepsia

Psychiatric disorders: nightmare and sleep disorder

Metabolism and nutrition disorders: hypertriglyceridaemia and hyperglycaemia

In addition, fatigue and insomnia were observed at a greater frequency with dolutegravir plus abacavir/lamivudine when compared to the individual components. The frequency category for fatique and insomnia was 'very common' with the combination (previously 'common' with

| System organ class | Dolutegravir | Abacavir | Lamivudine | |
|------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------|--------------------------------------------------------------|--|
| Blood and lymphatic systems disorders | | | Uncommon: neutropenia, anaemia, thrombocytopenia | |
| Immune system disorders | Uncommon: hypersensitivity (see Warnings and Precautions), Immune Reconstitution Syndrome (see Warnings and Precautions) | Common: drug hypersensitivity (see Warnings and Precautions) | | |
| Metabolism and nutrition disorders | | Common: anorexia | | |
| Psychiatric disorders | Common: Insomnia, abnormal dreams, depression | | | |
| | Uncommon: Suicidal ideation or suicide attempt (particularly in patients with a pre-existing history of depression or psychiatric illness) | | | |
| Nervous system disorders | Very common: headache Common: dizziness | Common: headache | Common: headache | |
| Gastrointestinal disorders | Very common: nausea, diarrhoea Common: vomiting, flatulence, abdmonial pain, upper abdominal pain, abdominal discomfort | Common: nausea, vomiting, diarrhoea | Common: nausea, vomiting, upper abdominal pain, diarrhoea | |
| Hepatobiliary disorders | Uncommon: hepatitis | | Uncommon: transient rises in liver enzymes (AST, ALT) | |
| Skin and subcutaneous tissue disorders | Common: rash, pruritus | | Common: rash | |
| General disorders and administration site conditions | Common: fatigue | Common: fever, lethargy, fatigue | Common: fatigue, malaise, fever | |

Changes in laboratory chemistries

Increases in serum creatinine occurred within the first week of treatment with dolutegravir and remained stable through 96 weeks. In G114467, a mean change from baseline of 12.6µmol/L was observed after 96 weeks of treatment. These changes are not considered to b clinically relevant since they do not reflect a change in glomerular filtration rate (see Pharmacodynamics – Effects on Renal Function). Small increases in total bilirubin (without clinical jaundice) were observed on dolutegravir and raltegravir (but not efavirenz) arms in the ogramme. These changes are not considered clinically relevant as they likely reflect competition between dolutegravir and unconjugated bilirubin for a common clearance pathway (UGT1A1) (see Pharmacokinetics — Metabolism). Asymptomatic creatine phosphokinase (CPK) elevations mainly in association with exercise have also been reported with dolutegravir therapy.

Paediatric population There are no clinical study data on the effects of TRIUMEQ in the paediatric population. Individual components have been investigated in

Rased on limited available data with the doluteoravir single entity used in combination with other antiretroviral agents to treat adolescents (12 to less than 18 years of age), there were no additional types of adverse reactions beyond those observed in the adult population. The individual preparations of ABC and 3TC have been investigated separately, and as a dual nucleoside backbone, in combination antiretroviral therapy to treat ART- naive and ART- experienced HIV- infected paediatric patients (data available on the use of ABC and 3TC in children less than three months are limited). No additional types of undesirable effects have been observed beyond those characterised for the adult population

Post-marketing data

In addition to the adverse reactions included from clinical trial data, the adverse reactions listed in Table 5 below have been identified during post-approval use of dolutegravir, abacavir and lamivudine. These events have been chosen for inclusion due to a potential causal connection to dolutegravir, abacavir and/or lamivudine.

Adverse reactions based on post-marketing experience

| System organ class | Dolutegravir | Abacavir | Lamivudine |
|-------------------------------------------------|-------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------|
| Blood and lymphatic systems disorders | | | Very rare: pure red cell aplasia |
| Metabolism and nutrition disorders | | Common: hyperlactataemia Rare: ¹lactic acidosis | Common: hyperlactataemia Rare: ¹lactic acidosis |
| Nervous system disorders | | | Very rare: paraesthesiae, peripheral neuropathy has been reported although a causal relationship to treatment is uncertain |
| Gastrointestinal disorders | | Rare: pancreatitis, but a causal relationship to abacavir is uncertain | Rare: rises in serum amylase, pancreatitis, although a causal relationship to lamivudine is uncertain |
| Skin and subcutaneous tissue disorders | | Common: rash (without systemic symptoms) Very rare: erythema multiforme, Stevens-Johnson syndrome and toxic epidermal necrolysis | Common: alopecia |
| Musculoskeletal and connective tissue disorders | Uncommon: arthralgia Uncommon: myalgia | | Common: arthralgia, muscle disorders Rare: rhabdomyolysis |

Description of Selected Adverse Reaction . Hypersensitivity (see also Warninas and Precautions):

Musculoskeletal:

th abacavir and dolutegravir are associated with a risk for hypersensitivity reactions (HSR), which were observed more commonly with abacavir. Hypersensitivity reaction observed for each of these medicinal products (described below) share some common features such as fever and/or rash with other symptoms indicating multi-organ involvement. Time to onset was typically 10-14 days for both abacavir and dolutegravir-associated reactions, although reactions to abacavir may occur at any time during therapy Dolutearavir hypersensitivity:

Symptoms have included rash, constitutional findings, and sometimes, organ dysfunction, including severe liver reactions. Abacavir hypersensitivity: The signs and symptoms of this hypersensitivity reaction (HSR) are listed below. These have been identified either from clinical studies or post

Rash (usually maculopapular or urticarial)

Elevated creatinine, renal failure

narketing surveillance. Those reported in **at least 10% of patients** with a hypersensitivity reaction are in bold text Almost all patients developing hypersensitivity reactions will have fever and/or rash (usually maculopapular or urticarial) as part of the syndrome, however, reactions have occurred without rash or fever. Other kev symptoms include gastrointestinal, respiratory or constitutional symptoms such as lethargy and malais

Nausea, vomiting, diarrhoea, abdominal pain, mouth ulceration **Dyspnoea, cough**, sore throat, adult respiratory distress syndrome, respiratory failure Fever, fatigue, malaise, oedema, lymphadenopathy, hypotension, conjunctivitis, anaphylaxis Neurological/Psychiatry: Headache, paraesthesia Haematological: Lymphopenia Elevated liver function tests, hepatic failure

Myalgia, rarely myolysis, arthralgia, elevated creatine phosphokinase

Restarting abacavir following an abacavir HSR results in a prompt return of symptoms within hours. This recurrence of the HSR is usually more severe than on initial presentation, and may include life-threatening hypotension and death. Reactions have also occurred infrequently after restarting abacavir in patients who had only one of the key symptoms of hypersensitivity (see above) prior to stopping abacavir; and on very rare occasions have also been seen in patients who have restarted therapy with no preceding symptoms of a HSR (i.e., patients previously considered to be abacavir tolerant).

For details of clinical management in the event of a suspected abacavir HSR see Warnings and Precautions.

Symptoms and Signs e is currently limited experience with overdosage in dolutegravir. Limited experience of single higher doses (up to 250 mg in healthy subjects) revealed no specific symptoms or signs, apart from those listed as adverse reactions.

No specific symptoms or signs have been identified following acute overdose with abacavir or lamivudine. apart from those listed as adverse reactions

Further management should be as clinically indicated or as recommended by the national poisons centre, where available. If overdose occurs, the natient should be treated supportively with appropriate monitoring as pecessary. Since lamivudine is dialysable, continuous haemodialysis could be used in the treatment of overdose, although this has not been studied. It is not known whether abacavir can be removed by peritoneal dialysis or haemodialysis. As dolutegravir is highly bound to plasma proteins, it is unlikely that it will be significantly removed by dialysis.

Mechanism of Action

Antiviral Activity in combination with other antiviral agents

Dolutegravir inhibits HIV integrase by binding to the integrase active site and blocking the strand transfer step of retroviral Deoxyribonucleic acid (DNA) integration which is essential for the HIV replication cycle. Strand transfer biochemical assays using purified HIV 1 integrase and pre-processed substrate DNA resulted in IC50 values of 2.7 nM and 12.6 nM. In vitro, dolutegravir dissociates slowly from the active site of the

Abacavir and lamivudine are NRTIs, and are potent, selective inhibitors of HIV-1 and HIV-2. Both abacavir and lamivudine are metabolised sequentially by intracellular kinases to the respective triphosphate (TP) which are the active moieties with extended intracellular half-lives ng once daily dosing (see Pharmacokinetics, Elimination). Lamivudine-TP and carbovir-TP (the active triphosphate form of abacavir) are substrates for and competitive inhibitors of HIV reverse transcriptase (RT). However, their main antiviral activity is through incorporation of the nophosphate form into the viral DNA chain, resulting in chain termination. Abacavir and lamivudine triphosphates show significantly less affinity for host cell DNA polymerases.

In a randomized, dose-ranging trial, HIV 1—infected subjects treated with dolutegravir monotherapy (ING111521) demonstrated rapid and dose-dependent antiviral activity, with mean declines from baseline to day 11 in HIV-1 RNA of 1.5, 2.0, and 2.5 log10 for dolutegravir 2 mg 10 mg, and 50 mg once daily, respectively. This antiviral response was maintained for 3 to 4 days after the last dose in the 50 mg group. Antiviral Activity in cell culture

Peripheral blood mononuclear cells (PBMCs) infected with HIV-1 strain BaL or HIV-1 strain NL432 gave DTG IC50s of 0.51 nM and 0.53 nM, respectively and MT-4 cell assays with HIV-1 strain IIIB resulted in IC50s of 0.71 and 2.1 nM. When dolutegravir was tested in PBMC assays against a panel consisting of 24 HIV-1 clinical isolates Group M (clade A. B. C. D. E. F and G) and group O] and 3 HIV-2 clinical isolates, the geometric mean ICSO was 0.20 nM and ICSO values ranged from 0.02 to 2.14 nM for HIV-1, while the metric mean IC50 was 0.18 nM and IC50 values ranged from 0.09 to 0.61 nM for HIV-2 isolates.

No drugs with inherent anti-HIV activity were antagonistic with dolutegravir (in vitro assessments were conducted in checkerboard format in combination with stavudine, abacavir, efavirenz, nevirapine, lopinavir, amprenavir, enfuvirtide, maraviroc, adefovir and raltegravir). In addition, antivirals without inherent anti-HIV activity (ribavirin) had no apparent effect on dolutegravir activity. The antiviral activity of abacavir in cell culture was not antagonized when combined with the nucleoside reverse transcriptase inhibitors (NRTIs

didanosine, emtricitabine, lamivudine, stavudine, tenofovir, zalcitabine or zidovudine, the non-nucleoside reverse transcriptase inhibitor (NNRTI nevirapine, or the protease inhibitor (PI) amprenavir. No antagonistic effects in vitro were seen with lamivudine and other antiretrovirals (tested agents: abacavir, didanosine, nevirapine, zalcitabine, and zidovudine). Effect of Human Serum and Serum Proteins The protein adjusted IC90 (PA-IC90) in PBMCs for dolutegravir was estimated to be 64 ng/mL. Dolutegravir trough concentration for a single

50 mg dose in integrase inhibitor naïve subjects was 1.20 µg/mL, 19 times higher than the estimated PA-IC90. Plasma protein binding studies

itro indicate that abacavir binds only low to moderately (\sim 49%) to human plasma proteins at therapeutic concentrations. Lamivudine exhibits linear pharmacokinetics over the therapeutic dose range and displays low plasma protein binding (less than 36%). Resistance in vitro (dolutegravir) Isolation from wild type HIV-1: Viruses highly resistant to dolutegravir were not observed during the 112 day passage of strain IIIB, with a

4.1-fold maximum fold change (FC) observed for the passaged resistant virus populations with substitutions at the conserved IN positions 5153Y and 5153F. Passage of the wild type HIV-1 strain NL432 in the presence of dolutegravir selected for E920 (passage population virus FC=3.1) and G193E (passage population virus FC=3.2) substitutions on Day 56. Additional passage of wildtype subtype B, C, and A/G viruses in the presence of dolutegravir selected for R263K, G118R, and S153T. Resistance in vivo (dolutegravir): integrase inhibitor naïve patients

No INI-resistant mutations or treatment emergent resistance to the NRTI backbone therapy were isolated with dolutegravir 50 mg once daily in treatment—naive studies (SPRING-1 SPRING-2 SINGLE FLAMING) and ARIA studies). In the SAILING study for treatment experienced (and integrase naïve) patients (n=354 in the dolutegravir arm), treatment emergent integrase mutations were observed at Week 48 in 4 of 17 subjects with virologic failure in the dolutegravir arm. Of these four 2 subjects had a unique R263K integrase substitution, with a maximum FC of 193 subject had a polymorphic V151V/I integrase substitution, with maximum FC of 0.92, and 1 subject had pre-existing integrase mutations and is assumed to have been integrase experienced or infected with integrase resistant virus by transmission (see Clinical Studies) Resistance in vitro and in vivo (abacavir and lamivudine)

Abacavir-resistant isolates of HIV-1 have been selected in vitro and in vivo and are associated with specific genotypic changes in the RT codon region (codons M184V, K65R, L74V and Y115F). During *in vitro* abacavir selection the M184V mutation occurred first and resulted in about a two-fold increase in ICSO, below the abacavir clinical cutoff of 4.5 FC. Continued passage in increasing concentrations of drug resulted in selection for double RT mutants 65R/184V and 74V/184V or triple RT mutant 74V/115Y/184V. Two mutations conferred a 7- to 8-FC in abacavir susceptibility and combinations of three mutations were required to confer more than an 8-FC in susceptibility. HIV-1 resistance to lamivudine involves the development of a M184I or M184V amino acid change close to the active site of the viral RT This

variant arises both in vitro and in HIV-1 infected patients treated with lamivudine-containing antiretroviral therapy. M184V mutants display greatly reduced susceptibility to lamivudine and show diminished viral replicative capacity in vitro. M184V is associated with a low level increase in abacavir resistance but does not confer clinical resistance for abacavir Isolates resistant to abacavir may also show reduced sensitivity to lamivudine. The combination of abacavir/lamivudine has demons decreased susceptibility to viruses with the substitutions K65R with or without the M184V/I substitution, and to viruses with L74V plus the

In a randomized, placebo-controlled, cross-over trial, 42 healthy subjects received single dose oral administrations of placebo, dolutegravi 250 mg suspension (exposures approximately 3-fold of the 50 mg once-daily dose at steady state), and moxifloxacin (400 mg, active control) in

random sequence. Dolutegravir did not prolong the QTc interval for 24 hours post dose. After baseline and placebo adjustment, the maximum mean QTc change based on Fridericia correction method (QTcF) was 1.99 msec (1-sided 95% upper Cl: 4.53 msec). Similar studies were not conducted with either abacavir or lamivudine. Effects on Renal Function The effect of dolutegravir on serum creatinine clearance (CrCl), glomerular filtration rate (GFR) using iohexol as the probe and effective renal plasm

flow (ERPF) using para-aminohippurate (PAH) as the probe was evaluated in an open-label, randomized, 3 arm, parallel, placebo-controlled study in 37 healthy subjects, who were administered dolutegravir 50 mg once daily (n=12), 50 mg twice daily (n=13) or placebo once daily (n=12) for 14 days. A modest decrease in CrCl was observed with dolutegravir within the first week of treatment, consistent with that seen in clinical studies

Metabolism

Elimination

Effects on Electrocardiogram

Dolutegravir at both doses had no significant effect on GFR or ERPF. These data support in vitro studies which suggest that the small increases in eatinine observed in clinical studies are due to the nonpathologic inhibition of the organic cation transporter 2 (OCT2) in the proximal renal ubules, which mediates the tubular secretion of creatinin The TRIUMEQ tablet has been shown to be bioequivalent to TIVICAY with KIVEXA tablets administered separately. This was demonstrated in a single dose, 2-way crossover bioequivalence study of TRIUMEO (fasted) versus 1 x 50 mg dolutegravir tablet, plus 1 x 600 mg abacavir/300 mg

mivudine tablet (fasted) in healthy subjects (n=62). In a separate cohort there was no clinically significant effect of a high fat meal on the exposure of dolutegravir. abacavir or lamivudine. These results indicate that TRIUMEO can be taken with or without food. The pharmacokinetic properties of dolutegravir, lamivudine and abacavir are described below

Absorption Dolutegravir, abacavir and lamivudine are rapidly absorbed following oral administration. The absolute bioavailability of dolutegravir has not een established. The absolute bioavailability of oral abacavir and lamivudine in adults is 83% and 80 to 85% respectively. The mean time t maximal serum concentrations (t_...) is about 2 to 3 hours (post dose for tablet formulation) for dolutegravir, 1.5 hours for abacavir and 1.0 hours

Following multiple oral doses of dolutegravir 50mg once daily, the geometric mean steady state pharmacokinetic parameter estimates are

53.6 micrograms.h/mL for AUC₃₄, 3.67 microgram/mL for C_{max}, and 1.11 microgram/mL for C₃₄. Following a single oral dose of 600 mg of

lamivudine 300 mg once daily for seven days the mean steady-state C_{max} is 2.04 micrograms/ml and the mean AUC_{24} is 8.87 micrograms.h/ml. Distribution The apparent volume of distribution of dolutegravir (following oral administration of suspension formulation, Vd/F) is estimated at 12.5 l travenous studies with abacavir and lamivudine showed that the mean apparent volume of distribution is 0.8 and 1.3 l/kg respectively Dolutegravir is highly bound (approximately 99.3%) to human plasma proteins based on in vitro data. Binding of dolutegravir to plasma protein

was independent of concentration. Total blood and plasma drug-related radioactivity concentration ratios averaged between 0.441 to 0.535 indicating minimal association of radioactivity with blood cellular components. Free fraction of dolutegravir in plasma is estimated a proximately 0.2 to 1.1% in healthy subjects, approximately 0.4 to 0.5% in subjects with moderate hepatic impairment, and 0.8 to 1.0% i ubiects with severe renal impairment and 0.5% in HIV-1 infected patients. Plasma protein binding studies *in vitro* indicate that abacavir binds only low to moderately (approximately 49%) to human plasma proteins at therapeutic concentrations. Lamivudine exhibits linear narmacokinetics over the therapeutic dose range and displays low plasma protein binding (less than 36%). Dolutegravir abacavir and lamivudine are present in cerebrospinal fluid (CSF). In 12 treatment-paive subjects receiving a regimen of dolutegravir

plus abacavir/lamivudine for 16 weeks, dolutegravir concentration in CSF averaged 15.4 ng/mL at Week 2 and 12.6 ng/mL at Week 16, ranging om 3.7 to 23.2 ng/ml. (comparable to unbound plasma concentration). CSF-plasma concentration ratio of DTG ranged from 0.11 to 2.0496 olutegravir concentrations in CSF exceeded the ICSO, supporting the median reduction from baseline in CSF HIV-1 RNA of 2.2 log after 2 weeks and 3.4 log after 16 weeks of the rany (see Pharmacodynamics). Studies with abacavir demonstrate a CSF to plasma ALIC ratio of between 3.0 to 44%. The observed values of the peak concentrations are 9 fold greater than the IC_{so} of abacavir of 0.08 micrograms/ml or 0.26 micromolar when abacavir is given at 600 mg twice daily. The mean ratio of CSF/serum lamiyudine concentrations 2 to 4 h after oral administration was approximately 12%. The true extent of CNS penetration of lamivudine and its relationship with any clinical efficacy is unknown. . Olutegravir is present in the female and male genital tract. AUC in cervicovaginal fluid, cervical tissue, and vaginal tissue were 6 to 10% of that

Dolutegravir is primarily metabolized via UGT1A1 with a minor CYP3A component (9.7% of total dose administered in a human mass balance dy). Dolutegravir is the predominant circulating compound in plasma; renal elimination of unchanged drug is low (<1% of the dose). Fifty-three percent of total oral dose is excreted unchanged in the faeces. It is unknown if all or part of this is due to unabsorbed drug or biliary cretion of the glucuronidate conjugate, which can be further degraded to form the parent compound in the gut lumen. Thirty-one percent of the total oral dose is excreted in the urine, represented by ether glucuronide of dolutegravir (18.9% of total dose). N-dealkylation metabolite .6% of total dose), and a metabolite formed by oxidation at the benzylic carbon (3.0% of total dose

in corresponding plasma at steady-state. AUC was 7% in semen and 17% in rectal tissue, of those in corresponding plasma at steady-state.

primary pathways of metabolism in man are by alcohol dehydrogenase and by glucuronidation to produce the 5'-carboxylic acid and uronide which account for about 66% of the administered dose. These metabolites are excreted in the urine. Metabolism of lamivudine is a minor route of elimination. Lamivudine is predominately cleared unchanged by renal excretion. The likelihood of metabolic interactions with lamivudine is low due to the small extent of hepatic metabolism (less than 10%).

Dolutegravir has a terminal half-life of \sim 14 hours and an apparent clearance (CL/F) of 0.56 L/hr.

Abacavir is primarily metabolised by the liver with less than 2% of the administered dose being renally excreted as unchanged compound. The

hepatic metabolism with subsequent excretion of metabolites primarily in the urine. The metabolites and unchanged abacavir account for about 83% of the administered abacavir dose in the urine. The remainder is eliminated in the faeces The observed lamivudine half-life of elimination is 5 to 7 hours. For patients receiving lamivudine 300 mg once daily, the terminal intracellular half-life of lamivudine-TP was prolonged to 16 to 19 hours. The mean systemic clearance of lamivudine is approximately 0.32 l/h/kg, The statistically higher responses on DTG+ABC/3TC were driven by withdrawals due to AEs, irrespective of viral load strata. At 144 weeks in the predominantly by renal clearance (greater than 70%) via the organic cationic transport system.

The mean half-life of abacavir is about 1.5 hours. The geometric mean terminal half-life of intracellular carbovir-TP at steady-state is 20.6 hours.

ollowing multiple oral doses of abacavir 300 mg twice a day, there is no significant accumulation of abacavir. Elimination of abacavir is vi

Special Patient Populations

In a paediatric study including 23 antiretroviral treatment-experienced HIV-1 infected children and adolescents aged 12 to 18 years of age, the pharmacokinetics of dolutegravir was evaluated in 10 children and showed that dolutegravir 50 mg once daily dosage resulte exposure in paediatric subjects comparable to that observed in adults who received dolutegravir 50 mg once daily (Table 6). Paediatric pharmacokinetic parameters (n=10)

| Age/weight | Dolutegravir Dose | Dolutegravir Pharmacokinetic Parameter Estimates Geometric Mean (CV%) | | | |
|------------------------------------------------------|----------------------|--------------------------------------------------------------------------|---------------------------|--------------------------|--|
| | | AUC ₍₀₋₂₄₎ μg.hr/mL | C _{max} µg/mL | C ₂₄ μg/mL | |
| 12 to <18 years ≥ 40 kg a | 50 mg once daily a | 46 (43) | 3.49 (38) | 0.90 (59) | |
| One subject weighing 27 kg received 25 mg once daily | | | | | |

Limited data are available in adolescents receiving a daily dose of 600 mg of abacavir and 300 mg of lamivudine. Pharmacokinetic parameters are comparable to those reported in adults.

Population pharmacokinetic analysis of dolutegravir using data in HIV-1 infected adults showed that there was no clinically relevant effect of age on dolutegravir exposure. Pharmacokinetic data for dolutegravir, abacavir and lamivudine in subjects of >65 years old are limited.

Hepatically impaired Pharmacokinetic data has been obtained for dolutegravir, abacavir and lamivudine alone. Based on data obtained for abacavir, TRIUMEQ is not

recommended in natients with moderate and severe henatic impairment Abacavir is metabolised primarily by the liver. The pharmacokinetics of abacavir have been studied in patients with mild hepatic impairment

Pugh score 5 to 6). The results showed that there was a mean increase of 1.89 fold in the abacavir AUC and 1.58 fold in the half-life of abacavir. The AUCs of the metabolites were not modified by the liver disease. However, the rates of formation and elimination of these were ecreased. Dosage reduction of abacavir may be required in patients with mild hepatic impairment. The separate preparation of ZIAGEN should herefore be used to treat these patients. The pharmacokinetics of abacavir have not been studied in patients with moderate or severe hepatic mpairment. Plasma concentrations of abacavir are expected to be variable and substantially increased in these patients. TRIUMEQ is therefore ot recommended in patients with moderate and severe hepatic impairment. Data obtained for lamivudine in natients with moderate to severe benatic impairment and for dolutegravir in natients with moderate benatic

impairment show that the pharmacokinetics are not significantly affected by hepatic dysfunction. Dolutegravir is primarily metabolized and eliminated by the liver. In a study comparing 8 subjects with moderate hepatic impairment (Child-Pugh category B) to 8 matched healthy adult ontrols, the single 50 mg dose exposure of dolutegravir was similar between the two groups. The effect of severe hepatic impairment on the pharmacokinetics of dolutegravir has not been studied.

Renally impaired

Pharmacokinetic data have been obtained for dolutegravir, abacavir and lamivudine alone. TRIUMEQ should not be used in patients with creatinine e of less than 50 mL/min because, whilst no dosage adjustment of dolutegravir or abacavir is necessary in patients with renal impairment dose reduction is required for the lamivudine component. Therefore the separate preparation of EPIVIR should be used to treat these patients. Studies with lamivudine show that plasma concentrations (AUC) are increased in patients with renal dysfunction due to decreased clearance. Abacavir is primarily metabolised by the liver, with approximately 2% of abacavir excreted unchanged in the urine. The pharmacokinetics of

Renal clearance of unchanged drug is a minor pathway of elimination for dolutegravir. A study of the pharmacokinetics of dolutegravir was performed in subjects with severe renal impairment (CLcr < 30 mL/min). No clinically important pharmacokinetic differences between subjects ith severe renal impairment (CLcr < 30 mL/min) and matching healthy subjects were observed. Dolutegravir has not been studied in patients on dialysis, though differences in exposure are not expected. olymorphisms in Drug Metabolising Enzymes

There is no evidence that common polymorphisms in drug metabolising enzymes alter dolutegravir pharmacokinetics to a clinically meaningful extent. In a meta-analysis using pharmacogenomics samples collected in clinical studies in healthy subjects, subjects with UGT1A1 (n=7) genotypes conferring poor dolutegravir metabolism had a 32% lower clearance of dolutegravir and 46% higher AUC compared with subjects ith genotypes associated with normal metabolism via UGT1A1 (n=41). Polymorphisms in CYP3A4, CYP3A5, and NR112 were not associated with differences in the pharmacokinetics of dolutegravir.

The dolutegravir exposure in healthy subjects appear to be slightly higher (~20%) in women than men based on data obtained in a healthy subject study (males n=17, females n=24). Population PK analyses using pooled pharmacokinetic data from Phase IIb and Phase III adult trials revealed no clinically relevant effect of gender on the exposure of dolutegravir. There is no evidence that a dose adjustment of dolutegravir, abacavir or lamivudine would be required based on the effects of gender on PK

Population PK analyses using pooled pharmacokinetic data from Phase IIb and Phase III adult trials revealed no clinically relevant effect of race on the exposure of dolutegravir. The pharmacokinetics of dolutegravir following single dose oral administration to Japanese subjects appear similar

to observed parameters in Western (US) subjects. There is no evidence that a dose adjustment of dolutegravir, abacavir or lamivudine would be required based on the effects of race on PK parameters. Co-infection with Hepatitis B or C

Population PK analysis indicated that hepatitis C virus co-infection had no clinically relevant effect on the exposure to dolutegravir. There are nited pharmacokinetic data on subjects with hepatitis B co-infection (see Warnings and Precautions for the use of TRIUMEQ in patients co-infected with hepatitis B).

Clinical Studies Antiretroviral naïve subjects

The efficacy of TRIUMEQ in HIV-infected, antiretroviral therapy naive subjects is based on the analyses of data from three studies; SINGLE (ING114467), SPRING-2 (ING113086) and FLAMINGO (ING114915).

n SINGLE, 833 subjects were randomized and received at least one dose of either dolutegravir 50 mg once daily with fixed-dose abacavir-lamivudine (DTG + ABC/3TC) or fixed-dose efavirenz-tenofovir-emtricitabine (EFV/TDF/FTC). At baseline, median patient age was 5 years, 16% were female, 32% non-white, 7% had hepatitis C co-infection and 4% were CDC Class C, these characteristics were similar

48 Weeks

between treatment groups. Virologic outcomes (including outcomes by key baseline covariates) are described below Virologic Outcomes of Randomized Treatment of SINGLE at 48 Weeks (Snapshot algorithm)

DTC FO ADC/DTC

| | DTG 50 mg + ABC/3TC Once Daily N=414 | EFV/TDF/FTC Once Daily N=419 | Lamivudine has not shown any genotoxic activity show any carcinogenic potential at exposures app Reproductive Toxicology | | |
|-----------------------------------------------------------------|--------------------------------------------|------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------|--|--|
| HIV-1 RNA <50 copies/mL | 88% | 81% | Fertility | | |
| Treatment Difference* 7.4% (95% Cl: 2.5%, 12.3%) | | | Fertility studies in the rat have shown that dolute Dolutegravir did not affect male or female fertility | | |
| Virologic non response† | 5% | 6% | clinical exposure when dolutegravir is administere | | |
| No virologic data at Weeks 48 window | 7% | 13% | Pregnancy In reproductive toxicity studies in animals, dolu | | |
| Reasons | | | Oral administration of dolutegravir to pregnant ra developmental toxicity or teratogenicity (50 time abacavir and lamivudine. based on AUC). | | |
| Discontinued study/study drug due to adverse event or death‡ | 2% | 10% | Oral administration of dolutegravir to pregnant ra developmental toxicity or teratogenicity (0.74 tim | | |
| Discontinued study/study drug for other reasons§ | 5% | 3% | abacavir and lamivudine, based on AUC). In rabbi weight gain) was observed at 1000 mg/kg (0.74 ti | | |
| Missing data during window but on study | 0 | <1% | with abacavir and lamivudine, based on AUC). Abacavir demonstrated toxicity to the developing | | |
| | HIV-1 RNA <50 copies/mL by baseline cov | ariates | approximately 28 times human therapeutic exposi findings included foetal oedema, variations and m | | |
| Baseline Plasma Viral Load (copies/mL) | n / N (%) | n / N (%) | which there were no effects on pre or post-natal d humans. Similar findings were not observed in rab | | |
| ≤100,000 | 253 / 280 (90%) | 238 / 288 (83%) | Lamivudine was not teratogenic in animal studies | | |
| >100,000 | 111 / 134 (83%) | 100 / 131 (76%) | levels comparable to those achieved in man. How 32 times the clinical exposure (based on C _{max}). | | |
| Baseline CD4+ (cells/ mm³) | | | Animal toxicology and/or pharmacology | | |
| <200 | 45 / 57 (79%) | 48 / 62 (77%) | The effect of prolonged daily treatment with high o | | |
| 200 to <350 | 143 / 163 (88%) | 126 / 159 (79%) | 26 weeks) and in monkeys (up to 38 weeks). The produce systemic exposures approximate | | |
| ≥350 | 176 / 194 (91%) | 164 / 198 (83%) | combination with abacavir and lamivudine, based | | |
| Gender | | | administration, mg/kg or mg/m² metrics are appro the human mg/kg equivalent dose (based on 50 kg | | |
| Male | 307 / 347 (88%) | 291 / 356 (82%) | Mild myocardial degeneration in the heart of mice | | |
| Female | 57 / 67 (85%) | 47 / 63 (75%) | exposures were equivalent to 7 to 21 times huma | | |
| Race | | | lamivudine. The clinical relevance of this finding h | | |
| White | 255 / 284 (90%) | 238 / 285 (84%) | List of Excipients | | |
| African- American/African Heritage/Other | 109 / 130 (84%) | 99 / 133 (74%) | Tablet core: D-Mannitol Microcrystalline cellulose | | |
| Age (years) | | | Povidone K29/32 | | |
| <50 | 319 / 361 (88%) | 302 / 375 (81%) | Sodium starch glycolate Magnesium stearate | | |

t Includes subjects who discontinued prior to Week 48 for lack or loss of efficacy and subjects who are ≥50 copies in the 48 week window. ‡ Includes subjects who discontinued due to an adverse event or death at any time point from Day 1 through the Week 48 analysis window if this resulted in no virologic data on treatment during the analysis window. reasons such as withdrew consent, loss to follow-up, moved, protocol deviatio Notes: ABC/3TC = abacavir 600 mg, lamivudine 300 mg in the form of Kivexa/Epzicom fixed dose combination (FDC)

FV/TDF/FTC = efavirenz 600 mg, tenofovir 300 mg, emtricitabine 200 mg in the form of Atripla FDC. N = Number of subjects in each treatment group In the primary 48 weeks analysis in the SINGLE study, the proportion of patients with virologic suppression (HIV-1 RNA <50 copies/mL) in the

45 / 53 (85%)

olutegravir + ABC/3TC arm (88%), was superior to the EFV/TDF/FTC arm (81%), p=0.003, similar treatment difference was observed in subjects defined by baseline HIV RNA level (< or > 100.000 copies/mL). The median time to viral suppression was 28 days in the group receiving dolutegravir + ABC/3TC and 84 days in the EFV/TDF/FTC arm (p<0.0001). The adjusted mean change in CD4+ T cell count from baseline were 267 cells/mm³ in the group receiving dolutegravir + ABC/3TC and 208 cells/mm³ for the EFV/TDF/FTC arm in SINGLE at 48 weeks [adjusted ifference between arm (with 95% CI), 58.9 cells (33.4 cells to 84.4 cells), p<0.001]. Both the time to viral suppression and change from baseline analyses were pre-specified and adjusted for multiplicity At 96 weeks, 80% of study participants on the DTG + ABC/3TC regimen were virologically suppressed (<50 copies/mL) vs. 72% of participants on

FV/TDF/FTC) [difference and 95% CI; 8.0% (+2.3% to +13.8%); the difference in the endpoint remained statistically significant, [p=0.006].

open-label phase, virologic suppression was maintained, the dolutegravir + ABC/3TC arm (71%) was superior to the EFV/TDF/FTC arm (63%), In SPRING-2, 822 adults were randomized and received at least one dose of either dolutegravir 50 mg once daily or raltegravir 400 mg twice daily.

both administered with fixed-dose dual NRTI therapy (either ABC/3TC or TDF/FTC). Of these subjects, 169/411 in the group receiving dolutegraving nd 164/411 in the group receiving raltegravir were receiving ABC/3TC background regimen. At baseline, median patient age was 36 years, 14% were female, 15% non-white, 12% had hepatitis B and/or C co-infection, and 2% were CDC Class C; these characteristics were similar between treatment groups. Overall virologic suppression (HIV-1 RNA <50 copies/mL) observed with either background regimen in the dolutegravir group (88%) was non-inferior to the raltegravir group (85%) at 48 weeks. The adjusted difference in proportion and 95% CI were 2.5 (-2.2, 7.1). At 96 weeks, virologic suppression in the dolutegravir group (81%) was still non-inferior to the raltegravir group (76%). The adjusted difference in proportion and 95% CI were 4.5 (-1.1, 10.0). Response rates at 48 weeks (and 96 weeks) were 86% (and 74%) for dolutegravir + ABC/3TC and 87% (and 76%) for raltegravir + ABC/3TC, respectively.

In both the SINGLE and SPRING-2 studies of virologic suppression (HIV-1 RNA <50 copies/mL), treatment differences were comparable across baseline characteristics (gender, race and age).

Through 96 weeks in SINGLE and SPRING-2, no INI-resistant mutations or treatment emergent resistance in background therapy were isolated on the dolutegravir-containing arms. In SPRING-2, four subjects on the raltegravir arm failed with major NRTI mutations and one subject developed raltegravir resistance; in SINGLE, six subjects on the EFV/TDF/FTC arm failed with mutations associated with NNRTI resistance and one developed a major NRTI mutation.

In FLAMINGO, an open-label and active-controlled study, 485 HIV-1 infected antiretroviral naïve adults were randomized and received at least one dose of either dolutegravir 50 mg once daily or darunavir/ritonavir (DRV/r) 800 mg/100 mg once daily, both administered with fixed-dose dual NRTI therapy (either ABC/3TC orTDF/FTC). Of these subjects, 33% in both groups were receiving ABC/3TC background regimen. At baseline, median patient age was 34 years, 15% were female, 28% non-white, 10% had hepatitis B and/or C co-infection, and 3% were CDC Class C; these haracteristics were similar between treatment groups. Overall virologic suppression (HIV-1 RNA <50 copies/mL) in the dolutegravir group (90%) was superior to the DRV/r group (83%) at 48 weeks. The adjusted difference in proportion and 95% CI were 7.1 (+0.9 + 13.2) [p=0.025]. At 96 weeks virologic suppression in the dolutegravir group (80%) was superior to the DRV/r group (68%). The median time to viral suppression was 28 days in the DTG treatment group and 85 days in the DRV/r arm (p<0.001). Response rates at 48 weeks were 90% for dolutegravir + ABC/3TC and 85% for DRV/r/ABC/3TC and at 96 weeks were 82% for dolutegravir + ABC/3TC and 75% for DRV/r/ABC/3TC. No subjects in the study had treatment-emergent primary resistance mutations.

Sustained virological response was demonstrated in the SPRING-1 study (ING112276), in which 88% of patients receiving dolutegravir 50 mg (n=51) once daily had HIV-1 RNA <50 copies/mL, compared to 72% of patients in the efavirenz group (n=50) at 96 weeks. No INI-resistant mutations or treatment emergent resistance in background therapy were isolated with dolutegravir 50 mg once daily through 96 weeks. Antiretroviral naive female subjects

In Aria (ING117172), a randomized, open-label, active-controlled, multicenter, parallel group, non-inferiority study; 499 HIV-1 infected ART naïve adult women were randomized 1:1 to receive either; DTG/ABC/3TC FDC 50 mg/600 mg/300 mg; or atazanavir 300 mg plus ritonair 100mg plus tenofovir disproxil fumarate/ emtricitabine 300 mg/200 mg (ATV+RTV+TDF/FTC FDC), all administered once daily. Demographic characteristics were similar across treatment groups, at baseline the median patient age was 37 years, 45% white and 42% African American/African Heritage 93% tested negative for hepatitis C (HCV) infection and 84% of subjects were in CDC Class A.

At 48 weeks overall virologic suppression (HIV-1 RNA <50 copies/mL) in the DTG/ABC/3TC FDC group (82%) was shown to be statistically superior to the ATC+RTV+TDF/FTC FDC group (71%). The adjusted difference in proportion and 95% CI were between 10.5 (3.1% to 17.8%) [p=0.005]. Antiretroviral experienced subjects

The efficacy of the TRIUMEO is also supported by data from a randomized, international, double-blind, active-controlled trial, SAILING (ING111762). In the SAILING study, 719 HIV-1 infected, ART-experienced, integrase inhibitor naive adults were randomized and reci 50 mg once daily or raltegravir 400 mg twice daily with investigator selected background regimen (BR) consisting of 2 agents (including at least one fully active agent). At baseline, median patient age was 43 years, 32% were female, 50% non-white, 16% had hepatitis B and/o co-infection, and 46% were CDC Class C. All subjects had at least two class ART resistance, and 49% of subjects had at least 3-class ART resistance at aseline. Virologic suppression (HIV-1 RNA < 50 copies/mL) in the dolutegravir arm (71%) was statistically superior to the raltegravir arm (64%), at Week 48 (p=0.030). Virologic suppression (HIV-1 RNA <50 copies/mL) treatment differences were comparable across the baseline characteristic of gender, race, and HIV sub type. In STRIIVING (201147) a 48-week, randomized, open-label, active controlled, multicenter, non-inferiority study; 555 HIV-1 infected, virologically

suppressed (HIV-1 RNA <50 c/mL) subjects were randomly assigned (1:1) to continue their current ART regimen (2 NRTIs plus either a PI, NNRTI r INI), or switch to ABC/DTG/3TC FDC once daily (Early Switch). he majority of subjects in the intent-to-treat exposed (ITT-E) population were white (65%) and male (86%); the median age was 45 (range 22-80) years. At Baseline, 31% of subjects had CD4+ counts of <500 cell/mm³. Overall, most subjects had negative test results at screening for

HBV and HCV infection (93%), were in CDC Class A (73%), and identified homosexual activity as an HIV risk factor (72%).

dolutegravir was evaluated in combination regimens in HIV-1 infected infants, children and adolescents.

in vivo rat micronucleus test with abacavir and lamivudine in combination were negative.

Carcinogenesis/mutagenesis

 $Virologic \, suppression \, (HIV-1 \, RNA < 50 \, copies/mL) \, in \, the \, ABC/DTG/3TC \, FDC \, group \, (85\%) \, was \, statistically \, non-inferior \, to \, the \, current \, ART \, groups \, (1975) \, and \, (1975) \, are a statistically \, continuous \, (1975) \, and \, (1975) \, are a statistically \, continuous \, (1975) \, and \, (1975) \, are a statistically \, continuous \, (1975) \, are a statistically \, (1975) \, are a statistically \, continuous \, (1975) \, are a statistically \, (1975) \, are a statistically \, (1975) \, ar$ (88%) at 24 weeks. The adjusted difference in proportion and 95% CI [ABC/DTG/3TC vs current ART] were 3.4%; 95% CI: [-9.1, 2.4]. After reeks all remaining subjects switched to ABC/DTG/3TC FDC (Late Switch). Similar levels of virologic suppression were maintained in both the Early and Late Switch groups at 48 weeks. In CAL30001 and ESS30008 ABC/3TC and ABC + 3TC were effectively used in combination therapy to maintain viral suppression in treatment rienced subjects with low rates of treatment emergent viral resistance mutations

In a Phase I/II 48 week multicentre, open-label study (P1093/ING112578), the pharmacokinetic parameters, safety, tolerability and efficacy of

At 24 weeks, 16 of 23 (69%) adolescents (12 to less than 18 years of age) treated with dolutegravir once daily (35 mg n=4, 50 mg n=19) plus

Pre-clinical Safety Data With the exception of a negative in vivo rat micronucleus test for the combination of abacavir and lamivudine, there are no data available on the effects of the combination of dolutegravir, abacavir and lamivudine in animals.

Dolutegravir was not mutagenic or clastogenic using in vitro tests in bacteria and cultured mammalian cells. and an in vivo rodent micronucleus ssay. Dolutegravir was not carcinogenic in long term studies in the mouse and rat. Neither abacavir nor lamivudine were mutagenic in bacterial tests, but like many nucleoside analogues they show activity in the *in vitro* mammalian tests such as the mouse lymphoma assay. This is consistent with the known activity of other nucleoside analogues. The results of an

Carcinogenicity studies with orally administered abacavir in mice and rats showed an increase in the incidence of malignant and non-malignant umours. Malignant tumours occurred in the preputial gland of males and the clitoral gland of females of both species, and in the liver, urinary bladder. lymph nodes and the subcutis of female rats. The majority of these tumours occurred at the highest abacavir dose of 330 mg/kg/day in mice and 600 mg/kg/day in rats. These dose levels were equivalent to 21 to 28 times the expected systemic exposure in humans when abacavir is administered in combination with dolutegravir and

lamivudine. The exception was preputial gland tumours in mice which occurred at a dose of 110 mg/kg. Exposure at this dose is approximately times the expected human systemic exposure. There is no structural counterpart for this gland in humans. While the carcinogenic potential in humans is unknown, these data suggest that a carcinogenic risk to humans is outweighed by the potential clinical benefit. Lamivudine has not shown any genotoxic activity in the in vivo studies. The results of long-term carcinogenicity studies in rats and mice did not show any carcinogenic potential at exposures approximately 12 to 72 times higher than clinical plasma levels.

Fertility studies in the rat have shown that dolutegravir, abacavir and lamivudine had no effect on male or female fertility. Dolutegravir did not affect male or female fertility in rats at doses up to 1000 mg/kg/day, the highest dose tested (44 times the 50 mg human clinical exposure when dolutegravir is administered in combination with abacavir and lamivudine based on AUC).

In reproductive toxicity studies in animals, dolutegravir, abacavir and lamivudine were shown to cross the placenta. Oral administration of dolutegravir to pregnant rats at doses up to 1000 mg/kg daily from days 6 to 17 of gestation did not elicit maternal toxicity, levelopmental toxicity or teratogenicity (50 times the 50 mg human clinical exposure when dolutegravir is administered in combination wit abacavir and lamivudine, based on AUC). Oral administration of dolutegravir to pregnant rabbits at doses up to 1000 mg/kg daily from days 6 to 18 of gestation did not elicit

evelopmental toxicity or teratogenicity (0.74 times the 50 mg human clinical exposure when dolutegravir is admin abacavir and lamivudine, based on AUC). In rabbits, maternal toxicity (decreased food consumption, scant/no faeces/urine, suppressed body weight gain) was observed at 1000 mg/kg (0.74 times the 50 mg human clinical exposure when dolutegravir is administered in combination with abacavir and lamivudine, based on AUC). Abacavir demonstrated toxicity to the developing embryo and foetus only in rats at maternally toxic doses of 500 mg/kg/day and above. This dose is pproximately 28 times human therapeutic exposure based on AUC, for a 600 mg dose in combination with dolutegravir and lamivudine. The findings included foetal oedema, variations and malformations, resorptions, decreased foetal body weight and an increase in still births. The dose at

humans. Similar findings were not observed in rabbits. Lamivudine was not teratogenic in animal studies, but there were indications of an increase in early embryonic deaths in rabbits at exposure evels comparable to those achieved in man. However, there was no evidence of embryonic loss in rats at exposure levels of approximately 32 times the clinical exposure (based on C_{max}).

which there were no effects on pre or post-natal development was 160 mg/kg/day. This dose is equivalent to an exposure of about 9 times that in

The effect of prolonged daily treatment with high doses of dolutegravir has been evaluated in repeat oral dose toxicity studies in rats (up to 26 weeks) and in monkeys (up to 38 weeks). The primary effect of dolutegravir was gastrointestinal intolerance or irritation in rats and monkeys at doses that produce systemic exposures approximately 38 and 1.5 times the 50 mg human clinical exposure when dolutegravir is administered ombination with abacavir and lamivudine, based on AUC, respectively. Because gastrointestinal (GI) intolerance is considered to be due to local drug administration, mg/kg or mg/m² metrics are appropriate determinates of safety cover for this toxicity. GI intolerance in monkeys occurred at 30 time the human mg/kg equivalent dose (based on 50 kg human), and 11 times the human mg/m² equivalent dose for a total daily clinical dose of 50 mg. Mild myocardial degeneration in the heart of mice and rats was observed following administration of abacavir for two years. The systemic exposures were equivalent to 7 to 21 times human exposure at 600 mg when abacavir is administered in combination with dolutegravir and lamivudine. The clinical relevance of this finding has not been determined PHARMACEUTICAL PARTICULARS

List of Excipients Tablet core:

Microcrystalline cellulose Povidone K29/32 Sodium starch glycolate Magnesium stearate <u>Tablet coating:</u> Opadry II Purple 85F90057 containing Polyvinyl alcohol – part hydrolyzed Titanium oxide

Macrogol/PEG Iron Oxide Black Iron Oxide Red Incompatibilities

No incompatibilities have been identified Shelf Life The expiry date is indicated on the packaging.

Special Precautions for Storage Do not store above 30°C.

Store in the original package in order to protect from moisture. Keep the bottle tightly closed. Do not remove the desiccant Nature and Contents of Container TRIUMEQ tablets are available in white high density polyethylene (HDPE) bottles containing a desiccant.

Instructions for Use/Handling There are no special requirements for use or handling of this product.

Not all presentations are available in every country.

Glaxo Wellcome, S.A.*, Avenida de Extremadura 3, 09400 Aranda De Duero, Burgos, Spain

*Member of the GSK group of companies TRIUMEQ, TIVICAY, ZIAGEN, EPIVIR, KIVEXA, EPZICOM, TRIZIVIR, COMBIVIR are trade marks of the ViiV Healthcare group of companies © 2017 ViiV Healthcare group of companies. All Rights Reserved

ersion number: GDS08/IPI08

