

Read all of this leaflet carefully before you start taking this medicine: - Keep this leaflet. You may need to read it again. - If you have any further questions, ask your doctor or phar

macist.

- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are

of to studies may hard the same as yours.

- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

1. WHAT ADENURIC® IS AND WHAT IT IS USED

Adenuric® works by reducing uric acid levels. Keeping uric

acid levels low by taking Adenuric\* once every day stops crystals building up, and over time it reduces symptoms. Keeping uric acid levels sufficiently low for a long period

In this leaflet

2. Before you take Adenuric

How to take Adenuric<sup>®</sup>
 Possible side effects

5. How to store Adenuric

6. Further information

. What Adenuric® is and what it is used for

abnormalities
- If you are being treated for high uric acid levels as a result of Lesch-Nyhan syndrome (a rare inherited condition in which there is too much uric acid in the blood)

If you have thyroid problems

- If you have thyroid problems
Should you experience allergie reactions to febuxostat, stop taking this medicine (see also section 4).
Possible symptoms of allergie reactions might be:
- rash including severe forms (e.g. blisters, nodules, itchye-etfoliative rash), itchiness
- swelling of limbs or face
- swelling of limbs or face
- difficulties in breathing
- fever with enlarged lymph nodes
- fever with enlarged lymph nodes
- fever with enlarged lymph nodes

Adenuric® 120 mg tablets is also used to treat and prevent high blood levels of uric acid that may occur when you start

When chemotherapy is given, cancer cells are destroyed, and uric acid levels increase in the blood accordingly, un-

not take Adenuric\*:
rou are allergic (hypersensitive) to febuxostat, or any of
other ingredients of this medicine (listed in section 6).

Take special care with Adenurie\*:
Tell your doctor before you start to take this medicine:
- If you have or have had feart failure or heart problems.
- If you have or have had renal disease and/or serious allergic reaction to Allopurinol (a medication used for the treat-

ment of Gout)

- If you have or have had liver disease or liver function test

eceive chemotherapy for blood cancers.

less the formation of uric acid is prevented.

2. BEFORE YOU TAKE ADENURIC

Adenuric® is indicated in adults.

FOR Adequire Containing the active substance februosstat, is a santhine oxidase (XO) inhibitor indicated for the chronic management of hyperuricemia in patients with gout. Ademure 'tablets are used to treat gout, which is associated and the contained that the

can also shrink tophi

cardiac and circulatory arrest. Your doctor might decide to permanently stop treatment with Adenuric.

with Adeuruic\*.

There have been rare reports of potentially life-threatening skin rashes (Stevens-Johnson Syndrome) with the use of Tebucostat, appearing initially a reddish integrétilie spots or may also include ulcers in the mouth, throat, nose, gentials and conjunctivities (red and swollen eyes). The rash may progress to widespread blistering or pecified of the skin. If you have developed Stevens-Johnson Syndrome with the

use of febuxostat, you must not be restarted on febuxosta at any time. If you developed a rash or these skin symp-toms, seek immediate advice from a doctor and tell that you are taking this medicine. If you are having a gout attack at the moment (a sudden

nset of severe pain, tenderness, redness, warmth and swelling in a joint), wait for the gout attack to subside be-fore first starting treatment with Adenuric\*. For some people, gout attacks may flare up when starting certain medicines that control uric acid levels. Not everyone gets flares, but you could get a flare-up even if you are taking Adenuric® and especially during the first weeks or nonths of treatment. It is important to keep taking Ade-nuric\* even if you have a flare, as Adenuric\* is still workng to lower uric acid. Over time, gout flares will occur les and be less painful if you keep taking Adenuric

every day. Your doctor will often prescribe other medicines, if they are

Your doctor will often prescribe other medicines, if they are needed, to help prevent or treat the symptoms of flares (such as pain and swelling in a joint). In patients with very high artate levels (e.g. those undergo-intermedicines) and the state of the state of

Children and adolescents:
Do not give this medicine to children under the age of 18 because the safety and efficacy have not been established.

Taking other medicines: Taking other medicines:

Please tell your dector or pharmacist if you are taking, or have recently taken, any other medicines, including medicines are recently taken, and the properties of the

Azathionrine (used to reduce immune response)

Theophylline (used to treat asthma)

Pregnancy and breastfeeding: It is not known if Adenuric \* may harm your unborn child. If you miss a dose of Adenuric<sup>®</sup> take it as soon as you re member unless it is almost time for your next dose, in

Tell your doctor if you think you are pregnant or if you are planning to become pregnant as Adenuric should not be used during pregnancy. It is not known if Adenuric may in breast milk. You should not use Adenur if you are breastfeeding, or if you are planning to breast-

Driving and using machines: Be aware that you may experience dizziness, sleepiness, blurred vision and numbness or tingling sensation during treatment and should not drive or operate machines if affected.

Important information about some of the ingredients of

Adenuric\*: Adenuric\* tablets contain lactose (a type of sugar). If you have been told that you have an intolerance to some sugars contact your doctor before taking this medicine.

3. HOW TO TAKE ADENURIC®

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not

The usual dose is one tablet daily. The tablets should be taken by mouth and can be taken with or without food

Adenuric® is available as either an 80 mg tablet or a 120 mg tablet. Your doctor will have prescribed the strength most

Continue to take Adenuric® every day even when you are not experiencing gout flare or attack.

. Prevention and treatment of high uric acid levels in pa-\*Trevention and treatment of min the act levels in pa-tients undergoing cancer chemotherapy: Adenuric\* is available as a 120 mg tablet. Adenuric\* who days before chemotherapy and continue its use accreding to your doctor's advice. Usually treatment is short-term.

If you take more Adenuric\* than you should:

In the event of an accidental overdose ask your doctor what to do, or contact your nearest accident and emergency department.

If you forget to take Adenuric®:

which case miss out the forgotten dose and take your next dose at the normal time. Do not take a double dose to make up for a forgotten dose.

up to a rogomet most.

If you stop taking Adenurie\*:

Do not stop paking Adenurie\*:

up to the the tree of your doctor even if you feel better. If you stop taking Adenurie\*

to the tree of your doctor even in your doctor even in your feel better. If you stop taking Adenurie\*

on your doctor in you want you want

4. POSSIBLE SIDE EFFECTS

4. POSSIBLE SIDE EFFECTS
Like all medicines, Adenuric\* can cause side effects, although not everybody gets them.

Stop taking this medicine and contact your doctor immediately fill the state of the st ately if the following rare (may affect up to 1 in 1,000 peo-ple) side effects occur, because a serious allergic reaction might follow

- anaphylactic reactions, drug hypersensitivity (see also section 2)

- potentially life-threatening skin rashes characterized by formation of blisters and shedding of the skin and inner su

formation of bisiers and shedding of the skin and miner suffices of body excitive, e.g. mouth and genitals, painful ulfaces of body excitive, e.g. mouth and genitals, painful ulfaces of body excitive, e.g. mouth and genitals, painful ulfaces, one throat and fatigue (Slevens-Johnson Syndrome,
foxer, one throat and fatigue (Slevens-Johnson Syndrome,
foxer body for the state of the state o

Common side effects (may affect up to 1 in 10 people) are: abnormal liver test results, diarrhea, headache, ras sea, increase in gout symptoms and localized swelling due to retention of fluids in tissues (oedema).

Uncommon side effects (may affect up to 1 in 100 people)

are:
- decreased appetite, change in blood sugar levels (diabetes)
of which a symptom may be excessive thirst, increased
blood fat levels, weight increase

difficulty in sleening sleeniness

dizziness, numbness, tingling, reduced or altered sensation (hypoesthesia, hemiparesis or paraesthesia), altered or reduced sense of taste

abnormal ECG heart tracing, irregular or rapid heartbeats.

 automina ECO final tracting, irregular of rapin neartoeats, feeling your heart beat (palpitation)
 hot flushes or flushing (e.g. redness of the face or neck), increased blood pressure, bleeding (hemorrhage, seen only in patients taking chemotherapy for blood disorders, cough, shortness of breath, chest discomfort or pain, inflammation of nasal passage and/or throat (upper respira-tory tract infection), bronchitis

dry mouth, abdominal pain/discomfort, heartburn/indiges tion, constipation, more frequent passing of stools, vomit

ing, stomach discomfort
- itching, hives, skin inflammation or discolouration, small -tiching, hives, skin inflammation or discolouration, small red or purple spot on the skin, small, flat red spots on the skin, flat, red area on the skin that is covered with small confluent bumps, rash, areas of redness and spots on the skin, other type of skin conditions
 -muscle cramp, muscle weakness, pain/ache in muscles/joints, bursitis or arthritis (inflammation of joints

muscles/joints, bursitis or arthritis (inflammation of joints usually accompanied by pain, swelling and/or stiffness), pain in extremity, back pain, muscle spasm - blood in the urine, abnormal frequent urination, abnormal urine tests (increased level of proteins in the urine), a reduction in the ability of the kidneys to function properly

fatigue, chest pain, chest discomfort stones in the gallbladder or in bile ducts (cholelithiasis)

Increase in blood thyroid stimulating hormone (TSH) changes in blood chemistry or amount of blood cells or

- changes in broad chemistry of annot platelets (abnormal blood test results) - kidney stones - erectile difficulties

erectife difficulties

Rars side effects (may affect up to 1 in 1,000 people) are:
muscle damage, a condition which on rare occasions can
serious. Il may cause muscle problems and particularly,
stream of the problems and particularly,
ture it may be caused by an abnormal muscle breakdown.
Contact your decir immediately if you experience muscle
pain, tendemess or weakness
re layers of the skin, especially
around the lips, cycs, genitals, hands, feet or tongue, with
possible sudden difficult breathing
high lever in combination with measles-like skin rash, enlarged (ymph nodes, liver enlargement, bepatits up to liver
to skin, with or without cosinaphilia) in the blood (cluskey
tosis, with or without cosinaphilia) in the blood (cluskey)

tosis, with or without eosinophilia)
- reddening of the skin (erythema), rash in various types

(e.g. itchy, with white spots, with blisters, with blisters conaining pus, with shedding of the skin, measles-like rash), widespread erythema, necrosis, and bullous detachment of the epidermis and mucous membranes, resulting in exfoliation and possible sensis (Stevens-Johnson Syndrome/Toxic epidermal necrolysis)

feeling thirst

or pharmacist

ringing in the ears blurred vision, change in vision

- mouth ulceration inflammation of the pancreas: common symptoms are abdominal pain, nausea and vomiting

increased sweating weight decrease, increased appetite, uncontrolled loss of

muscle and/or joint stiffness

abnormally low blood cell counts (white or red blood cells

urgent need to urinate

ease in urine amount due to inflammation

- changes or decrease in urine amount in the kidneys (tubulointerstitial neph - inflammation of the liver (hepatitis) - yellowing of the skin (jaundice) - liver damage

5 HOW TO STORE ADENUBIC®

5. FURTHER INFORMATION

The other ingredients are:

If any of the side effects gets serious, or if you notice any

5. HOW TO STORE ADENURIC\*
Keep out of reach and sight of children.
Do not store above 30°C.
Do not use after the expiry date which is stated on the carton and the tablet blister foil. The expiry date refers to the last day of that month.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

What Adenurie® contains
The active substance is febuxostat.
Each tablet contains 80 mg or 120 mg of febuxostat.

ines, their benefits and risks.

Do not by yourself interrupt the period of treatment preside effects not listed in this leaflet, please tell your doctor

Do not repeat the same prescription without consulting

Tablet core: lactose monohydrate, microcrystalline cellu-

lose, magnesium stearate, hydroxypropylcellulose, croscarmellose sodium and colloidal hydrated silica.

Film-coating: Onadry II vellow, 85F42129 containing

polyvinyl alcohol, titanium dioxide (E171), macrogols 3350, talc and iron oxide vellow (E172).

pack). Not all presentations may be marketed.

What Adenurie\* looks like and contents of the pack
Adenurie\* limit coated tablets are pale yellow to yellow in
Olour aid capable shaped.
The 80 with a fine-coated tablets are marked on one side with
80, win a fine-coated tablets are marked on one side with
80, win a fine-coated tablets are marked on one side with
120.
Adenurie\* is a plantial of one side with 120.
Statemark of the statemark of the side of the si

This is a medicament

Medicament is a product which affects your health and its consumption contrary to instructions is dangerous for

Follow strictly the doctor's prescription, the method of ise and the instructions of the pharmacist who sold the

nedicament.
The doctor and the pharmacist are the experts in medi-

- Keep all medicaments out of reach of children. Council of Arab Health Ministers Union of Arab Pharmacists

Manufactured by Patheon, France. 40 Boulevard De Champaret, 38300 Bourgoin Jallieu,

Packaged by ALGÖRITHM S.A.L. Zouk Mosbeh, Lebanon. Under License from Teijin Pharma Limited, Tokyo, Japan. \*Registered trademark of Teijin Limited, Tokyo, Japan.

PREPARED BY ALGORITHM		CC-15-01-003 & CC-15-01-004		
PRI Version # Final	PRINTED@ Luna Press # Final		addition of TLS indications scored Tablets	
Font : Times New Roman 7.7 pt			300160	ide dis
Color: Black Dimensions: 135 X 560 mm	UNFOLDED	Specs: W/F p	aper 43-47	g/m <sup>,</sup>
Δ	pproved by /dat	<u>e</u>		
TEIJIN:				
Supply Chain:				
Prod-TS/Prod Dev.:				
Marketing:				
Regulatory:				
Quality Assurance:				