

# Dexaméthasone MediS 4 mg/1ml

## Dexamethasone Phosphate

### FORM AND PRESENTATION :

Dexamethasone MediS 4 mg/1ml - injectable solution - Box of 50 ampoules.

### COMPOSITION :

	Dexaméthasone MédiS
Dexaméthasone phosphate Sodique	4,37 mg
methyl P-hydroxybenzoate	1,5 mg
propyl P-hydroxybenzoate	0,2 mg
Excipients Disodium edetate, Sodium citrate tribasique dihydrate, creatinine q.s	
W.F.I. q.s.f	1 ml

### PHARMACOLOGICAL PROPERTIES :

Synthesis steroïdal used for its anti-inflammatory action.

### THERAPEUTIC INDICATIONS :

Those of the corticotherapy by systemic or local route when an intense and fast corticotherapy is required.

### CONTRAINDICATIONS :

Hypersensitivity to one of the components of this drug, administration of live virus vaccine, mycosic or infectious states not controlled with a treatment, drop, evolutive peptic ulcer, psychotic states, cirrhosis, alcoholic, acute virus A, B, C hepatitis, thrombopenic purpura (I.M route).

No absolute contraindication when glucocorticoïd is considered life-threatening to the patient.

### CAUTION AND PRECAUTIONS OF USE:

- It's advised to diminish the salt in a diet which should be essentially rich in proteine and calcium, and weak in glucose.
- Use with care in children , menopansel wemen, ulcer or psychiatric antecedents, hypertension, diabete, osteoporosis, epilepsy.
- Patients receiving glucocorticoids should be monitored closely for the apparence of secondary bacterial or viral infectios.
- The chronic use of glucocorticoide abolish the hypothalamic-pituitary axis. Than an adrenocortical insufficiency is frequently observed when the drug is suddenly stopped.

**Pregnancy :** With human species, the retrospective studies have not detected any malformative risk linked to the taking of corticoids during the first trimester. A neonatal surrenalian insufficiency has been observed after a high-dose corticotherapy.

it is preferable to keep off nursing on treatment because the

corticoids pass in the maternal milk.

### **MEDICINAL INTERACTIONS :**

You should inform your doctor about the drugs that you take and especially : cardiologic drugs, drugs decreasing the blood potassium, aspirin, anticoagulant, anti diabetic, Phenobarbital, phenitoïn, primidone or rifamycin.

Let know your doctor if you are in period of de vaccination.

### **ADVERSE EFFECTS :**

Those of the corticotherapy, included water and sodium retention, hypertension, congestive cardiac insufficiency, osteoporosis, peptic ulcer, delay of cicatrization, aggravation of diabetic. Menstrual irregularities, decrease of the resistance of infection, increase of weight and glaucoma.

Let know your doctor about any unusual sensation.

### **DOSAGE AND MODE OF ADMINISTRATION :**

*Mode of administration* : intravenous or intramuscular route, local injection (intra-articular, soft tissue ).

*Dosage* : To conform to the medical prescription

### **DELIVERY CONDITIONS :**

List I, table A, delivered strictly on medical prescription

### **CONSERVATION :**

Keep away from light, heat and cold.

### **PRESERVATION :**

Dexamethasone MediS Box of 50 ampoules of 1 ml 923 313 2 H

### **Marketing Authorisation Holder and Manufacturer:**

LES LABORATOIRES MEDIS

Tunis road - KM 7 - BP 206 - 8000 Nabeul - Tunisia

Tel : (216) 72 23 50 06. Fax: (216) 72 23 50 16.

E-mail : [contact@labomedis.com](mailto:contact@labomedis.com)

**Date of the last revision : 09/2019**

#### **This is a drug**

- A drug is a specific product agent.
- A drug is a product acting on your health and its use, contrary to prescriptions may be dangerous for you.
- Strictly respect the doctor's prescription and the instructions of use he has prescribed.
- Follow your pharmacist's know this drug ; its indications and contra-indications.
- Do not discontinue the drug intake by yourself during the prescription period.
- Do not repeat the prescription or increase the dosage without consulting your doctor.

**KEEP ANY DRUG OUT OF THE REACH OF CHILDREN.**