

Sediproct

Suppositories

150225030016

Composition

Each suppository contains

Active Ingredients

Cinchocaine HCl	5 mg
Hydrocortisone	5 mg
Framycetine sulphate	10 mg
Esculoside	10 mg

Excipients

- Hard fats
- Sorbitan monooleate

Indications

The reduction of swelling, pain and hemorrhoids and other rectal lesions. The management of acute and chronic nonspecific proctitis, acute internal hemorrhoids, cryptitis, fissures and incomplete fistulas, internal and external pruritus ani. May be used in preoperative and postoperative hemorrhoidectomy and repair of fissures.

Contraindications, Warnings

Sensitivity to any of the excipients.

Hydrocortisone must not be used in the presence of tuberculosis, fungal or viral infections.

In pregnant animals, administration of corticosteroids can cause abnormalities of fetal development. The relevance of this finding to human beings has not been established. However, topical steroids should not be used extensively in pregnancy especially large amounts or for long periods.

As with all preparations containing corticosteroids, the possibility of systemic absorption should be considered. In particular, long-term continuous therapy should be avoided in infants.

Although rare, adrenal suppression may occur even with local steroids.

Precautions

Discontinue if sensitization occurs. Hydrocortisone should not be used until an adequate proctologic examination is completed and a diagnosis made. Other specific measures against infections, allergy, and other casual factors must not be neglected.

The possibility, however rare, that prolonged use of this preparation might produce systemic corticosteroid effects, should be borne in mind. Patients should be advised to inform subsequent physicians of previous use of hydrocortisone.

Pregnancy

The safe use of topical during pregnancy has not been fully established. Therefore, during pregnancy, they should not be used unnecessarily on extended areas, in large amounts or for prolonged periods of time.

Adverse Effects

Certain patients may experience burning upon application, especially if the mucous membrane is not intact.

Dosage

1 suppository morning and another one evening and after each bowel movement.

Presentation

Carton box containing 6 suppositories in aluminium strip + pamphlet.

Manufactured by

Medical Union Pharmaceuticals,
Abu-Sultan, Ismailia, Egypt.

Issue Date: 7 / 1 / 2001, Revision Date: --/--/2013

